

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

**(19) World Intellectual Property Organization
International Bureau**



(43) International Publication Date
30 January 2003 (30.01.2003)

PCT

(10) International Publication Number
WO 03/008537 A2

(51) International Patent Classification⁷: **C12N**

(72) **Inventors:** **SIMARD, John, J.L.**; 11918 Laughton Way, Northridge, CA 91326 (US). **DIAMOND, David, C.**; 23135 Schoenborn Street, West Hills, CA 91304 (US). **LIU, Liping**; 22228 Vicotry Blvd., H-111, Woodland Hills, CA 91367 (US). **XIE, Zhidong**; 22228 Victory Blvd., H-111, Woodland Hills, CA 91367 (US).

(21) International Application Number: PCT/US02/10189

(22) International Filing Date: 29 March 2002 (29.03.2002)

(25) Filing Language: English

(74) Agent: TAHMASSEBI, Sam; Knobbe, Martens, Olson & Bear, LLP, 16th Floor, 620 Newport Center Drive, Newport Beach, CA 92660 (US).

(26) Publication Language: English

(81) Designated State (national): AU.






(30) Priority Data:		
60/282,211	6 April 2001 (06.04.2001)	US
60/337,017	7 November 2001 (07.11.2001)	US
60/363,210	7 March 2002 (07.03.2002)	US

Published:
— *without international search report and to be republished upon receipt of that report*

(71) Applicant: CTL IMMUNOTHERAPIES CORP.
[US/US]; 9340 DeSoto Avenue, Chatsworth, CA 91311
(US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: EPITOPE SEQUENCES

[illegible]

Pool sequencing of PSMA_281-310 digested for 60 min by Proteasome

(57) Abstract: Disclosed herein are polypeptides, including epitopes, clusters, and antigens. Also disclosed are compositions including said polypeptides and methods for their use.

WO 03/008537 A2

EPITOPE SEQUENCES

Field of the Invention

5 The present invention generally relates to peptides, and nucleic acids encoding peptides, that are useful epitopes of target-associated antigens. More specifically, the invention relates to epitopes that have a high affinity for MHC class I and that are produced by target-specific proteasomes.

Description of the Related Art

Neoplasia and the Immune System

10 The neoplastic disease state commonly known as cancer is thought to result generally from a single cell growing out of control. The uncontrolled growth state typically results from a multi-step process in which a series of cellular systems fail, resulting in the genesis of a neoplastic cell. The resulting neoplastic cell rapidly reproduces itself, forms one or more tumors, and eventually may cause the death of the host.

15 Because the progenitor of the neoplastic cell shares the host's genetic material, neoplastic cells are largely unassailed by the host's immune system. During immune surveillance, the process in which the host's immune system surveys and localizes foreign materials, a neoplastic cell will appear to the host's immune surveillance machinery as a "self" cell.

Viruses and the Immune System

20 In contrast to cancer cells, virus infection involves the expression of clearly non-self antigens. As a result, many virus infections are successfully dealt with by the immune system with minimal clinical sequela. Moreover, it has been possible to develop effective vaccines for many of those infections that do cause serious disease. A variety of vaccine approaches have been used successfully to combat various diseases. These approaches include subunit vaccines consisting of individual proteins produced through recombinant DNA technology. Notwithstanding these
25 advances, the selection and effective administration of minimal epitopes for use as viral vaccines has remained problematic.

In addition to the difficulties involved in epitope selection stands the problem of viruses that have evolved the capability of evading a host's immune system. Many viruses, especially viruses that establish persistent infections, such as members of the herpes and pox virus families,
30 produce immunomodulatory molecules that permit the virus to evade the host's immune system. The effects of these immunomodulatory molecules on antigen presentation may be overcome by the targeting of select epitopes for administration as immunogenic compositions. To better understand the interaction of neoplastic cells and virally infected cells with the host's immune system, a discussion of the system's components follows below.

35 The immune system functions to discriminate molecules endogenous to an organism ("self" molecules) from material exogenous or foreign to the organism ("non-self" molecules). The

immune system has two types of adaptive responses to foreign bodies based on the components that mediate the response: a humoral response and a cell-mediated response. The humoral response is mediated by antibodies, while the cell-mediated response involves cells classified as lymphocytes. Recent anticancer and antiviral strategies have focused on mobilizing the host immune system as a means of anticancer or antiviral treatment or therapy.

The immune system functions in three phases to protect the host from foreign bodies: the cognitive phase, the activation phase, and the effector phase. In the cognitive phase, the immune system recognizes and signals the presence of a foreign antigen or invader in the body. The foreign antigen can be, for example, a cell surface marker from a neoplastic cell or a viral protein. Once the system is aware of an invading body, antigen specific cells of the immune system proliferate and differentiate in response to the invader-triggered signals. The last stage is the effector stage in which the effector cells of the immune system respond to and neutralize the detected invader.

An array of effector cells implements an immune response to an invader. One type of effector cell, the B cell, generates antibodies targeted against foreign antigens encountered by the host. In combination with the complement system, antibodies direct the destruction of cells or organisms bearing the targeted antigen. Another type of effector cell is the natural killer cell (NK cell), a type of lymphocyte having the capacity to spontaneously recognize and destroy a variety of virus infected cells as well as malignant cell types. The method used by NK cells to recognize target cells is poorly understood.

Another type of effector cell, the T cell, has members classified into three subcategories, each playing a different role in the immune response. Helper T cells secrete cytokines which stimulate the proliferation of other cells necessary for mounting an effective immune response, while suppressor T cells down-regulate the immune response. A third category of T cell, the cytotoxic T cell (CTL), is capable of directly lysing a targeted cell presenting a foreign antigen on its surface.

The Major Histocompatibility Complex and T Cell Target Recognition

T cells are antigen-specific immune cells that function in response to specific antigen signals. B lymphocytes and the antibodies they produce are also antigen-specific entities. However, unlike B lymphocytes, T cells do not respond to antigens in a free or soluble form. For a T cell to respond to an antigen, it requires the antigen to be processed to peptides which are then bound to a presenting structure encoded in the major histocompatibility complex (MHC). This requirement is called "MHC restriction" and it is the mechanism by which T cells differentiate "self" from "non-self" cells. If an antigen is not displayed by a recognizable MHC molecule, the T cell will not recognize and act on the antigen signal. T cells specific for a peptide bound to a recognizable MHC molecule bind to these MHC-peptide complexes and proceed to the next stages of the immune response.

There are two types of MHC, class I MHC and class II MHC. T Helper cells (CD4⁺) predominately interact with class II MHC proteins while cytolytic T cells (CD8⁺) predominately

interact with class I MHC proteins. Both classes of MHC protein are transmembrane proteins with a majority of their structure on the external surface of the cell. Additionally, both classes of MHC proteins have a peptide binding cleft on their external portions. It is in this cleft that small fragments of proteins, endogenous or foreign, are bound and presented to the extracellular environment.

5 Cells called "professional antigen presenting cells" (pAPCs) display antigens to T cells using the MHC proteins but additionally express various co-stimulatory molecules depending on the particular state of differentiation/activation of the pAPC. When T cells, specific for the peptide bound to a recognizable MHC protein, bind to these MHC-peptide complexes on pAPCs, the specific co-stimulatory molecules that act upon the T cell direct the path of differentiation/activation taken by the
10 T cell. That is, the co-stimulation molecules affect how the T cell will act on antigenic signals in future encounters as it proceeds to the next stages of the immune response.

As discussed above, neoplastic cells are largely ignored by the immune system. A great deal of effort is now being expended in an attempt to harness a host's immune system to aid in combating the presence of neoplastic cells in a host. One such area of research involves the formulation of
15 anticancer vaccines.

Anticancer Vaccines

Among the various weapons available to an oncologist in the battle against cancer is the immune system of the patient. Work has been done in various attempts to cause the immune system to combat cancer or neoplastic diseases. Unfortunately, the results to date have been
20 largely disappointing. One area of particular interest involves the generation and use of anticancer vaccines.

To generate a vaccine or other immunogenic composition, it is necessary to introduce to a subject an antigen or epitope against which an immune response may be mounted. Although neoplastic cells are derived from and therefore are substantially identical to normal cells on a genetic level, many neoplastic cells are known to present tumor-associated antigens (TuAAs). In
25 theory, these antigens could be used by a subject's immune system to recognize these antigens and attack the neoplastic cells. In reality, however, neoplastic cells generally appear to be ignored by the host's immune system.

A number of different strategies have been developed in an attempt to generate vaccines with activity against neoplastic cells. These strategies include the use of tumor-associated antigens as immunogens. For example, U.S. Patent No. 5,993,828, describes a method for producing an immune response against a particular subunit of the Urinary Tumor Associated Antigen by administering to a subject an effective dose of a composition comprising inactivated tumor cells having the Urinary Tumor Associated Antigen on the cell surface and at least one tumor associated
35 antigen selected from the group consisting of GM-2, GD-2, Fetal Antigen and Melanoma

Associated Antigen. Accordingly, this patent describes using whole, inactivated tumor cells as the immunogen in an anticancer vaccine.

Another strategy used with anticancer vaccines involves administering a composition containing isolated tumor antigens. In one approach, MAGE-A1 antigenic peptides were used as an immunogen. (See Chaux, P., *et al.*, "Identification of Five MAGE-A1 Epitopes Recognized by Cytolytic T Lymphocytes Obtained by *In Vitro* Stimulation with Dendritic Cells Transduced with MAGE-A1," J. Immunol., 163(5):2928-2936 (1999)). There have been several therapeutic trials using MAGE-A1 peptides for vaccination, although the effectiveness of the vaccination regimes was limited. The results of some of these trials are discussed in Vose, J.M., "Tumor Antigens Recognized by T Lymphocytes," 10th European Cancer Conference, Day 2, Sept. 14, 1999.

In another example of tumor associated antigens used as vaccines, Scheinberg, *et al.* treated 12 chronic myelogenous leukemia (CML) patients already receiving interferon (IFN) or hydroxyurea with 5 injections of class I-associated bcr-abl peptides with a helper peptide plus the adjuvant QS-21. Scheinberg, D.A., *et al.*, "BCR-ABL Breakpoint Derived Oncogene Fusion Peptide Vaccines Generate Specific Immune Responses in Patients with Chronic Myelogenous Leukemia (CML) [Abstract 1665], American Society of Clinical Oncology 35th Annual Meeting, Atlanta (1999). Proliferative and delayed type hypersensitivity (DTH) T cell responses indicative of T-helper activity were elicited, but no cytolytic killer T cell activity was observed within the fresh blood samples.

Additional examples of attempts to identify TuAAs for use as vaccines are seen in the recent work of Cebon, *et al.* and Scheibenbogen, *et al.* Cebon, *et al.* immunized patients with metastatic melanoma using intradermally administered MART-1₂₆₋₃₅ peptide with IL-12 in increasing doses given either subcutaneously or intravenously. Of the first 15 patients, 1 complete remission, 1 partial remission, and 1 mixed response were noted. Immune assays for T cell generation included DTH, which was seen in patients with or without IL-12. Positive CTL assays were seen in patients with evidence of clinical benefit, but not in patients without tumor regression. Cebon, *et al.*, "Phase I Studies of Immunization with Melan-A and IL-12 in HLA A2+ Positive Patients with Stage III and IV Malignant Melanoma," [Abstract 1671], American Society of Clinical Oncology 35th Annual Meeting, Atlanta (1999).

Scheibenbogen, *et al.* immunized 18 patients with 4 HLA class I restricted tyrosinase peptides, 16 with metastatic melanoma and 2 adjuvant patients. Scheibenbogen, *et al.*, "Vaccination with Tyrosinase peptides and GM-CSF in Metastatic Melanoma: a Phase II Trial," [Abstract 1680], American Society of Clinical Oncology 35th Annual Meeting, Atlanta (1999). Increased CTL activity was observed in 4/15 patients, 2 adjuvant patients, and 2 patients with evidence of tumor regression. As in the trial by Cebon, *et al.*, patients with progressive disease did

not show boosted immunity. In spite of the various efforts expended to date to generate efficacious anticancer vaccines, no such composition has yet been developed.

Antiviral Vaccines

5 Vaccine strategies to protect against viral diseases have had many successes. Perhaps the most notable of these is the progress that has been made against the disease small pox, which has been driven to extinction. The success of the polio vaccine is of a similar magnitude.

Viral vaccines can be grouped into three classifications: live attenuated virus vaccines, such as vaccinia for small pox, the Sabin poliovirus vaccine, and measles mumps and rubella; whole killed or inactivated virus vaccines, such as the Salk poliovirus vaccine, hepatitis A virus
10 vaccine and the typical influenza virus vaccines; and subunit vaccines, such as hepatitis B. Due to their lack of a complete viral genome, subunit vaccines offer a greater degree of safety than those based on whole viruses.

The paradigm of a successful subunit vaccine is the recombinant hepatitis B vaccine based on the viruses envelope protein. Despite much academic interest in pushing the reductionist
15 subunit concept beyond single proteins to individual epitopes, the efforts have yet to bear much fruit. Viral vaccine research has also concentrated on the induction of an antibody response although cellular responses also occur. However, many of the subunit formulations are particularly poor at generating a CTL response.

Summary of the Invention

20 Previous methods of priming professional antigen presenting cells (pAPCs) to display target cell epitopes have relied simply on causing the pAPCs to express target-associated antigens (TAAs), or epitopes of those antigens which are thought to have a high affinity for MHC I molecules. However, the proteasomal processing of such antigens results in presentation of epitopes on the pAPC that do not correspond to the epitopes present on the target cells.

25 Using the knowledge that an effective cellular immune response requires that pAPCs present the same epitope that is presented by the target cells, the present invention provides epitopes that have a high affinity for MHC I, and that correspond to the processing specificity of the housekeeping proteasome, which is active in peripheral cells. These epitopes thus correspond to those presented on target cells. The use of such epitopes in vaccines can activate the cellular
30 immune response to recognize the correctly processed TAA and can result in removal of target cells that present such epitopes. In some embodiments, the housekeeping epitopes provided herein can be used in combination with immune epitopes, generating a cellular immune response that is competent to attack target cells both before and after interferon induction. In other embodiments the epitopes are useful in the diagnosis and monitoring of the target-associated disease and in the
35 generation of immunological reagents for such purposes.

Embodiments of the invention relate to isolated epitopes and antigens or polypeptides that comprise the epitopes. Preferred embodiments include an epitope or antigen having the sequence as disclosed in TABLE 1. Other embodiments can include an epitope cluster comprising a polypeptide from Table 1. Further, embodiments include a polypeptide having substantial similarity to the already mentioned epitopes, antigens, or clusters. Other preferred embodiments include a polypeptide having functional similarity to any of the above. Still further embodiments relate to a nucleic acid encoding the polypeptide of any of the epitopes, clusters, antigens, and polypeptides from Table 1 and mentioned herein.

The epitope can be immunologically active. The polypeptide comprising the epitope can be less than about 30 amino acids in length, more preferably, the polypeptide is 8 to 10 amino acids in length, for example. Substantial or functional similarity can include addition of at least one amino acid, for example, and the at least one additional amino acid can be at an N-terminus of the polypeptide. The substantial or functional similarity can include a substitution of at least one amino acid.

The epitope, cluster, or polypeptide comprising the same can have affinity to an HLA-A2 molecule. The affinity can be determined by an assay of binding, by an assay of restriction of epitope recognition, by a prediction algorithm, and the like. The epitope, cluster, or polypeptide comprising the same can have affinity to an HLA-B7, HLA-B51 molecule, and the like.

In preferred embodiments the polypeptide can be a housekeeping epitope. The epitope or polypeptide can correspond to an epitope displayed on a tumor cell, to an epitope displayed on a neovasculature cell, and the like. The epitope or polypeptide can be an immune epitope. The epitope, cluster and/or polypeptide can be a nucleic acid.

Other embodiments relate to pharmaceutical compositions comprising the polypeptides, including an epitope from Table 1, a cluster, or a polypeptide comprising the same and a pharmaceutically acceptable adjuvant, carrier, diluent, excipient, and the like. The adjuvant can be a polynucleotide. The polynucleotide can include a dinucleotide. The dinucleotide can be CpG, for example. The adjuvant can be encoded by a polynucleotide. The adjuvant can be a cytokine and the cytokine can be, for example, GM-CSF.

The pharmaceutical compositions can further include a professional antigen-presenting cell (pAPC). The pAPC can be a dendritic cell, for example. The pharmaceutical composition can further include a second epitope. The second epitope can be a polypeptide. The second epitope can be a nucleic acid. The second epitope can be a housekeeping epitope, an immune epitope, and the like.

Still further embodiments relate to pharmaceutical compositions that include any of the nucleic acids discussed herein, including those that encode polypeptides that comprise epitopes or

antigens from Table 1. Such compositions can include a pharmaceutically acceptable adjuvant, carrier, diluent, excipient, and the like.

Other embodiments relate to recombinant constructs that include such a nucleic acid as described herein, including those that encode polypeptides that comprise epitopes or antigens from Table 1. The constructs can further include a plasmid, a viral vector, an artificial chromosome, and the like. The construct can further include a sequence encoding at least one feature, such as for example, a second epitope, an IRES, an ISS, an NIS, ubiquitin.

Further embodiments relate to purified antibodies that specifically bind to at least one of the epitopes in Table 1A. Other embodiments relate to purified antibodies that specifically bind to a peptide-MHC protein complex comprising an epitope disclosed in Table 1A or any other suitable epitope. The antibody from any embodiment can be a monoclonal antibody.

Still other embodiments relate to multimeric MHC-peptide complexes that include an epitope, such as, for example, an epitope disclosed in Table 1.

Embodiments relate to isolated T cells expressing a T cell receptor specific for an MHC-peptide complex. The complex can include an epitope, such as, for example, an epitope disclosed in Table 1 of claim 1. The T cell can be produced by an *in vitro* immunization. The T cell can be isolated from an immunized animal. Embodiments relate to T cell clones, including cloned T cells, such as those discussed above. Embodiments also relate to polyclonal population of T cells. Such populations can include a T cell, as described above, for example.

Still further embodiments relate to pharmaceutical compositions that include a T cell, such as those described above, for example, and a pharmaceutically acceptable adjuvant, carrier, diluent, excipient, and the like.

Embodiments of the invention relate to isolated protein molecules comprising the binding domain of a T cell receptor specific for an MHC-peptide complex. The complex can include an epitope disclosed in Table 1. The protein can be multivalent. Other embodiments relate to isolated nucleic acids encoding such proteins. Still further embodiments relate to recombinant constructs that include such nucleic acids.

Other embodiments of the invention relate to host cells expressing the recombinant construct described herein, including constructs encoding an epitope, cluster or polypeptide comprising the same, disclosed in Table 1, for example. The host cell can be a dendritic cell, macrophage, tumor cell, tumor-derived cell, and the like. The host cell can be a bacterium, fungus, protozoan and the like. Embodiments also relate to pharmaceutical compositions that include a host cell, such as those discussed herein, and a pharmaceutically acceptable adjuvant, carrier, diluent, excipient, and the like.

Still other embodiments relate to vaccines or immunotherapeutic compositions that include at least one component, such as, for example, an epitope disclosed in Table 1 or otherwise

described herein; a cluster that includes such an epitope, an antigen or polypeptide that includes such an epitope; a composition described above and herein; a construct, a T cell, or a host cell as described above and herein.

Further embodiments relate to methods of treating an animal. The methods can include administering to an animal a vaccine or immunotherapeutic composition, including those disclosed above and herein. The administering step can include a mode of delivery, such as, for example, transdermal, intranodal, perinodal, oral, intravenous, intradermal, intramuscular, intraperitoneal, mucosal, aerosol inhalation, instillation, and the like. The method can further include a step of assaying to determine a characteristic indicative of a state of a target cell or target cells. The method can include a first assaying step and a second assaying step, wherein the first assaying step precedes the administering step, and wherein the second assaying step follows the administering step. The method can further include a step of comparing the characteristic determined in the first assaying step with the characteristic determined in the second assaying step to obtain a result. The result can be for example, evidence of an immune response, a diminution in number of target cells, a loss of mass or size of a tumor comprising target cells, a decrease in number or concentration of an intracellular parasite infecting target cells, and the like.

Embodiments relate to methods of evaluating immunogenicity of a vaccine or immunotherapeutic composition. The methods can include administering to an animal a vaccine or immunotherapeutic, such as those described above and elsewhere herein, and evaluating immunogenicity based on a characteristic of the animal. The animal can be HLA-transgenic.

Other embodiments relate to methods of evaluating immunogenicity that include *in vitro* stimulation of a T cell with the vaccine or immunotherapeutic composition, such as those described above and elsewhere herein, and evaluating immunogenicity based on a characteristic of the T cell. The stimulation can be a primary stimulation.

Still further embodiments relate to methods of making a passive/adoptive immunotherapeutic. The methods can include combining a T cell or a host cell, such as those described above and elsewhere herein, with a pharmaceutically acceptable adjuvant, carrier, diluent, excipient, and the like.

Other embodiments relate to methods of determining specific T cell frequency, and can include the step of contacting T cells with a MHC-peptide complex comprising an epitope disclosed in Table 1, or a complex comprising a cluster or antigen comprising such an epitope. The contacting step can include at least one feature, such as, for example, immunization, restimulation, detection, enumeration, and the like. The method can further include ELISPOT analysis, limiting dilution analysis, flow cytometry, *in situ* hybridization, the polymerase chain reaction, any combination thereof, and the like.

Embodiments relate to methods of evaluating immunologic response. The methods can include the above-described methods determining specific T cell frequency carried out prior to and subsequent to an immunization step.

5 Another embodiment relates to methods of evaluating immunologic response. The methods can include determining frequency, cytokine production, or cytolytic activity of T cells, prior to and subsequent to a step of stimulation with MHC-peptide complexes comprising an epitope, such as, for example an epitope from Table 1, a cluster or a polypeptide comprising such an epitope.

10 Further embodiments relate to methods of diagnosing a disease. The methods can include contacting a subject tissue with at least one component, including, for example, a T cell, a host cell, an antibody, a protein, including those described above and elsewhere herein; and diagnosing the disease based on a characteristic of the tissue or of the component. The contacting step can take place *in vivo*. The contacting step can take place *in vitro*.

15 Still other embodiments relate to methods of making a vaccine. The methods can include combining at least one component, an epitope, a composition, a construct, a T cell, a host cell; including any of those described above and elsewhere herein, with a pharmaceutically acceptable adjuvant, carrier, diluent, excipient, and the like.

20 Embodiments relate to computer readable media having recorded thereon the sequence of any one of SEQ ID NOS: 1 -602, in a machine having a hardware or software that calculates the physical, biochemical, immunologic, or molecular genetic properties of a molecule embodying said sequence.

25 Still other embodiments relate to methods of treating an animal. The methods can include combining the method of treating an animal that includes administering to the animal a vaccine or immunotherapeutic composition, such as described above and elsewhere herein, combined with at least one mode of treatment, including, for example, radiation therapy, chemotherapy, biochemotherapy, surgery, and the like.

Further embodiments relate to isolated polypeptides that include an epitope cluster from a target-associated antigen having the sequence as disclosed in any one of Tables 25-44, wherein the amino acid sequence includes not more than about 80% of the amino acid sequence of the antigen.

30 Other embodiments relate to vaccines or immunotherapeutic products that include an isolated peptide as described above and elsewhere herein. Still other embodiments relate to isolated polynucleotides encoding a polypeptide as described above and elsewhere herein. Other embodiments relate vaccines or immunotherapeutic products that include these polynucleotides. The polynucleotide can be DNA or RNA.

Brief Description of the Drawings

Figure 1 is a sequence alignment of NY-ESO-1 and several similar protein sequences.

Figure 2 graphically represents a plasmid vaccine backbone useful for delivering nucleic acid-encoded epitopes.

5 Figures 3A and 3B are FACS profiles showing results of HLA-A2 binding assays for tyrosinase₂₀₇₋₂₁₅ and tyrosinase₂₀₈₋₂₁₆.

Figure 4 is a T=120 min. time point mass spectrum of the fragments produced by proteasomal cleavage of SSX-2₃₁₋₆₈.

Figure 5 shows a binding curve for HLA-A2:SSX-2₄₁₋₄₉ with controls.

10 Figure 6 shows specific lysis of SSX-2₄₁₋₄₉-pulsed targets by CTL from SSX-2₄₁₋₄₉-immunized HLA-A2 transgenic mice.

Figure 7A, B, and C show results of N-terminal pool sequencing of a T=60 min. time point aliquot of the PSMA₁₆₃₋₁₉₂ proteasomal digest.

15 Figure 8 shows binding curves for HLA-A2:PSMA₁₆₈₋₁₇₇ and HLA-A2:PSMA₂₈₈₋₂₉₇ with controls.

Figure 9 shows results of N-terminal pool sequencing of a T=60 min. time point aliquot of the PSMA₂₈₁₋₃₁₀ proteasomal digest.

Figure 10 shows binding curves for HLA-A2:PSMA₄₆₁₋₄₆₉, HLA-A2:PSMA₄₆₀₋₄₆₉, and HLA-A2:PSMA₆₆₃₋₆₇₁, with controls.

20 Figure 11 shows the results of a γ -IFN-based ELISPOT assay detecting PSMA₄₆₃₋₄₇₁-reactive HLA-A1⁺ CD8⁺ T cells.

Figure 12 shows blocking of reactivity of the T cells used in figure 10 by anti-HLA-A1 mAb, demonstrating HLA-A1-restricted recognition.

Figure 13 shows a binding curve for HLA-A2:PSMA₆₆₃₋₆₇₁, with controls.

25 Figure 14 shows a binding curve for HLA-A2:PSMA₆₆₂₋₆₇₁, with controls.

Figure 15. Comparison of anti-peptide CTL responses following immunization with various doses of DNA by different routes of injection.

Figure 16. Growth of transplanted gp33 expressing tumor in mice immunized by i.ln. injection of gp33 epitope-expressing, or control, plasmid.

30 Figure 17. Amount of plasmid DNA detected by real-time PCR in injected or draining lymph nodes at various times after i.ln. of i.m. injection, respectively.

Detailed Description of the Preferred Embodiment

Definitions

35 Unless otherwise clear from the context of the use of a term herein, the following listed terms shall generally have the indicated meanings for purposes of this description.

PROFESSIONAL ANTIGEN-PRESENTING CELL (pAPC) – a cell that possesses T cell costimulatory molecules and is able to induce a T cell response. Well characterized pAPCs include dendritic cells, B cells, and macrophages.

PERIPHERAL CELL – a cell that is not a pAPC.

5 HOUSEKEEPING PROTEASOME – a proteasome normally active in peripheral cells, and generally not present or not strongly active in pAPCs.

IMMUNE PROTEASOME – a proteasome normally active in pAPCs; the immune proteasome is also active in some peripheral cells in infected tissues.

10 EPITOPE – a molecule or substance capable of stimulating an immune response. In preferred embodiments, epitopes according to this definition include but are not necessarily limited to a polypeptide and a nucleic acid encoding a polypeptide, wherein the polypeptide is capable of stimulating an immune response. In other preferred embodiments, epitopes according to this definition include but are not necessarily limited to peptides presented on the surface of cells, the peptides being non-covalently bound to the binding cleft of class I MHC, such that they can
15 interact with T cell receptors.

MHC EPITOPE – a polypeptide having a known or predicted binding affinity for a mammalian class I or class II major histocompatibility complex (MHC) molecule.

HOUSEKEEPING EPITOPE – In a preferred embodiment, a housekeeping epitope is defined as a polypeptide fragment that is an MHC epitope, and that is displayed on a cell in which
20 housekeeping proteasomes are predominantly active. In another preferred embodiment, a housekeeping epitope is defined as a polypeptide containing a housekeeping epitope according to the foregoing definition, that is flanked by one to several additional amino acids. In another preferred embodiment, a housekeeping epitope is defined as a nucleic acid that encodes a housekeeping epitope according to the foregoing definitions.

25 IMMUNE EPITOPE – In a preferred embodiment, an immune epitope is defined as a polypeptide fragment that is an MHC epitope, and that is displayed on a cell in which immune proteasomes are predominantly active. In another preferred embodiment, an immune epitope is defined as a polypeptide containing an immune epitope according to the foregoing definition, that is flanked by one to several additional amino acids. In another preferred embodiment, an immune
30 epitope is defined as a polypeptide including an epitope cluster sequence, having at least two polypeptide sequences having a known or predicted affinity for a class I MHC. In yet another preferred embodiment, an immune epitope is defined as a nucleic acid that encodes an immune epitope according to any of the foregoing definitions.

TARGET CELL – a cell to be targeted by the vaccines and methods of the invention.
35 Examples of target cells according to this definition include but are not necessarily limited to: a

neoplastic cell and a cell harboring an intracellular parasite, such as, for example, a virus, a bacterium, or a protozoan.

TARGET-ASSOCIATED ANTIGEN (TAA) – a protein or polypeptide present in a target cell.

5 TUMOR-ASSOCIATED ANTIGENS (TuAA) – a TAA, wherein the target cell is a neoplastic cell.

HLA EPITOPE – a polypeptide having a known or predicted binding affinity for a human class I or class II HLA complex molecule.

10 ANTIBODY – a natural immunoglobulin (Ig), poly- or monoclonal, or any molecule composed in whole or in part of an Ig binding domain, whether derived biochemically or by use of recombinant DNA. Examples include *inter alia*, F(ab), single chain Fv, and Ig variable region-phage coat protein fusions.

15 ENCODE – an open-ended term such that a nucleic acid encoding a particular amino acid sequence can consist of codons specifying that (poly)peptide, but can also comprise additional sequences either translatable, or for the control of transcription, translation, or replication, or to facilitate manipulation of some host nucleic acid construct.

20 SUBSTANTIAL SIMILARITY – this term is used to refer to sequences that differ from a reference sequence in an inconsequential way as judged by examination of the sequence. Nucleic acid sequences encoding the same amino acid sequence are substantially similar despite differences in degenerate positions or modest differences in length or composition of any non-coding regions. Amino acid sequences differing only by conservative substitution or minor length variations are substantially similar. Additionally, amino acid sequences comprising housekeeping epitopes that differ in the number of N-terminal flanking residues, or immune epitopes and epitope clusters that differ in the number of flanking residues at either terminus, are substantially similar. Nucleic acids
25 that encode substantially similar amino acid sequences are themselves also substantially similar.

30 FUNCTIONAL SIMILARITY – this term is used to refer to sequences that differ from a reference sequence in an inconsequential way as judged by examination of a biological or biochemical property, although the sequences may not be substantially similar. For example, two nucleic acids can be useful as hybridization probes for the same sequence but encode differing amino acid sequences. Two peptides that induce cross-reactive CTL responses are functionally similar even if they differ by non-conservative amino acid substitutions (and thus do not meet the substantial similarity definition). Pairs of antibodies, or TCRs, that recognize the same epitope can be functionally similar to each other despite whatever structural differences exist. In testing for functional similarity of immunogenicity one would generally immunize with the “altered” antigen
35 and test the ability of the elicited response (Ab, CTL, cytokine production, etc.) to recognize the target antigen. Accordingly, two sequences may be designed to differ in certain respects while

retaining the same function. Such designed sequence variants are among the embodiments of the present invention.

Table 1A. SEQ ID NOS.* including eptopes in Examples 1-7, 13.

SEQ ID NO	IDENTITY	SEQUENCE
1	Tyr 207-216	FLPWHRLFLL
2	Tyrosinase protein	Accession number**: P14679
3	SSX-2 protein	Accession number: NP_003138
4	PSMA protein	Accession number: NP_004467
5	Tyrosinase cDNA	Accession number: NM_000372
6	SSX-2 cDNA	Accession number: NM_003147
7	PSMA cDNA	Accession number: NM_004476
8	Tyr 207-215	FLPWHRLFL
9	Tyr 208-216	LPWHRLFLL
10	SSX-2 31-68	YFSKEEWEKMKASEKIFYVYMKRKYEAMTKLGFKATLP
11	SSX-2 32-40	FSKEEWEKM
12	SSX-2 39-47	KMKASEKIF
13	SSX-2 40-48	MKASEKIFY
14	SSX-2 39-48	KMKASEKIFY
15	SSX-2 41-49	KASEKIFYV
16	SSX-2 40-49	MKASEKIFYV
17	SSX-2 41-50	KASEKIFYVY
18	SSX-2 42-49	ASEKIFYVY
19	SSX-2 53-61	RKYEAMTKL
20	SSX-2 52-61	KRKYEAMTKL
21	SSX-2 54-63	KYEAMTKLGF
22	SSX-2 55-63	YEAMTKLGF
23	SSX-2 56-63	EAMTKLGF
24	HBV18-27	FLPSDYFPSV
25	HLA-B44 binder	AEMGKYSFY
26	SSX-1 41-49	KYSEKISYV
27	SSX-3 41-49	KVSEKIVYV
28	SSX-4 41-49	KSSEKIVYV
29	SSX-5 41-49	KASEKIIVY
30	PSMA163-192	AFSPQGMPEGDLVYVNYARTEDFFKLERDM
31	PSMA 168-190	GMPEGDLVYVNYARTEDFFKLER
32	PSMA 169-177	MPEGDLVYV
33	PSMA 168-177	GMPEGDLVYV
34	PSMA 168-176	GMPEGDLVY
35	PSMA 167-176	QGMPEGDLVY
36	PSMA 169-176	MPEGDLVY

37	PSMA 171-179	EGDLVYVNY
38	PSMA 170-179	PEGDLVYVNY
39	PSMA 174-183	LVYVNYARTE
40	PSMA 177-185	VNYARTEDF
41	PSMA 176-185	YVNYARTEDF
42	PSMA 178-186	NYARTEDFF
43	PSMA 179-186	YARTEDFF
44	PSMA 181-189	RTEDFFKLE
45	PSMA 281-310	RGIAEAVGLPSIPVHPIGYYDAQKLEKMG
46	PSMA 283-307	IAEAVGLPSIPVHPIGYYDAQKLE
47	PSMA 289-297	LPSIPVHPI
48	PSMA 288-297	GLPSIPVHPI
49	PSMA 297-305	IGYYDAQKL
50	PSMA 296-305	PIGYYDAQKL
51	PSMA 291-299	SIPVHPIGY
52	PSMA 290-299	PSIPVHPIGY
53	PSMA 292-299	IPVHPIGY
54	PSMA 299-307	YYDAQKLE
55	PSMA454-481	SSIEGNYTLRVDCTPLMYSLVHLTKEL
56	PSMA 456-464	IEGNYTLRV
57	PSMA 455-464	SIEGNYTLRV
58	PSMA 457-464	EGNYTLRV
59	PSMA 461-469	TLRVDCTPL
60	PSMA 460-469	YTLRVDCTPL
61	PSMA 462-470	LRVDCTPLM
62	PSMA 463-471	RVDCTPLMY
63	PSMA 462-471	LRVDCTPLMY
64	PSMA653-687	FDKSNPIVLRMMNDQLMFLERAFIDPLGLPDRPFY
65	PSMA 660-681	VLRMMNDQLMFLERAFIDPLGL
66	PSMA 663-671	MMNDQLMFL
67	PSMA 662-671	RMMNDQLMFL
68	PSMA 662-670	RMMNDQLMF
69	Tyr 1-17	MLLAVLYCLLWSFQTSA

Table 1B. SEQ ID NOS.* Including epitopes in Examples 14 and 15.

SEQ ID NO	IDENTITY	SEQUENCE
70	GP100 protein	** Accession number: P40967
71	MAGE-1 protein	Accession number: P43355
72	MAGE-2 protein	Accession number: P43356
73	MAGE-3 protein	Accession number: P43357
74	NY-ESO-1 protein	Accession number: P78358
75	LAGE-1a protein	Accession number: CAA11116

76	LAGE-1b protein	Accession number: CAA11117
77	PRAME protein	Accession number: NP_006106
78	PSA protein	Accession number: P07288
79	PSCA protein	Accession number: O43653
80	GP100 cds	Accession number: U20093
81	MAGE-1 cds	Accession number: M77481
82	MAGE-2 cds	Accession number: L18920
83	MAGE-3 cds	Accession number: U03735
84	NY-ESO-1 cDNA	Accession number: U87459
85	PRAME cDNA	Accession number: NM_006115
86	PSA cDNA	Accession number: NM_001648
87	PSCA cDNA	Accession number: AF043498
88	GP100 630-638	LPHSSSHWL
89	GP100 629-638	QLPHSSSHWL
90	GP100 614-622	LIYRRRLMK
91	GP100 613-622	SLIYRRRLMK
92	GP100 615-622	IYRRRLMK
93	GP100 630-638	LPHSSSHWL
94	GP100 629-638	QLPHSSSHWL
95	MAGE-1 95-102	ESLFRAVI
96	MAGE-1 93-102	ILESFRVI
97	MAGE-1 93-101	ILESFRV
98	MAGE-1 92-101	CILESFRV
99	MAGE-1 92-100	CILESFR
100	MAGE-1 263-271	EFLWGPRAL
101	MAGE-1 264-271	FLWGPRAL
102	MAGE-1 264-273	FLWGPRALAE
103	MAGE-1 265-274	LWGPRALAE
104	MAGE-1 268-276	PRALAETSY
105	MAGE-1 267-276	GPRALAETSY
106	MAGE-1 269-277	RALAETSYV
107	MAGE-1 271-279	LAETSYVKV
108	MAGE-1 270-279	ALAETSYVKV
109	MAGE-1 272-280	AETSYVKVL
110	MAGE-1 271-280	LAETSYVKVL
111	MAGE-1 274-282	TSYVKVLEY
112	MAGE-1 273-282	ETSYVKVLEY
113	MAGE-1 278-286	KVLEYVIKV
114	MAGE-1 168-177	SYVLVTCLGL
115	MAGE-1 169-177	YVLVTCLGL
116	MAGE-1 170-177	VLVTCLGL
117	MAGE-1 240-248	TQDLVQEKY

118	MAGE-1 239-248	LTQDLVQEKY
119	MAGE-1 232-240	YGEPRKLLT
120	MAGE-1 243-251	LVQEKYLEY
121	MAGE-1 242-251	DLVQEKYLEY
122	MAGE-1 230-238	SAYGEPRKL
123	MAGE-1 278-286	KVLEYVIKV
124	MAGE-1 277-286	VKVLEYVIKV
125	MAGE-1 276-284	YVKVLEYVI
126	MAGE-1 274-282	TSYVKVLEY
127	MAGE-1 273-282	ETSYVKVLEY
128	MAGE-1 283-291	VIKVSARVR
129	MAGE-1 282-291	YVIKVSARVR
130	MAGE-2 115-122	ELVHFLLL
131	MAGE-2 113-122	MVELVHFLLL
132	MAGE-2 109-116	ISRKMVEL
133	MAGE-2 108-116	AI SRKMVEL
134	MAGE-2 107-116	AAISRK MVEL
135	MAGE-2 112-120	KMVELVHFL
136	MAGE-2 109-117	ISRKMVELV
137	MAGE-2 108-117	AI SRKMVELV
138	MAGE-2 116-124	LVHFLLLKY
139	MAGE-2 115-124	ELVHFLLLKY
140	MAGE-2 111-119	RKMVELVHF
141	MAGE-2 158-166	LQLVFGIEV
142	MAGE-2 157-166	YLQLVFGIEV
143	MAGE-2 159-167	QLVFGIEVV
144	MAGE-2 158-167	LQLVFGIEVV
145	MAGE-2 164-172	IEVV E VPI
146	MAGE-2 163-172	GIEVV E VPI
147	MAGE-2 162-170	FGIEVV E V
148	MAGE-2 154-162	ASEYLQLVF
149	MAGE-2 153-162	KASEYLQLVF
150	MAGE-2 218-225	BEKIWEEL
151	MAGE-2 216-225	APEEKIWEEL
152	MAGE-2 216-223	APEEKIWE
153	MAGE-2 220-228	KIWEELSML
154	MAGE-2 219-228	EKIWEELSML
155	MAGE-2 271-278	FLWGPRAL
156	MAGE-2 271-279	FLWGPRALI
157	MAGE-2 278-286	LIETSYVKV
158	MAGE-2 277-286	ALIETSYVKV
159	MAGE-2 276-284	RALIETSYV

160	MAGE-2 279-287	IETSYVKVL
161	MAGE-2 278-287	LIETSYVKVL
162	MAGE-3 271-278	FLWGPRAL
163	MAGE-3 270-278	EFLWGPRAL
164	MAGE-3 271-279	FLWGPRALV
165	MAGE-3 276-284	RALVETSYV
166	MAGE-3 272-280	LWGPRALVE
167	MAGE-3 271-280	FLWGPRALVE
168	MAGE-3 27 2-281	LWGPRALVET
169	NY-ESO-1 82-90	GPESRLLEF
170	NY-ESO-1 83-91	PESRLLEFY
171	NY-ESO-1 82-91	GPESRLLEFY
172	NY-ESO-1 84-92	ESRLLEFYL
173	NY-ESO-1 86-94	RLLEFYLAM
174	NY-ESO-1 88-96	LEFYLAMPF
175	NY-ESO-1 87-96	LLEFYLAMPF
176	NY-ESO-1 93-102	AMPFATPMEA
177	NY-ESO-1 94-102	MPFATPMEA
178	NY-ESO-1 115-123	PLPVPGVLL
179	NY-ESO-1 114-123	PPLPVPGVLL
180	NY-ESO-1 116-123	LPVPGVLL
181	NY-ESO-1 103-112	ELARRSLAQD
182	NY-ESO-1 118-126	VPGVLLKEF
183	NY-ESO-1 117-126	PVPGVLLKEF
184	NY-ESO-1 116-123	LPVPGVLL
185	NY-ESO-1 127-135	TVSGNILTI
186	NY-ESO-1 126-135	FTVSGNILTI
187	NY-ESO-1 120-128	GVLLKEFTV
188	NY-ESO-1 121-130	VLLKEFTVSG
189	NY-ESO-1 122-130	LLKEFTVSG
190	NY-ESO-1 118-126	VPGVLLKEF
191	NY-ESO-1 117-126	PVPGVLLKEF
192	NY-ESO-1 139-147	AADHRQLQL
193	NY-ESO-1 148-156	SISSCLQQL
194	NY-ESO-1 147-156	LSISSCLQQL
195	NY-ESO-1 138-147	TAADHRQLQL
196	NY-ESO-1 161-169	WITQCFLPV
197	NY-ESO-1 157-165	SLLMWITQC
198	NY-ESO-1 150-158	SSCLQQLSL
199	NY-ESO-1 154-162	QQLSLLMWI
200	NY-ESO-1 151-159	SCLQQLSLL
201	NY-ESO-1 150-159	SSCLQQLSLL

202	NY-ESO-1 163-171	TQCFLPVFL
203	NY-ESO-1 162-171	ITQCFLPVFL
204	PRAME 219-227	PMQDIKMIL
205	PRAME 218-227	MPMQDIKMIL
206	PRAME 428-436	QHLIGLSNL
207	PRAME 427-436	LQHLIGLSNL
208	PRAME 429-436	HLIGLSNL
209	PRAME 431-439	IGLSNLTHV
210	PRAME 430-439	LIGLSNLTHV
211	PSA 53-61	VLVHPQWVL
212	PSA 52-61	GVLVHPQWVL
213	PSA 52-60	GVLVHPQWV
214	PSA 59-67	WVLTAHCI
215	PSA 54-63	LVHPQWVLTA
216	PSA 53-62	VLVHPQWVLT
217	PSA 54-62	LVHPQWVLT
218	PSA 66-73	CIRNKSVI
219	PSA 65-73	HCIRNKSVI
220	PSA 56-64	HPQWVLTAA
221	PSA 63-72	AAHCIRNKSV
222	PSCA 116-123	LLWGPGQL
223	PSCA 115-123	LLLWGPGQL
224	PSCA 114-123	GLLLWGPGQL
225	PSCA 99-107	ALQPAAAIL
226	PSCA 98-107	HALQPAAAIL
227	Tyr 128-137	APEKDKFFAY
228	Tyr 129-137	PEKDKFFAY
229	Tyr 130-138	EKDKFFAYL
230	Tyr 131-138	KDKFFAYL
231	Tyr 205-213	PAFLPWHRL
232	Tyr 204-213	APAFLPWHRL
233	Tyr 214-223	FLLRWEQEIQ
234	Tyr 212-220	RLFLLRWEQ
235	Tyr 191-200	GSEIWRDIDF
236	Tyr 192-200	SEIWRDIDF
237	Tyr 473-481	RIWSWLLGA
238	Tyr 476-484	SWLLGAAMV
239	Tyr 477-486	WLLGAAMVGA
240	Tyr 478-486	LLGAAMVGA
241	PSMA 4-12	LLHETDSAV
242	PSMA 13-21	ATARRPRWL
243	PSMA 53-61	TPKHNMKAF

244	PSMA 64-73	ELKAENIKKF
245	PSMA 69-77	NIKKFLH ¹ NF
246	PSMA 68-77	ENIKKFLH ¹ NF
247	PSMA 220-228	AGAKGVILY
248	PSMA 468-477	PLMYSLVHNL
249	PSMA 469-477	LMYSLVHNL
250	PSMA 463-471	RVDCTPLMY
251	PSMA 465-473	DCTPLMYSL
252	PSMA 507-515	SGMPRISKL
253	PSMA 506-515	FSGMPRISKL
254	NY-ESO-1 136-163	RLTAADHRQLQLSISSCLQQLSLLMWIT
255	NY-ESO-1 150-177	SSCLQQLSLLMWITQCFLPVFLAQPPSG

¹ This H was reported as Y in the SWISSPROT database.

Table 1C. SEQ ID NOS.* including epitopes in Example14.

SEQ ID NO.	IDENTITY	SEQUENCE
256	Mage-1 125-132	KAEMLESV
257	Mage-1 124-132	TKAEMLESV
258	Mage-1 123-132	VTKAEMLESV
259	Mage-1 128-136	MLESVIKNY
260	Mage-1 127-136	EMLESVIKNY
261	Mage-1 125-133	KAEMLESVI
262	Mage-1 146-153	KASESLQL
263	Mage-1 145-153	GKASESLQL
264	Mage-1 147-155	ASESLQLVF
265	Mage-1 153-161	LVFGIDVKE
266	Mage-1 114-121	LLKYRARE
267	Mage-1 106-113	VADLVGFL
268	Mage-1 105-113	KVADLVGFL
269	Mage-1 107-115	ADLVGFLL
270	Mage-1 106-115	VADLVGFLL
271	Mage-1 114-123	LLKYRAREPV
272	Mage-3 278-286	LVETSYVKV
273	Mage-3 277-286	ALVETSYVKV
274	Mage-3 285-293	KVLHHMVKI
275	Mage-3 283-291	YVKVLHHMV
276	Mage-3 275-283	PRALVETSY
277	Mage-3 274-283	GPRALVETSY
278	Mage-3 278-287	LVETSYVKVL
279	ED-B 4'-5	TIPEVPQL
280	ED-B 5'-5	DTIPEVPQL
281	ED-B 1-10	EVPQLTDLFS
282	ED-B 23-30	TPLNSSTI
283	ED-B 18-25	IGLRWTPL
284	ED-B 17-25	SIGLRWTPL
285	ED-B 25-33	LNSSTIIGY
286	ED-B 24-33	PLNSSTIIGY

287	ED-B 23-31	TPLNSSSTII
288	ED-B 31-38	IGYRITVV
289	ED-B 30-38	IIGYRITVV
290	ED-B 29-38	TIIGYRITVV
291	ED-B 31-39	IGYRITVVA
292	ED-B 30-39	IIGYRITVVA
293	CEA 184-191	SLPVSPRL
294	CEA 183-191	QSLPVSPRL
295	CEA 186-193	PVSPRLQL
296	CEA 185-193	LPVSPRLQL
297	CEA 184-193	SLPVSPRLQL
298	CEA 185-192	LPVSPRLQ
299	CEA 192-200	QLSNGNRTL
300	CEA 191-200	LQLSNGNRTL
301	CEA 179-187	WVNNQSLPV
302	CEA 186-194	PVSPRLQLS
303	CEA 362-369	SLPVSPRL
304	CEA 361-369	QSLPVSPRL
305	CEA 364-371	PVSPRLQL
306	CEA 363-371	LPVSPRLQL
307	CEA 362-371	SLPVSPRLQL
308	CEA 363-370	LPVSPRLQ
309	CEA 370-378	QLSNDNRTL
310	CEA 369-378	LQLSNDNRTL
311	CEA 357-365	WVNNQSLPV
312	CEA 360-368	NQSLPVSPR
313	CEA 540-547	SLPVSPRL
314	CEA 539-547	QSLPVSPRL
315	CEA 542-549	PVSPRLQL
316	CEA 541-549	LPVSPRLQL
317	CEA 540-549	SLPVSPRLQL
318	CEA 541-548	LPVSPRLQ
319	CEA 548-556	QLSNGNRTL
320	CEA 547-556	LQLSNGNRTL
321	CEA 535-543	WVNGQSLPV
322	CEA 533-541	LWWVNGQSL
323	CEA 532-541	YLWWVNGQSL
324	CEA 538-546	GQSLPVSPR
325	Her-2 30-37	DMKLRLPA
326	Her-2 28-37	GTDMLRLPA
327	Her-2 42-49	HLDMLRHL
328	Her-2 41-49	THLDMLRHL
329	Her-2 40-49	ETHLDMLRHL
330	Her-2 36-43	PASPETHL
331	Her-2 35-43	LPASPETHL
332	Her-2 34-43	RLPASPETHL
333	Her-2 38-46	SPETHDML
334	Her-2 37-46	ASPETHDML
335	Her-2 42-50	HLDMLRHLY
336	Her-2 41-50	THLDMLRHLY
337	Her-2 719-726	ELRKVKVL

338	Her-2 718-726	TELKVKVL
339	Her-2 717-726	ETELRKVKVL
340	Her-2 715-723	LKETELRKV
341	Her-2 714-723	ILKETELRKV
342	Her-2 712-720	MRILKETEL
343	Her-2 711-720	QMRILKETEL
344	Her-2 717-725	ETELRKVKV
345	Her-2 716-725	KETELRKVKV
346	Her-2 706-714	MPNQAQMRI
347	Her-2 705-714	AMPNQAQMRI
348	Her-2 706-715	MPNQAQMRI
349	HER-2 966-973	RPRFREL V
350	HER-2 965-973	CRPRFREL V
351	HER-2 968-976	RFREL VSEF
352	HER-2 967-976	PRFREL VSEF
353	HER-2 964-972	ECRPRFREL
354	NY-ESO-1 67-75	GAASGLNGC
355	NY-ESO-1 52-60	RASGPGGGA
356	NY-ESO-1 64-72	PHGGAASGL
357	NY-ESO-1 63-72	GPHGGAASGL
358	NY-ESO-1 60-69	APRGPHGGAA
359	PRAME 112-119	VRPRRWKL
360	PRAME 111-119	EVRPRRWKL
361	PRAME 113-121	RPRRWKLQV
362	PRAME 114-122	PRRWKLQVL
363	PRAME 113-122	RPRRWKLQVL
364	PRAME 116-124	RWKLQVLDL
365	PRAME 115-124	RRWKLOVLDL
366	PRAME 174-182	PVEVLVDLF
367	PRAME 199-206	VKRKKNVL
368	PRAME 198-206	KVKRKNVL
369	PRAME 197-206	EKVKRKKNVL
370	PRAME 198-205	KVKRKNV
371	PRAME 201-208	RKKNVRL
372	PRAME 200-208	KRKNVRL
373	PRAME 199-208	VKRKKNVRL
374	PRAME 189-196	DELFSLI
375	PRAME 205-213	VLRLCCKKL
376	PRAME 204-213	NVLRLCCKKL
377	PRAME 194-202	YLIEKVKRK
378	PRAME 74-81	QAWPFTCL
379	PRAME 73-81	VQAWPFTCL
380	PRAME 72-81	MVQAWPFTCL
381	PRAME 81-88	LPLGVLMK
382	PRAME 80-88	CLPLGVLMK
383	PRAME 79-88	TCLPLGVLMK
384	PRAME 84-92	GVLKMGQHL
385	PRAME 81-89	LPLGVLMKG
386	PRAME 80-89	CLPLGVLMKG
387	PRAME 76-85	WPFTCLPLGV
388	PRAME 51-59	ELFPPLFMA

389	PRAME 49-57	PRELFPPLF
390	PRAME 48-57	LPRELFPPLF
391	PRAME 50-58	RELFPPLFM
392	PRAME 49-58	PRELFPPLFM
393	PSA 239-246	RPSLYTKV
394	PSA 238-246	ERPSLYTKV
395	PSA 236-243	LPERPSLY
396	PSA 235-243	ALPERPSLY
397	PSA 241-249	SLYTKVVHY
398	PSA 240-249	PSLYTKVVHY
399	PSA 239-247	RPSLYTKVV
400	PSMA 211-218	GNKVKNAQ
401	PSMA 202-209	IARYGKVF
402	PSMA 217-225	AQLAGAKGV
403	PSMA 207-215	KVFRGNKVK
404	PSMA 211-219	GNKVKNAQL
405	PSMA 269-277	TPGYPANAY
406	PSMA 268-277	LTPGYPANAY
407	PSMA 271-279	GYPANAYAY
408	PSMA 270-279	PGYPANAYAY
409	PSMA 266-274	DPLTPGYPA
410	PSMA 492-500	SLYESWTKK
411	PSMA 491-500	KSLYESWTKK
412	PSMA 486-494	EGFEGKSLY
413	PSMA 485-494	DEGFEGKSLY
414	PSMA 498-506	TKKSPSPEF
415	PSMA 497-506	WTKKSPSPEF
416	PSMA 492-501	SLYESWTKKS
417	PSMA 725-732	WGEVKRQI
418	PSMA 724-732	AWGEVKRQI
419	PSMA 723-732	KAWGEVKRQI
420	PSMA 723-730	KAWGEVKR
421	PSMA 722-730	SKAWGEVKR
422	PSMA 731-739	QIYVAAFTV
423	PSMA 733-741	YVAAFTVQA
424	PSMA 725-733	WGEVKRQIY
425	PSMA 727-735	EVKRQIYVA
426	PSMA 738-746	TVQAAAETL
427	PSMA 737-746	FTVQAAAETL
428	PSMA 729-737	KRQIYVAAF
429	PSMA 721-729	PSKAWGEVK
430	PSMA 723-731	KAWGEVKRQ
431	PSMA 100-108	WKEFGLDSV
432	PSMA 99-108	QWKEFGLDSV
433	PSMA 102-111	EFGLDSVELA
434	SCP-1 126-134	ELRQKESKL
435	SCP-1 125-134	AELRQKESKL
436	SCP-1 133-141	KLQENRKII
437	SCP-1 298-305	QLEEKTKL
438	SCP-1 297-305	NQLEEKTKL
439	SCP-1 288-296	LLEESRDKV

440	SCP-1 287-296	FLLEESRDKV
441	SCP-1 291-299	ESRDKVNQL
442	SCP-1 290-299	EESRDKVNQL
443	SCP-1 475-483	EKEVHDLEY
444	SCP-1 474-483	REKEVHDLEY
445	SCP-1 480-488	DLEYSYCHY
446	SCP-1 477-485	EVHDLEYSY
447	SCP-1 477-486	EVHDLEYSYC
448	SCP-1 502-509	KLSSKREL
449	SCP-1 508-515	ELKNTEYF
450	SCP-1 507-515	RELKNTEYF
451	SCP-1 496-503	KRGQRPKL
452	SCP-1 494-503	LPKRGQRPKL
453	SCP-1 509-517	LKNTEYFTL
454	SCP-1 508-517	ELKNTEYFTL
455	SCP-1 506-514	KRELKNTEY
456	SCP-1 502-510	KLSSKRELK
457	SCP-1 498-506	GQRPKLSSK
458	SCP-1 497-506	RGQRPKLSSK
459	SCP-1 500-508	RPKLSSKRE
460	SCP-1 573-580	LEYVREEL
461	SCP-1 572-580	ELEYVREEL
462	SCP-1 571-580	NELEYVREEL
463	SCP-1 579-587	ELKQKREDEV
464	SCP-1 575-583	YVREELKQK
465	SCP-1 632-640	QLNVYEIKV
466	SCP-1 630-638	SKQLNVYEI
467	SCP-1 628-636	AESKQLNVY
468	SCP-1 627-636	TAESKQLNVY
469	SCP-1 638-645	IKVNKLEL
470	SCP-1 637-645	EIKVNKLEL
471	SCP-1 636-645	YEIKVNKLEL
472	SCP-1 642-650	KLELELESA
473	SCP-1 635-643	VYEIKVNKL
474	SCP-1 634-643	NVYEIKVNKL
475	SCP-1 646-654	ELESAKQKF
476	SCP-1 642-650	KLELELESA
477	SCP-1 646-654	ELESAKQKF
478	SCP-1 771-778	KEKCLKREA
479	SCP-1 777-785	EAKENTATL
480	SCP-1 776-785	REAKENTATL
481	SCP-1 773-782	KLKREAKENT
482	SCP-1 112-119	EAEEKIKKW
483	SCP-1 101-109	GLSRVYSKL
484	SCP-1 100-109	EGLSRVYSKL
485	SCP-1 108-116	KLYKEAEKI
486	SCP-1 98-106	NSEGLSRVY
487	SCP-1 97-106	ENSEGLSRVY
488	SCP-1 102-110	LSRVYSKLY
489	SCP-1 101-110	GLSRVYSKLY
490	SCP-1 96-105	LENSEGLSRV

491	SCP-1 108-117	KLYKEAEKIK
492	SCP-1 949-956	REDRWAVI
493	SCP-1 948-956	MRREDRWAVI
494	SCP-1 947-956	KMREDRWAVI
495	SCP-1 947-955	KMREDRWAV
496	SCP-1 934-942	TTPGSTLKF
497	SCP-1 933-942	LTPGSTLKF
498	SCP-1 937-945	GSTLKGAI
499	SCP-1 945-953	IRKMREDRW
500	SCP-1 236-243	RLEMHFKL
501	SCP-1 235-243	SRLEMHFKL
502	SCP-1 242-250	KLKEDYEKI
503	SCP-1 249-257	KIQHLEQY
504	SCP-1 248-257	EKIQHLEQY
505	SCP-1 233-242	ENSRLEMHF
506	SCP-1 236-245	RLEMHFKLKE
507	SCP-1 324-331	LEDIKVSL
508	SCP-1 323-331	ELEDIKVSL
509	SCP-1 322-331	KELEDIKVSL
510	SCP-1 320-327	LTKELEDI
511	SCP-1 319-327	HLTKELEDI
512	SCP-1 330-338	SLQRSVSTQ
513	SCP-1 321-329	TKELEDIKV
514	SCP-1 320-329	LTKELEDIKV
515	SCP-1 326-335	DIKVSQRSV
516	SCP-1 281-288	KMKDLTFL
517	SCP-1 280-288	NKMKDLTFL
518	SCP-1 279-288	ENKMKDLTFL
519	SCP-1 288-296	LLEESRDKV
520	SCP-1 287-296	FLLEESRDKV
521	SCP-1 291-299	ESRDKVNQL
522	SCP-1 290-299	EESRDKVNQL
523	SCP-1 277-285	EKENKMKDL
524	SCP-1 276-285	TEKENKMKDL
525	SCP-1 279-287	ENKMKDLTF
526	SCP-1 218-225	IEKMITAF
527	SCP-1 217-225	NIEKMITAF
528	SCP-1 216-225	SNIEKMITAF
529	SCP-1 223-230	TAFEELRV
530	SCP-1 222-230	ITAFEELRV
531	SCP-1 221-230	MITAFEELRV
532	SCP-1 220-228	KMITAFEEL
533	SCP-1 219-228	EKMITAFEEL
534	SCP-1 227-235	ELRVQAENS
535	SCP-1 213-222	DLNSNIEKMI
536	SCP-1 837-844	WTSAKNTL
537	SCP-1 846-854	TPLPKAYTV
538	SCP-1 845-854	STPLPKAYTV
539	SCP-1 844-852	LSTPLPKAY
540	SCP-1 843-852	TLSTPLPKAY
541	SCP-1 842-850	NTLSTPLPK

542	SCP-1 841-850	KNTLSTPLPK
543	SCP-1 828-835	ISKDKRDY
544	SCP-1 826-835	HGISKDKRDY
545	SCP-1 832-840	KRDYLWTS
546	SCP-1 829-838	SKDKRDYLWT
547	SCP-1 279-286	ENKMKDLT
548	SCP-1 260-268	EINDKEKQV
549	SCP-1 274-282	QITEKENKM
550	SCP-1 269-277	SLLLIQITE
551	SCP-1 453-460	FEKIAEEL
552	SCP-1 452-460	QFEKIAEEL
553	SCP-1 451-460	KQFEKIAEEL
554	SCP-1 449-456	DNKQFEKI
555	SCP-1 448-456	YDNKQFEKI
556	SCP-1 447-456	LYDNKQFEKI
557	SCP-1 440-447	LGEKETLL
558	SCP-1 439-447	VLGEKETLL
559	SCP-1 438-447	KVLGEKETLL
560	SCP-1 390-398	LLRTEQORL
561	SCP-1 389-398	ELLRTEQORL
562	SCP-1 393-401	TEQORLENY
563	SCP-1 392-401	RTEQORLENY
564	SCP-1 402-410	EDQLILTM
565	SCP-1 397-406	RLENYEDQLI
566	SCP-1 368-375	KARAAHSF
567	SCP-1 376-384	VVTEFETTV
568	SCP-1 375-384	FVVTEFETTV
569	SCP-1 377-385	VTEFETTVC
570	SCP-1 376-385	VVTEFETTVC
571	SCP-1 344-352	DLQIATNTI
572	SCP-1 347-355	IATNTICQL
573	SCP-1 346-355	QIATNTICQL
574	SSX4 57-65	VMTKLGFKY
575	SSX4 53-61	LNVEVMTKL
576	SSX4 52-61	KLNVEVMTKL
577	SSX4 66-74	TLPPFMRSK
578	SSX4 110-118	KIMPKKPAE
579	SSX4 103-112	SLQRIFPKIM
580	Tyr 463-471	YIKSYLEQA
581	Tyr 459-467	SFQDYIKSY
582	Tyr 458-467	DSFQDYIKSY
583	Tyr 507-514	LPEEKQPL
584	Tyr 506-514	QLPEEKQPL
585	Tyr 505-514	KQLPEEKQPL
586	Tyr 507-515	LPEEKQPLL
587	Tyr 506-515	QLPEEKQPLL
588	Tyr 497-505	SLLCRHKRK
589	ED-B domain of Fibronectin	EVPQLTDLSFVDITDSSIGLRWTPLNSSSTHGYRI TVVAAGEGIPFEDFVDSSVGYYTGTGLEPGID YDISVITLINGGESAPTTLTQQT
590	ED-B domain of	CTFDNLSPGLEYNVSVYTVKDDKESVPISDTIIP

	Fibronectin with flanking sequence from Fribronectin	EVPQLTDLFSFVDITDSSIGLRWTPPLNSSTIGYRI TVVAAGEGIPFEDFVDSSVGYYTVTGLEPGID YDISVITLINGGESAPTTLTQQT AVPPPTDLRFTNIGPDTMRVTW
591	ED-B domain of Fibronectin cds	Accession number: X07717
592	CEA protein	Accession number: P06731
593	CEA cDNA	Accession number: NM_004363
594	Her2/Neu protein	Accession number: P04626
595	Her2/Neu cDNA	Accession number: M11730
596	SCP-1 protein	Accession number: Q15431
597	SCP-1 cDNA	Accession number: X95654
598	SSX-4 protein	Accession number: O60224
599	SSX-4 cDNA	Accession number: NM_005636

*Any of SEQ ID NOS. 1, 8, 9, 11-23, 26-29, 32-44, 47-54, 56-63, 66-68 88-253, and 256-588 can be useful as epitopes in any of the various embodiments of the invention. Any of SEQ ID NOS. 10, 30, 31, 45, 46, 55, 64, 65, 69, 254, and 255 can be useful as sequences containing epitopes or epitope clusters, as described in various embodiments of the invention.

**All accession numbers used here and throughout can be accessed through the NCBI databases, for example, through the Entrez seek and retrieval system on the world wide web.

Note that the following discussion sets forth the inventors' understanding of the operation of the invention. However, it is not intended that this discussion limit the patent to any particular theory of operation not set forth in the claims.

In pursuing the development of epitope vaccines others have generated lists of predicted epitopes based on MHC binding motifs. Such peptides can be immunogenic, but may not correspond to any naturally produced antigenic fragment so that whole antigen will not elicit a similar response or sensitize a target cell to cytolysis by CTL. Therefore such lists do not differentiate between those sequences that can be useful as vaccines and those that cannot. Efforts to determine which of these predicted epitopes are in fact naturally produced have often relied on screening their reactivity with tumor infiltrating lymphocytes (TIL). However, TIL are strongly biased to recognize immune epitopes whereas tumors (and chronically infected cells) will generally present housekeeping epitopes. Thus, unless the epitope is produced by both the housekeeping and immuno- proteasomes, the target cell will generally not be recognized by CTL induced with TIL-identified epitopes. The epitopes of the present invention, in contrast, are generated by the action a specified proteasome, indicating that they can be naturally produced, and enabling their appropriate use. The importance of the distinction between housekeeping and immune epitopes to vaccine design in more fully set forth in PCT publication WO 01/82963A2.

The epitopes of the invention include or encode polypeptide fragments of TAAs that are precursors or products of proteasomal cleavage by a housekeeping or immune proteasome, and that have known or predicted affinity for at least one allele of MHC I. In some embodiments, the epitopes include or encode a polypeptide of about 6 to 25 amino acids in length, preferably about 7

to 20 amino acids in length, more preferably about 8 to 15 amino acids in length, and still more preferably 9 or 10 amino acids in length. However, it is understood that the polypeptides can be larger as long as they do not contain sequences that cause the polypeptides to be directed away from the proteasome or to be destroyed by the proteasome. For immune epitopes, if the larger peptides do not contain such sequences, they can be processed in the pAPC by the immune proteasome. Housekeeping epitopes may also be embedded in longer sequences provided that the sequence is adapted to facilitate liberation of the epitope's C-terminus by action of the immunoproteasome. The sequences of these epitopes can be subjected to computer analysis in order to calculate physical, biochemical, immunologic, or molecular genetic properties such as mass, isoelectric point, predicted mobility in electrophoresis, predicted binding to other MHC molecules, melting temperature of nucleic acid probes, reverse translations, similarity or homology to other sequences, and the like.

In constructing the polynucleotides encoding the polypeptide epitopes of the invention, the gene sequence of the associated TAA can be used, or the polynucleotide can be assembled from any of the corresponding codons. For a 10 amino acid epitope this can constitute on the order of 10^6 different sequences, depending on the particular amino acid composition. While large, this is a distinct and readily definable set representing a miniscule fraction of the $>10^{18}$ possible polynucleotides of this length, and thus in some embodiments, equivalents of a particular sequence disclosed herein encompass such distinct and readily definable variations on the listed sequence. In choosing a particular one of these sequences to use in a vaccine, considerations such as codon usage, self-complementarity, restriction sites, chemical stability, etc. can be used as will be apparent to one skilled in the art.

The invention contemplates producing peptide epitopes. Specifically these epitopes are derived from the sequence of a TAA, and have known or predicted affinity for at least one allele of MHC I. Such epitopes are typically identical to those produced on target cells or pAPCs.

Compositions Containing Active Epitopes

Embodiments of the present invention provide polypeptide compositions, including vaccines, therapeutics, diagnostics, pharmacological and pharmaceutical compositions. The various compositions include newly identified epitopes of TAAs, as well as variants of these epitopes. Other embodiments of the invention provide polynucleotides encoding the polypeptide epitopes of the invention. The invention further provides vectors for expression of the polypeptide epitopes for purification. In addition, the invention provides vectors for the expression of the polypeptide epitopes in an APC for use as an anti-tumor vaccine. Any of the epitopes or antigens, or nucleic acids encoding the same, from Table 1A can be used. Other embodiments relate to methods of making and using the various compositions.

A general architecture for a class I MHC-binding epitope can be described, and has been reviewed more extensively in Madden, D.R. *Annu. Rev. Immunol.* 13:587-622, 1995. Much of the binding energy arises from main chain contacts between conserved residues in the MHC molecule and the N- and C-termini of the peptide. Additional main chain contacts are made but vary among MHC alleles. Sequence specificity is conferred by side chain contacts of so-called anchor residues with pockets that, again, vary among MHC alleles. Anchor residues can be divided into primary and secondary. Primary anchor positions exhibit strong preferences for relatively well-defined sets of amino acid residues. Secondary positions show weaker and/or less well-defined preferences that can often be better described in terms of less favored, rather than more favored, residues. Additionally, residues in some secondary anchor positions are not always positioned to contact the pocket on the MHC molecule at all. Thus, a subset of peptides exists that bind to a particular MHC molecule and have a side chain-pocket contact at the position in question and another subset exists that show binding to the same MHC molecule that does not depend on the conformation the peptide assumes in the peptide-binding groove of the MHC molecule. The C-terminal residue (P₉) is preferably a primary anchor residue. For many of the better studied HLA molecules (e.g. A2, A68, B27, B7, B35, and B53) the second position (P2) is also an anchor residue. However, central anchor residues have also been observed including P3 and P5 in HLA-B8, as well as P5 and P₋₃ in the murine MHC molecules H-2D^b and H-2K^b, respectively. Since more stable binding will generally improve immunogenicity, anchor residues are preferably conserved or optimized in the design of variants, regardless of their position.

Because the anchor residues are generally located near the ends of the epitope, the peptide can buckle upward out of the peptide-binding groove allowing some variation in length. Epitopes ranging from 8-11 amino acids have been found for HLA-A68, and up to 13 amino acids for HLA-A2. In addition to length variation between the anchor positions, single residue truncations and extensions have been reported and the N- and C-termini, respectively. Of the non-anchor residues, some point up out of the groove, making no contact with the MHC molecule but being available to contact the TCR, very often P1, P4, and P₋₁ for HLA-A2. Others of the non-anchor residues can become interposed between the upper edges of the peptide-binding groove and the TCR, contacting both. The exact positioning of these side chain residues, and thus their effects on binding, MHC fine conformation, and ultimately immunogenicity, are highly sequence dependent. For an epitope to be highly immunogenic it must not only promote stable enough TCR binding for activation to occur, but the TCR must also have a high enough off-rate that multiple TCR molecules can interact sequentially with the same peptide-MHC complex (Kalergis, A.M. et al., *Nature Immunol.* 2:229-234, 2001). Thus without further information about the ternary complex, both conservative and non-conservative substitutions at these positions merit consideration when designing variants.

The polypeptide epitope variants can be made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations. Variants can be derived from substitution, deletion or insertion of one or more amino acids as compared with the native sequence. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a threonine with a serine. Such replacements are referred to as conservative amino acid replacements, and all appropriate conservative amino acid replacements are considered to be embodiments of one invention. Insertions or deletions can optionally be in the range of about 1 to 4, preferably 1 to 2, amino acids. It is generally preferable to maintain the "anchor positions" of the peptide which are responsible for binding to the MHC molecule in question. Indeed, immunogenicity of peptides can be improved in many cases by substituting more preferred residues at the anchor positions (Franco, et al., *Nature Immunology*, 1(2):145-150, 2000). Immunogenicity of a peptide can also often be improved by substituting bulkier amino acids for small amino acids found in non-anchor positions while maintaining sufficient cross-reactivity with the original epitope to constitute a useful vaccine. The variation allowed can be determined by routine insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the polypeptide epitope. Because the polypeptide epitope is often 9 amino acids, the substitutions preferably are made to the shortest active epitope, for example, an epitope of 9 amino acids.

Variants can also be made by adding any sequence onto the N-terminus of the polypeptide epitope variant. Such N-terminal additions can be from 1 amino acid up to at least 25 amino acids. Because peptide epitopes are often trimmed by N-terminal exopeptidases active in the pAPC, it is understood that variations in the added sequence can have no effect on the activity of the epitope. In preferred embodiments, the amino acid residues between the last upstream proteasomal cleavage site and the N-terminus of the MHC epitope do not include a proline residue. Serwold, T. et al., *Nature Immunol.* 2:644-651, 2001. Accordingly, effective epitopes can be generated from precursors larger than the preferred 9-mer class I motif.

Peptides are useful to the extent that they correspond to epitopes actually displayed by MHC I on the surface of a target cell or a pACP. A single peptide can have varying affinities for different MHC molecules, binding some well, others adequately, and still others not appreciably (Table 2). MHC alleles have traditionally been grouped according to serologic reactivity which does not reflect the structure of the peptide-binding groove, which can differ among different alleles of the same type. Similarly, binding properties can be shared across types; groups based on shared binding properties have been termed supertypes. There are numerous alleles of MHC I in the human population; epitopes specific to certain alleles can be selected based on the genotype of the patient.

Table 2.
Predicted Binding of Tyrosinase₂₀₇₋₂₁₆ (SEQ ID NO. 1) to Various MHC types

MHC I type	*Half time of dissociation (min)
A1	0.05
A*0201	1311.
A*0205	50.4
A3	2.7
A*1101 (part of the A3 supertype)	0.012
A24	6.0
B7	4.0
B8	8.0
B14 (part of the B27 supertype)	60.0
B*2702	0.9
B*2705	30.0
B*3501 (part of the B7 supertype)	2.0
B*4403	0.1
B*5101 (part of the B7 supertype)	26.0
B*5102	55.0
B*5801	0.20
B60	0.40
B62	2.0

*HLA Peptide Binding Predictions (internet [http:// access at bimas.dcrt.nih.gov/molbio/hla_bin](http://access.bimas.dcrt.nih.gov/molbio/hla_bin))

5

In further embodiments of the invention, the epitope, as peptide or encoding polynucleotide, can be administered as a vaccine or immunogenic composition, alone or in combination with various adjuvants, carriers, or excipients. It should be noted that although the term vaccine may be used herein, the discussion can be applied and used with any of the other compositions mentioned herein. Particularly advantageous adjuvants include various cytokines and oligonucleotides containing immunostimulatory sequences (as set forth in greater detail in the co-pending applications referenced herein). Additionally the polynucleotide encoded epitope can be contained in a virus (e.g. *vaccinia* or adenovirus) or in a microbial host cell (e.g. *Salmonella* or *Listeria monocytogenes*) which is then used as a vector for the polynucleotide (Dietrich, G. et al. Nat. Biotech. 16:181-185, 1998). Alternatively a pAPC can be transformed, *ex vivo*, to express the epitope, or pulsed with peptide epitope, to be itself administered as a vaccine. To increase efficiency of these processes, the encoded epitope can be carried by a viral or bacterial vector, or complexed with a ligand of a receptor found on pAPC. Similarly the peptide epitope can be complexed with or conjugated to a pAPC ligand. A vaccine can be composed of more than a single epitope.

10

15

20

Particularly advantageous strategies for incorporating epitopes, and combining them with epitope clusters, into a vaccine are disclosed in U.S. Patent Application No. 09/560,465 entitled "EPIPOPE SYNCHRONIZATION IN ANTIGEN PRESENTING CELLS," filed on April 28,

2000. Epitope clusters for use in connection with this invention are disclosed in U.S. Patent Application No. 09/561,571 entitled "EPITOPE CLUSTERS," filed on April 28, 2000.

Preferred embodiments of the present invention are directed to vaccines and methods for causing a pAPC or population of pAPCs to present housekeeping epitopes that correspond to the epitopes displayed on a particular target cell. Any of the epitopes or antigens in Table 1A, can be used for example. In one embodiment, the housekeeping epitope is a TuAA epitope processed by the housekeeping proteasome of a particular tumor type. In another embodiment, the housekeeping epitope is a virus-associated epitope processed by the housekeeping proteasome of a cell infected with a virus. This facilitates a specific T cell response to the target cells. Concurrent expression by the pAPCs of multiple epitopes, corresponding to different induction states (pre- and post-attack), can drive a CTL response effective against target cells as they display either housekeeping epitopes or immune epitopes.

By having both housekeeping and immune epitopes present on the pAPC, this embodiment can optimize the cytotoxic T cell response to a target cell. With dual epitope expression, the pAPCs can continue to sustain a CTL response to the immune-type epitope when the tumor cell switches from the housekeeping proteasome to the immune proteasome with induction by IFN, which, for example, may be produced by tumor-infiltrating CTLs.

In a preferred embodiment, immunization of a patient is with a vaccine that includes a housekeeping epitope. Many preferred TAAs are associated exclusively with a target cell, particularly in the case of infected cells. In another embodiment, many preferred TAAs are the result of deregulated gene expression in transformed cells, but are found also in tissues of the testis, ovaries and fetus. In another embodiment, useful TAAs are expressed at higher levels in the target cell than in other cells. In still other embodiments, TAAs are not differentially expressed in the target cell compare to other cells, but are still useful since they are involved in a particular function of the cell and differentiate the target cell from most other peripheral cells; in such embodiments, healthy cells also displaying the TAA may be collaterally attacked by the induced T cell response, but such collateral damage is considered to be far preferable to the condition caused by the target cell.

A preferred embodiment of the present invention includes a method of administering a vaccine including a housekeeping epitope to induce a therapeutic immune response. The vaccine is administered to a patient in a manner consistent with the standard vaccine delivery protocols that are well known in the art. Methods of administering epitopes of TAAs include, without limitation, transdermal, intranodal, perinodal, oral, intravenous, intradermal, intramuscular, intraperitoneal, and mucosal administration. A particularly useful method of vaccine delivery to elicit a CTL response is disclosed in PCT Publication No. WO 99/01283, entitled "A METHOD OF INDUCING A CTL RESPONSE," filed on July 10, 1998.

Because the epitope synchronization system has utility in inducing a cell mediated immune response, a vaccine to induce a specific T cell response to a target cell is likewise included in a preferred embodiment of the present invention. The vaccine contains a housekeeping epitope in a concentration effective to cause a pAPC or populations of pAPCs to display housekeeping epitopes. Advantagously, the vaccine can include a plurality of housekeeping epitopes or one or more housekeeping epitopes optionally in combination with one or more immune epitopes. Formulations of the vaccine contain peptides and/or nucleic acids in a concentration sufficient to cause pAPCs to present the epitopes. The formulations preferably contain epitopes in a total concentration of about 1µg-1mg/100µl of vaccine preparation. Conventional dosages and dosing for peptide vaccines and/or nucleic acid vaccines can be used with the present invention, and such dosing regimens are well understood in the art. In one embodiment, a single dosage for an adult human may advantageously be from about 1 to about 5000 µl of such a composition, administered one time or multiple times, e.g., in 2, 3, 4 or more dosages separated by 1 week, 2 weeks, 1 month, or more. insulin pump delivers 1 ul per hour (lowest frequency) ref intranodal method patent.

The compositions and methods of the invention disclosed herein further contemplate incorporating adjuvants into the formulations in order to enhance the performance of the vaccines. Specifically, the addition of adjuvants to the formulations is designed to enhance the delivery or uptake of the epitopes by the pAPCs. The adjuvants contemplated by the present invention are known by those of skill in the art and include, for example, GMCSF, GCSF, IL-2, IL-12, BCG, tetanus toxoid, osteopontin, and ETA-1.

In some embodiments of the invention, the vaccines can include a recombinant organism, such as a virus, bacterium or parasite, genetically engineered to express an epitope in a host. For example, *Listeria monocytogenes*, a gram-positive, facultative intracellular bacterium, is a potent vector for targeting TuAAs to the immune system. In a preferred embodiment, this vector can be engineered to express a housekeeping epitope to induce therapeutic responses. The normal route of infection of this organism is through the gut and can be delivered orally. In another embodiment, an adenovirus (Ad) vector encoding a housekeeping epitope for a TuAA can be used to induce anti-virus or anti-tumor responses. Bone marrow-derived dendritic cells can be transduced with the virus construct and then injected, or the virus can be delivered directly via subcutaneous injection into an animal to induce potent T-cell responses. Another embodiment employs a recombinant vaccinia virus engineered to encode amino acid sequences corresponding to a housekeeping epitope for a TAA. Vaccinia viruses carrying constructs with the appropriate nucleotide substitutions in the form of a minigene construct can direct the expression of a housekeeping epitope, leading to a therapeutic T cell response against the epitope.

The immunization with DNA requires that APCs take up the DNA and express the encoded proteins or peptides. It is possible to encode a discrete class I peptide on the DNA. By

immunizing with this construct, APCs can be caused to express a housekeeping epitope, which is then displayed on class I MHC on the surface of the cell for stimulating an appropriate CTL response. Constructs generally relying on termination of translation or non-proteasomal proteases for generation of proper termini of housekeeping epitopes have been described in U.S. Patent application No. 09/561,572 entitled EXPRESSION VECTORS ENCODING EPITOPES OF
5 TARGET-ASSOCIATED ANTIGENS, filed on April 28, 2000.

It can be desirable to express housekeeping peptides in the context of a larger protein. Processing can be detected even when a small number of amino acids are present beyond the terminus of an epitope. Small peptide hormones are usually proteolytically processed from longer
10 translation products, often in the size range of approximately 60-120 amino acids. This fact has led some to assume that this is the minimum size that can be efficiently translated. In some embodiments, the housekeeping peptide can be embedded in a translation product of at least about 60 amino acids. In other embodiments the housekeeping peptide can be embedded in a translation product of at least about 50, 30, or 15 amino acids.

Due to differential proteasomal processing, the immune proteasome of the pAPC produces peptides that are different from those produced by the housekeeping proteasome in peripheral body cells. Thus, in expressing a housekeeping peptide in the context of a larger protein, it is preferably expressed in the APC in a context other than its full length native sequence, because, as a housekeeping epitope, it is generally only efficiently processed from the native protein by the
15 housekeeping proteasome, which is not active in the APC. In order to encode the housekeeping epitope in a DNA sequence encoding a larger protein, it is useful to find flanking areas on either side of the sequence encoding the epitope that permit appropriate cleavage by the immune proteasome in order to liberate that housekeeping epitope. Such a sequence ensuring epitope synchronization is referred to hereinafter as a SYNCHROTOPE™. Altering flanking amino acid
20 residues at the N-terminus and C-terminus of the desired housekeeping epitope can facilitate appropriate cleavage and generation of the housekeeping epitope in the APC. Sequences embedding housekeeping epitopes can be designed *de novo* and screened to determine which can be successfully processed by immune proteasomes to liberate housekeeping epitopes.

Alternatively, another strategy is very effective for identifying sequences allowing
30 production of housekeeping epitopes in APC. A contiguous sequence of amino acids can be generated from head to tail arrangement of one or more housekeeping epitopes. A construct expressing this sequence is used to immunize an animal, and the resulting T cell response is evaluated to determine its specificity to one or more of the epitopes in the array. By definition, these immune responses indicate housekeeping epitopes that are processed in the pAPC effectively.
35 The necessary flanking areas around this epitope are thereby defined. The use of flanking regions of about 4-6 amino acids on either side of the desired peptide can provide the necessary

information to facilitate proteasome processing of the housekeeping epitope by the immune proteasome. Therefore, a SYNCHROTOPE™ of approximately 16-22 amino acids can be inserted into, or fused to, any protein sequence effectively to result in that housekeeping epitope being produced in an APC. In alternate embodiments the whole head-to-tail array of epitopes, or just the epitopes immediately adjacent to the correctly processed housekeeping epitope can be similarly transferred from a test construct to a vaccine vector.

In a preferred embodiment, the housekeeping epitopes can be embedded between known immune epitopes, or segments of such, thereby providing an appropriate context for processing. The abutment of housekeeping and immune epitopes can generate the necessary context to enable the immune proteasome to liberate the housekeeping epitope, or a larger fragment, preferably including a correct C-terminus. It can be useful to screen constructs to verify that the desired epitope is produced. The abutment of housekeeping epitopes can generate a site cleavable by the immune proteasome. Some embodiments of the invention employ known epitopes to flank housekeeping epitopes in test substrates; in others, screening as described below are used whether the flanking regions are arbitrary sequences or mutants of the natural flanking sequence, and whether or not knowledge of proteasomal cleavage preferences are used in designing the substrates.

Cleavage at the mature N-terminus of the epitope, while advantageous, is not required, since a variety of N-terminal trimming activities exist in the cell that can generate the mature N-terminus of the epitope subsequent to proteasomal processing. It is preferred that such N-terminal extension be less than about 25 amino acids in length and it is further preferred that the extension have few or no proline residues. Preferably, in screening, consideration is given not only to cleavage at the ends of the epitope (or at least at its C-terminus), but consideration also can be given to ensure limited cleavage within the epitope.

Shotgun approaches can be used in designing test substrates and can increase the efficiency of screening. In one embodiment multiple epitopes can be assembled one after the other, with individual epitopes possibly appearing more than once. The substrate can be screened to determine which epitopes can be produced. In the case where a particular epitope is of concern a substrate can be designed in which it appears in multiple different contexts. When a single epitope appearing in more than one context is liberated from the substrate additional secondary test substrates, in which individual instances of the epitope are removed, disabled, or are unique, can be used to determine which are being liberated and truly constitute SYNCHROTOPE™s.

Several readily practicable screens exist. A preferred *in vitro* screen utilizes proteasomal digestion analysis, using purified immune proteasomes, to determine if the desired housekeeping epitope can be liberated from a synthetic peptide embodying the sequence in question. The position of the cleavages obtained can be determined by techniques such as mass spectrometry, HPLC, and

N-terminal pool sequencing; as described in greater detail in U. S. Patent Applications entitled METHOD OF EPITOPE DISCOVERY, EPITOPE SYNCHRONIZATION IN ANTIGEN PRESENTING CELLS, two Provisional U. S. Patent Applications entitled EPITOPE SEQUENCES, which are all cited and incorporated by reference above.

5 Alternatively, *in vivo* screens such as immunization or target sensitization can be employed. For immunization a nucleic acid construct capable of expressing the sequence in question is used. Harvested CTL can be tested for their ability to recognize target cells presenting the housekeeping epitope in question. Such targets cells are most readily obtained by pulsing cells expressing the appropriate MHC molecule with synthetic peptide embodying the mature
10 housekeeping epitope. Alternatively, cells known to express housekeeping proteasome and the antigen from which the housekeeping epitope is derived, either endogenously or through genetic engineering, can be used. To use target sensitization as a screen, CTL, or preferably a CTL clone, that recognizes the housekeeping epitope can be used. In this case it is the target cell that expresses the embedded housekeeping epitope (instead of the pAPC during immunization) and it must
15 express immune proteasome. Generally, the target cell can be transformed with an appropriate nucleic acid construct to confer expression of the embedded housekeeping epitope. Loading with a synthetic peptide embodying the embedded epitope using peptide loaded liposomes or a protein transfer reagent such as BIOPORTER™ (Gene Therapy Systems, San Diego, CA) represents an alternative.

20 Additional guidance on nucleic acid constructs useful as vaccines in accordance with the present invention are disclosed in U.S. Patent Application No. 09/561,572 entitled "EXPRESSION VECTORS ENCODING EPITOPES OF TARGET-ASSOCIATED ANTIGENS," filed on April 28, 2000. Further, expression vectors and methods for their design, which are useful in accordance with the present invention are disclosed in U.S. Patent Application No. 60/336,968 (attorney
25 docket number CTLIMM.022PR) entitled "EXPRESSION VECTORS ENCODING EPITOPES OF TARGET-ASSOCIATED ANTIGENS AND METHODS FOR THEIR DESIGN," filed on 11/7/2001, which is incorporated by reference in its entirety.

 A preferred embodiment of the present invention includes a method of administering a vaccine including an epitope (or epitopes) to induce a therapeutic immune response. The vaccine
30 is administered to a patient in a manner consistent with the standard vaccine delivery protocols that are known in the art. Methods of administering epitopes of TAAs including, without limitation, transdermal, intranodal, perinodal, oral, intravenous, intradermal, intramuscular, intraperitoneal, and mucosal administration, including delivery by injection, instillation or inhalation. A particularly useful method of vaccine delivery to elicit a CTL response is disclosed in Australian
35 Patent No. 739189 issued January 17, 2002; U.S. Patent Application No. 09/380,534, filed on

September 1, 1999; and a Continuation-in-Part thereof U.S. Patent Application No. 09/776,232 both entitled "A METHOD OF INDUCING A CTL RESPONSE," filed on February 2, 2001.

Reagents Recognizing Epitopes

5 In another aspect of the invention, proteins with binding specificity for the epitope and the epitope-MHC molecule complex are contemplated, as well as the isolated cells by which they can be expressed. In one set of embodiments these reagents take the form of immunoglobulins: polyclonal sera or monoclonal antibodies (mAb), methods for the generation of which are well known in the art. Generation of mAb with specificity for peptide-MHC molecule complexes is known in the art. See, for example, Aharoni et al. *Nature* 351:147-150, 1991; Andersen et al. *Proc.*
10 *Natl. Acad. Sci. USA* 93:1820-1824, 1996; Dadaglio et al. *Immunity* 6:727-738, 1997; Duc et al. *Int. Immunol.* 5:427-431, 1993; Eastman et al. *Eur. J. Immunol.* 26:385-393, 1996; Engberg et al. *Immunotechnology* 4:273-278, 1999; Porgdor et al. *Immunity* 6:715-726, 1997; Puri et al. *J. Immunol.* 158:2471-2476, 1997; and Polakova, K., et al. *J. Immunol.* 165 342-348, 2000; all of which are hereby incorporated by reference in their entirety.

15 In other embodiments the compositions can be used to induce and generate, *in vivo* and *in vitro*, T-cells specific for the any of the epitopes, including those listed in Table 1A, for example. Thus, embodiments also relate to and include isolated T cells, T cell clones, T cell hybridomas, or a protein containing the T cell receptor (TCR) binding domain derived from the cloned gene, as well as a recombinant cell expressing such a protein. Such TCR derived proteins can be simply the
20 extra-cellular domains of the TCR, or a fusion with portions of another protein to confer a desired property or function. One example of such a fusion is the attachment of TCR binding domains to the constant regions of an antibody molecule so as to create a divalent molecule. The construction and activity of molecules following this general pattern have been reported, for example, Plaksin, D. et al. *J. Immunol.* 158:2218-2227, 1997 and Lebowitz, M.S. et al. *Cell Immunol.* 192:175-184,
25 1999, which are hereby incorporated by reference in their entirety. The more general construction and use of such molecules is also treated in U.S. patent 5,830,755 entitled T CELL RECEPTORS AND THEIR USE IN THERAPEUTIC AND DIAGNOSTIC METHODS, which is hereby incorporated by reference in its entirety.

The generation of such T cells can be readily accomplished by standard immunization of
30 laboratory animals, and reactivity to human target cells can be obtained by immunizing with human target cells or by immunizing HLA-transgenic animals with the antigen/epitope. For some therapeutic approaches T cells derived from the same species are desirable. While such a cell can be created by cloning, for example, a murine TCR into a human T cell as contemplated above, *in vitro* immunization of human cells offers a potentially faster option. Techniques for *in vitro*
35 immunization, even using naive donors, are known in the field, for example, Stauss et al., *Proc. Natl. Acad. Sci. USA* 89:7871-7875, 1992; Salgaller et al. *Cancer Res.* 55:4972-4979, 1995; Tsai et

al., *J. Immunol.* 158:1796-1802, 1997; and Chung et al., *J. Immunother.* 22:279-287, 1999; which are hereby incorporated by reference in their entirety.

Any of these molecules can be conjugated to enzymes, radiochemicals, fluorescent tags, and toxins, so as to be used in the diagnosis (imaging or other detection), monitoring, and treatment of the pathogenic condition associated with the epitope. Thus a toxin conjugate can be administered to kill tumor cells, radiolabeling can facilitate imaging of epitope positive tumor, an enzyme conjugate can be used in an ELISA-like assay to diagnose cancer and confirm epitope expression in biopsied tissue. In a further embodiment, such T cells as set forth above, following expansion accomplished through stimulation with the epitope and/or cytokines, can be administered to a patient as an adoptive immunotherapy.

Reagents Comprising Epitopes

A further aspect of the invention provides isolated epitope-MHC complexes. In a particularly advantageous embodiment of this aspect of the invention, the complexes can be soluble, multimeric proteins such as those described in U. S. Patent No. 5,635,363 (tetramers) or U. S. Patent No. 6,015,884 (Ig-dimers), both of which are hereby incorporated by reference in their entirety. Such reagents are useful in detecting and monitoring specific T cell responses, and in purifying such T cells.

Isolated MHC molecules complexed with epitopic peptides can also be incorporated into planar lipid bilayers or liposomes. Such compositions can be used to stimulate T cells *in vitro* or, in the case of liposomes, *in vivo*. Co-stimulatory molecules (e.g. B7, CD40, LFA-3) can be incorporated into the same compositions or, especially for *in vitro* work, co-stimulation can be provided by anti-co-receptor antibodies (e.g. anti-CD28, anti-CD154, anti-CD2) or cytokines (e.g. IL-2, IL-12). Such stimulation of T cells can constitute vaccination, drive expansion of T cells *in vitro* for subsequent infusion in an immunotherapy, or constitute a step in an assay of T cell function.

The epitope, or more directly its complex with an MHC molecule, can be an important constituent of functional assays of antigen-specific T cells at either an activation or readout step or both. Of the many assays of T cell function current in the art (detailed procedures can be found in standard immunological references such as *Current Protocols in Immunology* 1999 John Wiley & Sons Inc., N.Y., which is hereby incorporated by reference in its entirety) two broad classes can be defined, those that measure the response of a pool of cells and those that measure the response of individual cells. Whereas the former conveys a global measure of the strength of a response, the latter allows determination of the relative frequency of responding cells. Examples of assays measuring global response are cytotoxicity assays, ELISA, and proliferation assays detecting cytokine secretion. Assays measuring the responses of individual cells (or small clones derived from them) include limiting dilution analysis (LDA), ELISPOT, flow cytometric detection of

unsecreted cytokine (described in U.S. Patent No. 5,445,939, entitled "METHOD FOR ASSESSMENT OF THE MONONUCLEAR LEUKOCYTE IMMUNE SYSTEM" and U.S. Patent Nos 5,656,446; and 5,843,689, both entitled "METHOD FOR THE ASSESSMENT OF THE MONONUCLEAR LEUKOCYTE IMMUNE SYSTEM," reagents for which are sold by Becton, Dickinson & Company under the tradename 'FASTIMMUNE', which patents are hereby incorporated by reference in their entirety) and detection of specific TCR with tetramers or Ig-dimers as stated and referenced above. The comparative virtues of these techniques have been reviewed in Yee, C. et al. *Current Opinion in Immunology*, 13:141-146, 2001, which is hereby incorporated by reference in its entirety. Additionally detection of a specific TCR rearrangement or expression can be accomplished through a variety of established nucleic acid based techniques, particularly in situ and single-cell PCR techniques, as will be apparent to one of skill in the art.

These functional assays are used to assess endogenous levels of immunity, response to an immunologic stimulus (e.g. a vaccine), and to monitor immune status through the course of a disease and treatment. Except when measuring endogenous levels of immunity, any of these assays presume a preliminary step of immunization, whether *in vivo* or *in vitro* depending on the nature of the issue being addressed. Such immunization can be carried out with the various embodiments of the invention described above or with other forms of immunogen (e.g., pAPC-tumor cell fusions) that can provoke similar immunity. With the exception of PCR and tetramer/Ig-dimer type analyses which can detect expression of the cognate TCR, these assays generally benefit from a step of *in vitro* antigenic stimulation which can advantageously use various embodiments of the invention as described above in order to detect the particular functional activity (highly cytolytic responses can sometimes be detected directly). Finally, detection of cytolytic activity requires epitope-displaying target cells, which can be generated using various embodiments of the invention. The particular embodiment chosen for any particular step depends on the question to be addressed, ease of use, cost, and the like, but the advantages of one embodiment over another for any particular set of circumstances will be apparent to one of skill in the art.

Tumor Associated Antigens

Epitopes of the present invention are derived from the TuAAs tyrosinase (SEQ ID NO. 2), SSX-2, (SEQ ID NO. 3), PSMA (prostate-specific membrane antigen) (SEQ ID NO. 4), GP100, (SEQ ID NO. 70), MAGE-1, (SEQ ID NO. 71), MAGE-2, (SEQ ID NO. 72), MAGE-3, (SEQ ID NO. 73), NY-ESO-1, (SEQ ID NO. 74), PRAME, (SEQ ID NO. 77), PSA, (SEQ ID NO. 78), and PSCA, (SEQ ID NO. 79). The natural coding sequences for these eleven proteins, or any segments within them, can be determined from their cDNA or complete coding (cds) sequences, SEQ ID NOS. 5-7, and 80-87, respectively.

Tyrosinase is a melanin biosynthetic enzyme that is considered one of the most specific markers of melanocytic differentiation. Tyrosinase is expressed in few cell types, primarily in

melanocytes, and high levels are often found in melanomas. The usefulness of tyrosinase as a TuAA is taught in U.S. Patent 5,747,271 entitled "METHOD FOR IDENTIFYING INDIVIDUALS SUFFERING FROM A CELLULAR ABNORMALITY SOME OF WHOSE ABNORMAL CELLS PRESENT COMPLEXES OF HLA-A2/TYROSINASE DERIVED PEPTIDES, AND METHODS FOR TREATING SAID INDIVIDUALS" which is hereby incorporated by reference in its entirety.

GP100, also known as PMel17, also is a melanin biosynthetic protein expressed at high levels in melanomas. GP100 as a TuAA is disclosed in U.S. Patent 5,844,075 entitled "MELANOMA ANTIGENS AND THEIR USE IN DIAGNOSTIC AND THERAPEUTIC METHODS," which is hereby incorporated by reference in its entirety.

SSX-2, also known as Hom-Mel-40, is a member of a family of highly conserved cancer-testis antigens (Gure, A.O. et al. *Int. J. Cancer* 72:965-971, 1997, which is hereby incorporated by reference in its entirety). Its identification as a TuAA is taught in U.S. Patent 6,025,191 entitled "ISOLATED NUCLEIC ACID MOLECULES WHICH ENCODE A MELANOMA SPECIFIC ANTIGEN AND USES THEREOF," which is hereby incorporated by reference in its entirety. Cancer-testis antigens are found in a variety of tumors, but are generally absent from normal adult tissues except testis. Expression of different members of the SSX family have been found variously in tumor cell lines. Due to the high degree of sequence identity among SSX family members, similar epitopes from more than one member of the family will be generated and able to bind to an MHC molecule, so that some vaccines directed against one member of this family can cross-react and be effective against other members of this family (see example 3 below).

MAGE-1, MAGE-2, and MAGE-3 are members of another family of cancer-testis antigens originally discovered in melanoma (MAGE is a contraction of melanoma-associated antigen) but found in a variety of tumors. The identification of MAGE proteins as TuAAs is taught in U.S. Patent 5,342,774 entitled NUCLEOTIDE SEQUENCE ENCODING THE TUMOR REJECTION ANTIGEN PRECURSOR, MAGE-1, which is hereby incorporated by reference in its entirety, and in numerous subsequent patents. Currently there are 17 entries for (human) MAGE in the SWISS Protein database. There is extensive similarity among these proteins so in many cases, an epitope from one can induce a cross-reactive response to other members of the family. A few of these have not been observed in tumors, most notably MAGE-H1 and MAGE-D1, which are expressed in testes and brain, and bone marrow stromal cells, respectively. The possibility of cross-reactivity on normal tissue is ameliorated by the fact that they are among the least similar to the other MAGE proteins.

NY-ESO-1, is a cancer-testis antigen found in a wide variety of tumors, also known as CTAG-1 (Cancer-Testis Antigen-1) and CAG-3 (Cancer Antigen-3). NY-ESO-1 as a TuAA is disclosed in U.S. Patent 5,804,381 entitled ISOLATED NUCLEIC ACID MOLECULE

ENCODING AN ESOPHAGEAL CANCER ASSOCIATED ANTIGEN, THE ANTIGEN ITSELF,
AND USES THEREOF which is hereby incorporated by reference in its entirety. A paralogous
locus encoding antigens with extensive sequence identity, LAGE-1a/s (SEQ ID NO. 75) and
LAGE-1b/L (SEQ ID NO. 76), have been disclosed in publicly available assemblies of the human
genome, and have been concluded to arise through alternate splicing. Additionally, CT-2 (or
CTAG-2, Cancer-Testis Antigen-2) appears to be either an allele, a mutant, or a sequencing
discrepancy of LAGE-1b/L. Due to the extensive sequence identity, many epitopes from NY-ESO-
1 can also induce immunity to tumors expressing these other antigens. See figure 1. The proteins
are virtually identical through amino acid 70. From 71-134 the longest run of identities between
NY-ESO-1 and LAGE is 6 residues, but potentially cross-reactive sequences are present. And
from 135-180 NY-ESO and LAGE-1a/s are identical except for a single residue, but LAGE-1b/L is
unrelated due to the alternate splice. The CAMEL and LAGE-2 antigens appear to derive from the
LAGE-1 mRNA, but from alternate reading frames, thus giving rise to unrelated protein sequences.
More recently, GenBank Accession AF277315.5, Homo sapiens chromosome X clone RP5-
865E18, RP5-1087L19, complete sequence, reports three independent loci in this region which are
labeled as LAGE1 (corresponding to CTAG-2 in the genome assemblies), plus LAGE2-A and
LAGE2-B (both corresponding to CTAG-1 in the genome assemblies).

PSMA (prostate-specific membranes antigen), a TuAA described in U.S. Patent 5,538,866
entitled "PROSTATE-SPECIFIC MEMBRANES ANTIGEN" which is hereby incorporated by
reference in its entirety, is expressed by normal prostate epithelium and, at a higher level, in
prostatic cancer. It has also been found in the neovasculature of non-prostatic tumors. PSMA can
thus form the basis for vaccines directed to both prostate cancer and to the neovasculature of other
tumors. This later concept is more fully described in a provisional U.S. Patent application No.
60/274,063 entitled ANTI-NEOVASCULAR VACCINES FOR CANCER, filed March 7, 2001,
and U.S. Application No. __/____, attorney docket number CTLIMM.015A, filed on March 7,
2002, entitled "ANTI-NEOVASCULAR PREPARATIONS FOR CANCER," both of which are
hereby incorporated by reference in their entirety. Alternate splicing of the PSMA mRNA also
leads to a protein with an apparent start at Met₅₈, thereby deleting the putative membrane anchor
region of PSMA as described in U.S. Patent 5,935,818 entitled "ISOLATED NUCLEIC ACID
MOLECULE ENCODING ALTERNATIVELY SPLICED PROSTATE-SPECIFIC
MEMBRANES ANTIGEN AND USES THEREOF" which is hereby incorporated by reference in
its entirety. A protein termed PSMA-like protein, Genbank accession number AF261715, is nearly
identical to amino acids 309-750 of PSMA and has a different expression profile. Thus the most
preferred epitopes are those with an N-terminus located from amino acid 58 to 308.

PRAME, also known as MAPE, DAGE, and OIP4, was originally observed as a melanoma
antigen. Subsequently, it has been recognized as a CT antigen, but unlike many CT antigens (e.g.,

MAGE, GAGE, and BAGE) it is expressed in acute myeloid leukemias. PRAME is a member of the MAPE family which consists largely of hypothetical proteins with which it shares limited sequence similarity. The usefulness of PRAME as a TuAA is taught in U.S. Patent 5,830,753 entitled "ISOLATED NUCLEIC ACID MOLECULES CODING FOR TUMOR REJECTION
5 ANTIGEN PRECURSOR DAGE AND USES THEREOF" which is hereby incorporated by reference in its entirety.

PSA, prostate specific antigen, is a peptidase of the kallikrein family and a differentiation antigen of the prostate. Expression in breast tissue has also been reported. Alternate names include gamma-seminoprotein, kallikrein 3, seminogelase, seminin, and P-30 antigen. PSA has a high
10 degree of sequence identity with the various alternate splicing products prostatic/glandular kallikrein-1 and -2, as well as kallikrein 4, which is also expressed in prostate and breast tissue. Other kallikreins generally share less sequence identity and have different expression profiles. Nonetheless, cross-reactivity that might be provoked by any particular epitope, along with the likelihood that that epitope would be liberated by processing in non-target tissues (most generally
15 by the housekeeping proteasome), should be considered in designing a vaccine.

PSCA, prostate stem cell antigen, and also known as SCAH-2, is a differentiation antigen preferentially expressed in prostate epithelial cells, and overexpressed in prostate cancers. Lower level expression is seen in some normal tissues including neuroendocrine cells of the digestive tract and collecting ducts of the kidney. PSCA is described in U.S. Patent 5,856,136 entitled "HUMAN
20 STEM CELL ANTIGENS" which is hereby incorporated by reference in its entirety.

Synaptonemal complex protein 1 (SCP-1), also known as HOM-TES-14, is a meiosis-associated protein and also a cancer-testis antigen (Tureci, O., et al. *Proc. Natl. Acad. Sci. USA* 95:5211-5216, 1998). As a cancer antigen its expression is not cell-cycle regulated and it is found frequently in gliomas, breast, renal cell, and ovarian carcinomas. It has some similarity to myosins,
25 but with few enough identities that cross-reactive epitopes are not an immediate prospect.

The ED-B domain of fibronectin is also a potential target. Fibronectin is subject to developmentally regulated alternative splicing, with the ED-B domain being encoded by a single exon that is used primarily in oncofetal tissues (Matsuura, H. and S. Hakomori *Proc. Natl. Acad. Sci. USA* 82:6517-6521, 1985; Carnemolla, B. et al. *J. Cell Biol.* 108:1139-1148, 1989; Loridon-
30 Rosa, B. et al. *Cancer Res.* 50:1608-1612, 1990; Nicolo, G. et al. *Cell Differ. Dev.* 32:401-408, 1990; Borsi, L. et al. *Exp. Cell Res.* 199:98-105, 1992; Oyama, F. et al. *Cancer Res.* 53:2005-2011, 1993; Mandel, U. et al. *APMIS* 102:695-702, 1994; Farnoud, M.R. et al. *Int. J. Cancer* 61:27-34, 1995; Pujuguet, P. et al. *Am. J. Pathol.* 148:579-592, 1996; Gabler, U. et al. *Heart* 75:358-362, 1996; Chevalier, X. *Br. J. Rheumatol.* 35:407-415, 1996; Midulla, M. *Cancer Res.* 60:164-169, 35 2000).

The ED-B domain is also expressed in fibronectin of the neovasculature (Kaczmarek, J. et al. *Int. J. Cancer* 59:11-16, 1994; Castellani, P. et al. *Int. J. Cancer* 59:612-618, 1994; Neri, D. et al. *Nat. Biotech.* 15:1271-1275, 1997; Karelina, T.V. and A.Z. Eisen *Cancer Detect. Prev.* 22:438-444, 1998; Tarli, L. et al. *Blood* 94:192-198, 1999; Castellani, P. et al. *Acta Neurochir. (Wien)* 142:277-282, 2000). As an oncofetal domain, the ED-B domain is commonly found in the fibronectin expressed by neoplastic cells in addition to being expressed by the neovasculature. Thus, CTL-inducing vaccines targeting the ED-B domain can exhibit two mechanisms of action: direct lysis of tumor cells, and disruption of the tumor's blood supply through destruction of the tumor-associated neovasculature. As CTL activity can decay rapidly after withdrawal of vaccine, interference with normal angiogenesis can be minimal. The design and testing of vaccines targeted to neovasculature is described in Provisional U.S. Patent Application No. 60/274,063 entitled "ANTI-NEOVASCULATURE VACCINES FOR CANCER" and in U.S. Patent Application No. ____/____, attorney docket number CTLIMM.015A, entitled "ANTI-NEOVASCULATURE PREPARATIONS FOR CANCER, filed on date even with this application (March 7, 2002). A tumor cell line is disclosed in Provisional U.S. Application No. ____/____, filed on March 7, 2002, attorney docket number CTLIMM.028PR, entitled "HLA-TRANSGENIC MURINE TUMOR CELL LINE," which is hereby incorporated by reference in its entirety.

Carcinoembryonic antigen (CEA) is a paradigmatic oncofetal protein first described in 1965 (Gold and Freedman, *J. Exp. Med.* 121: 439-462, 1965. Fuller references can be found in the Online Medelian Inheritance in Man; record *114890). It has officially been renamed carcinoembryonic antigen-related cell adhesion molecule 5 (CEACAM5). Its expression is most strongly associated with adenocarcinomas of the epithelial lining of the digestive tract and in fetal colon. CEA is a member of the immunoglobulin supergene family and the defining member of the CEA subfamily.

HER2/NEU is an oncogene related to the epidermal growth factor receptor (van de Vijver, et al., *New Eng. J. Med.* 319:1239-1245, 1988), and apparently identical to the c-ERBB2 oncogene (Di Fiore, et al., *Science* 237: 178-182, 1987). The over-expression of ERBB2 has been implicated in the neoplastic transformation of prostate cancer. As HER2 it is amplified and over-expressed in 25-30% of breast cancers among other tumors where expression level is correlated with the aggressiveness of the tumor (Slamon, et al., *New Eng. J. Med.* 344:783-792, 2001). A more detailed description is available in the Online Medelian Inheritance in Man; record *164870.

All references mentioned herein are hereby incorporated by reference in their entirety. Further, incorporated by reference in its entirety is U.S. Patent Application No. 10/005,905 (attorney docket number CTLIMM.021CP1) entitled "EPI TOPE SYNCHRONIZATION IN ANTIGEN PRESENTING CELLS," filed on November 7, 2001 and a continuation thereof, U.S. Application No. ____/____, filed on December 7, 2000, attorney docket number

CTLIMM.21CP1C, also entitled "EPITOPE SYNCHRONIZATION IN ANTIGEN PRESENTING CELLS."

Useful epitopes were identified and tested as described in the following examples. However, these examples are intended for illustration purposes only, and should not be construed as limiting the scope of the invention in any way.

EXAMPLES

Sequences of Specific Preferred Epitopes

Example 1

Manufacture of tyrosinase epitopes.

A. Synthetic production of epitopes

Peptides having an amino acid sequence of any of SEQ ID NO: 1, 8, 9, 11-23, 2-29, 32-44, 47-54, 56-63, 66-68 88-253, or 256-588 are synthesized using either FMOC or tBOC solid phase synthesis methodologies. After synthesis, the peptides are cleaved from their supports with either trifluoroacetic acid or hydrogen fluoride, respectively, in the presence of appropriate protective scavengers. After removing the acid by evaporation, the peptides are extracted with ether to remove the scavengers and the crude, precipitated peptide is then lyophilized. Purity of the crude peptides is determined by HPLC, sequence analysis, amino acid analysis, counterion content analysis and other suitable means. If the crude peptides are pure enough (greater than or equal to about 90% pure), they can be used as is. If purification is required to meet drug substance specifications, the peptides are purified using one or a combination of the following: re-precipitation; reverse-phase, ion exchange, size exclusion or hydrophobic interaction chromatography; or counter-current distribution.

Drug product formulation

GMP-grade peptides are formulated in a parenterally acceptable aqueous, organic, or aqueous-organic buffer or solvent system in which they remain both physically and chemically stable and biologically potent. Generally, buffers or combinations of buffers or combinations of buffers and organic solvents are appropriate. The pH range is typically between 6 and 9. Organic modifiers or other excipients can be added to help solubilize and stabilize the peptides. These include detergents, lipids, co-solvents, antioxidants, chelators and reducing agents. In the case of a lyophilized product, sucrose or mannitol or other lyophilization aids can be added. Peptide solutions are sterilized by membrane filtration into their final container-closure system and either lyophilized for dissolution in the clinic, or stored until use.

B. Construction of expression vectors for use as nucleic acid vaccines

The construction of three generic epitope expression vectors is presented below. The particular advantages of these designs are set forth in U.S. Patent Application No. 09/561,572

entitled "EXPRESSION VECTORS ENCODING EPITOPES OF TARGET-ASSOCIATED ANTIGENS," which has been incorporated by reference in its entirety above.

A suitable *E. coli* strain was then transfected with the plasmid and plated out onto a selective medium. Several colonies were grown up in suspension culture and positive clones were identified by restriction mapping. The positive clone was then grown up and aliquotted into storage vials and stored at -70°C.

A mini-prep (QIAprep Spin Mini-prep: Qiagen, Valencia, CA) of the plasmid was then made from a sample of these cells and automated fluorescent dideoxy sequence analysis was used to confirm that the construct had the desired sequence.

B.1 Construction of pVAX-EP1-IRES-EP2

Overview:

The starting plasmid for this construct is pVAX1 purchased from Invitrogen (Carlsbad, CA). Epitopes EP1 and EP2 were synthesized by GIBCO BRL (Rockville, MD). The IRES was excised from pIRES purchased from Clontech (Palo Alto, CA).

Procedure:

- 1 pIRES was digested with EcoRI and NotI. The digested fragments were separated by agarose gel electrophoresis, and the IRES fragment was purified from the excised band.
- 2 pVAX1 was digested with EcoRI and NotI, and the pVAX1 fragment was gel-purified.
- 3 The purified pVAX1 and IRES fragments were then ligated together.
- 4 Competent *E. coli* of strain DH5α were transformed with the ligation mixture.
- 5 Minipreps were made from 4 of the resultant colonies.
- 6 Restriction enzyme digestion analysis was performed on the miniprep DNA. One recombinant colony having the IRES insert was used for further insertion of EP1 and EP2. This intermediate construct was called pVAX-IRES.
- 7 Oligonucleotides encoding EP1 and EP2 were synthesized.
- 8 EP1 was subcloned into pVAX-IRES between AflII and EcoRI sites, to make pVAX-EP1-IRES;
- 9 EP2 was subcloned into pVAX-EP1-IRES between SalI and NotI sites, to make the final construct pVAX-EP1-IRES-EP2.
- 10 The sequence of the EP1-IRES-EP2 insert was confirmed by DNA sequencing.

B 2. Construction of pVAX-EP1-IRES-EP2-ISS-NIS

Overview:

The starting plasmid for this construct was pVAX-EP1-IRES-EP2 (Example 1). The ISS (immunostimulatory sequence) introduced into this construct is AACGTT, and the NIS (standing for nuclear import sequence) used is the SV40 72bp repeat sequence. ISS-NIS was synthesized by GIBCO BRL. See Figure 2.

Procedure:

- 1 pVAX-EP1-IRES-EP2 was digested with NruI; the linearized plasmid was gel-purified.
- 2 ISS-NIS oligonucleotide was synthesized.
- 3 The purified linearized pVAX-EP1-IRES-EP2 and synthesized ISS-NIS were ligated
5 together.
- 4 Competent E. coli of strain DH5 α were transformed with the ligation product.
- 5 Minipreps were made from resultant colonies.
- 6 Restriction enzyme digestions of the minipreps were carried out.
- 7 The plasmid with the insert was sequenced.

10 B3. Construction of pVAX-EP2-UB-EP1**Overview:**

The starting plasmid for this construct was pVAX1 (Invitrogen). EP2 and EP1 were synthesized by GIBCO BRL. Wild type Ubiquitin cDNA encoding the 76 amino acids in the construct was cloned from yeast.

15 Procedure:

- 1 RT-PCR was performed using yeast mRNA. Primers were designed to amplify the complete coding sequence of yeast Ubiquitin.
- 2 The RT-PCR products were analyzed using agarose gel electrophoresis. A band with the predicted size was gel-purified.
- 20 3 The purified DNA band was subcloned into pZERO1 at EcoRV site. The resulting clone was named pZERO-UB.
- 4 Several clones of pZERO-UB were sequenced to confirm the Ubiquitin sequence before further manipulations.
- 5 EP1 and EP2 were synthesized.
- 25 6 EP2, Ubiquitin and EP1 were ligated and the insert cloned into pVAX1 between BamHI and EcoRI, putting it under control of the CMV promoter.
- 7 The sequence of the insert EP2-UB-EP1 was confirmed by DNA sequencing.

Example 2**Identification of useful epitope variants.**

- 30 The 10-mer FLPWHRLFL (SEQ ID NO. 1) is identified as a useful epitope. Based on this sequence, numerous variants are made. Variants exhibiting activity in HLA binding assays (see Example 3, section 6) are identified as useful, and are subsequently incorporated into vaccines.

The HLA-A2 binding of length variants of FLPWHRLFL have been evaluated. Proteasomal digestion analysis indicates that the C-terminus of the 9-mer FLPWHRLFL (SEQ ID
35 NO. 8) is also produced. Additionally the 9-mer LPWHRLFL (SEQ ID NO. 9) can result from N-terminal trimming of the 10-mer. Both are predicted to bind to the HLA-A*0201 molecule,

however of these two 9-mers, FLPWHRLFL displayed more significant binding and is preferred (see Figs. 3A and B).

Sequence variants of FLPWHRLFL are constructed as follow. Consistent with the binding coefficient table (see Table 3) from the NIH/BIMAS MHC binding prediction program (see reference in example 3 below), binding can be improved by changing the L at position 9, an anchor position, to V. Binding can also be altered, though generally to a lesser extent, by changes at non-anchor positions. Referring generally to Table 3, binding can be increased by employing residues with relatively larger coefficients. Changes in sequence can also alter immunogenicity independently of their effect on binding to MHC. Thus binding and/or immunogenicity can be improved as follows:

By substituting F, L, M, W, or Y for P at position 3; these are all bulkier residues that can also improve immunogenicity independent of the effect on binding. The amine and hydroxyl-bearing residues, Q and N; and S and T; respectively, can also provoke a stronger, cross-reactive response.

By substituting D or E for W at position 4 to improve binding; this addition of a negative charge can also make the epitope more immunogenic, while in some cases reducing cross-reactivity with the natural epitope. Alternatively the conservative substitutions of F or Y can provoke a cross-reactive response.

By substituting F for H at position 5 to improve binding. H can be viewed as partially charged, thus in some cases the loss of charge can hinder cross-reactivity. Substitution of the fully charged residues R or K at this position can enhance immunogenicity without disrupting charge-dependent cross-reactivity.

By substituting I, L, M, V, F, W, or Y for R at position 6. The same caveats and alternatives apply here as at position 5.

By substituting W or F for L at position 7 to improve binding. Substitution of V, I, S, T, Q, or N at this position are not generally predicted to reduce binding affinity by this model (the NIH algorithm), yet can be advantageous as discussed above.

Y and W, which are equally preferred as the Fs at positions 1 and 8, can provoke a useful cross-reactivity. Finally, while substitutions in the direction of bulkiness are generally favored to improve immunogenicity, the substitution of smaller residues such as A, S, and C, at positions 3-7 can be useful according to the theory that contrast in size, rather than bulkiness per se, is an important factor in immunogenicity. The reactivity of the thiol group in C can introduce other properties as discussed in Chen, J.-L., et al. *J. Immunol.* 165:948-955, 2000.

Table 3. 9-mer Coefficient Table for HLA-A*0201*

HLA Coefficient table for file "A_0201_standard"									
Amino Acid Type	1 st	2 nd	3 rd	4 th	5 th	6 th	7 th	8 th	9 th
A	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
C	1.000	0.470	1.000	1.000	1.000	1.000	1.000	1.000	
D	0.075	0.100	0.400	4.100	1.000	1.000	0.490	1.000	
E	0.075	1.400	0.064	4.100	1.000	1.000	0.490	1.000	
F	4.600	0.050	3.700	1.000	3.800	1.900	5.800	5.500	
G	1.000	0.470	1.000	1.000	1.000	1.000	0.130	1.000	
H	0.034	0.050	1.000	1.000	1.000	1.000	1.000	1.000	
I	1.700	9.900	1.000	1.000	1.000	2.300	1.000	0.410	
K	3.500	0.100	0.035	1.000	1.000	1.000	1.000	1.000	
L	1.700	72.000	3.700	1.000	1.000	2.300	1.000	1.000	
M	1.700	52.000	3.700	1.000	1.000	2.300	1.000	1.000	
N	1.000	0.470	1.000	1.000	1.000	1.000	1.000	1.000	
P	0.022	0.470	1.000	1.000	1.000	1.000	1.000	1.000	
Q	1.000	7.300	1.000	1.000	1.000	1.000	1.000	1.000	
R	1.000	0.010	0.076	1.000	1.000	1.000	0.200	1.000	
S	1.000	0.470	1.000	1.000	1.000	1.000	1.000	1.000	
T	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
V	1.700	6.300	1.000	1.000	1.000	2.300	1.000	0.410	1
W	4.600	0.010	8.300	1.000	1.000	1.700	7.500	5.500	
Y	4.600	0.010	3.200	1.000	1.000	1.500	1.000	5.500	

*This table and other comparable data that are publicly available are useful in designing epitope variants and in determining whether a particular variant is substantially similar, or is functionally similar.

Example 3

Cluster Analysis (SSX-2₃₁₋₆₈).

1. Epitope cluster region prediction:

The computer algorithms: SYFPEITHI (internet [http:// access at syfpeithi.bmi-heidelberg.com/Scripts/MHCServer.dll/EpPredict.htm](http://access.at.syfpeithi.bmi-heidelberg.com/Scripts/MHCServer.dll/EpPredict.htm)), based on the book "MHC Ligands and Peptide Motifs" by H.G.Rammensee, J.Bachmann and S.Stevanovic; and HLA Peptide Binding Predictions (NIH) (internet [http:// access at bimas.dcrf.nih.gov/molbio/hla_bin](http://access.at.bimas.dcrf.nih.gov/molbio/hla_bin)), described in Parker, K. C., et al., *J. Immunol.* 152:163, 1994; were used to analyze the protein sequence of SSX-2 (GI:10337583). Epitope clusters (regions with higher than average density of peptide fragments with high predicted MHC affinity) were defined as described fully in U.S. Patent Application No. 09/561,571 entitled "EPITOPE CLUSTERS," filed on April 28, 2000. Using a epitope density ratio cutoff of 2, five and two clusters were defined using the SYFPETHI and NIH algorithms, respectively, and peptides score cutoffs of 16 (SYFPETHI) and 5 (NIH). The highest scoring peptide with the NIH algorithm, SSX-2₄₁₋₄₉, with an estimated halftime of dissociation of

>1000 min., does not overlap any other predicted epitope but does cluster with SSX-2₅₇₋₆₅ in the NIH analysis.

2. Peptide synthesis and characterization:

SSX-2₃₁₋₆₈, YFSKEEWEKMKASEKIFYVYMKRKYEAMTKLGFKATLP (SEQ ID. NO. 10) was synthesized by MPS (Multiple Peptide Systems, San Diego, CA 92121) using standard solid phase chemistry. According to the provided 'Certificate of Analysis', the purity of this peptide was 95%.

3. Proteasome digestion:

Proteasome was isolated from human red blood cells using the proteasome isolation protocol described in U.S. Patent Application No. 09/561,074 entitled "METHOD OF EPITOPE DISCOVERY," filed on April 28, 2000. SDS-PAGE, western-blotting, and ELISA were used as quality control assays. The final concentration of proteasome was 4 mg/ml, which was determined by non-interfering protein assay (Geno Technologies Inc.). Proteasomes were stored at -70°C in 25 µl aliquots.

SSX-2₃₁₋₆₈ was dissolved in Milli-Q water, and a 2 mM stock solution prepared and 20µL aliquots stored at -20°C.

1 tube of proteasome (25 µL) was removed from storage at -70°C and thawed on ice. It was then mixed thoroughly with 12.5µL of 2mM peptide by repipetting (samples were kept on ice). A 5µL sample was immediately removed after mixing and transferred to a tube containing 1.25µL 10%TFA (final concentration of TFA was 2%); the T=0 min sample. The proteasome digestion reaction was then started and carried out at 37°C in a programmable thermal controller. Additional 5µL samples were taken out at 15, 30, 60, 120, 180 and 240 min respectively, the reaction was stopped by adding the sample to 1.25µL 10% TFA as before. Samples were kept on ice or frozen until being analyzed by MALDI-MS. All samples were saved and stored at -20°C for HPLC analysis and N-terminal sequencing. Peptide alone (without proteasome) was used as a blank control: 2 µL peptide + 4µL Tris buffer (20 mM, pH 7.6) + 1.5µL TFA.

4. MALDI-TOF MS measurements:

For each time point 0.3 µL of matrix solution (10mg/ml α-cyano-4-hydroxycinnamic acid in AcCN/H₂O (70:30)) was first applied on a sample slide, and then an equal volume of digested sample was mixed gently with matrix solution on the slide. The slide was allowed to dry at ambient air for 3-5 min. before acquiring the mass spectra. MS was performed on a Lasermat 2000 MALDI-TOF mass spectrometer that was calibrated with peptide/protein standards. To improve the accuracy of measurement, the molecular ion weight (MH⁺) of the peptide substrate was used as an internal calibration standard. The mass spectrum of the T=120 min. digested sample is shown in figure 4.

5. MS data analysis and epitope identification:

To assign the measured mass peaks, the computer program MS-Product, a tool from the UCSF Mass Spectrometry Facility (<http://prospector.ucsf.edu/ucsfhtml3.4/msprod.htm>), was used to generate all possible fragments (N- and C-terminal ions, and internal fragments) and their corresponding molecular weights. Due to the sensitivity of the mass spectrometer, average molecular weight was used. The mass peaks observed over the course of the digestion were identified as summarized in Table 4.

Fragments co-C-terminal with 8-10 amino acid long sequences predicted to bind HLA by the SYFPEITHI or NIH algorithms were chosen for further study. The digestion and prediction steps of the procedure can be usefully practiced in any order. Although the substrate peptide used in proteasomal digest described here was specifically designed to include predicted HLA-A2.1 binding sequences, the actual products of digestion can be checked after the fact for actual or predicted binding to other MHC molecules. Selected results are shown in Table 5.

Table 4. SSX-2₃₁₋₆₈ Mass Peak Identification.

MS PEAK (measured)	PEPTIDE	SEQUENCE	CALCULATED MASS (MH ⁺)
988.23	31-37	YFSKEEW	989.08
1377.68±2.3 8	31-40	YFSKEEWEKM	1377.68
1662.45±1.3 0	31-43	YFSKEEWEKMKAS	1663.90
2181.72±0.8 5	31-47	YFSKEEWEKMKASEKIF	2181.52
2346.6	31-48	YFSKEEWEKMKASEKIFY	2344.71
1472.16±1.5 4	38-49	EKMASEKIFYV	1473.77
2445.78±1.1 8	31-49*	YFSKEEWEKMKASEKIFYV	2443.84
2607.	31-50	YFSKEEWEKMKASEKIFYVY	2607.02
1563.3	50-61	YMKRKYEAMTKL	1562.93
3989.9	31-61	YFSKEEWEKMKASEKIFYVYMKRKYEAMTKL	3987.77
1603.74±1.5 3	51-63	MKRKYEAMTKLGF	1603.98
1766.45±1.5	50-63	YMKRKYEAMTKLGF	1767.16
1866.32±1.2 2	49-63	VYMKRKYEAMTKLGF	1866.29
4192.6	31-63	YFSKEEWEKMKASEKIFYVYMKRKYEAMTKLG F	4192.00
4392.1	31-65**	YFSKEEWEKMKASEKIFYVYMKRKYEAMTKLG FKA	4391.25

Boldface sequence correspond to peptides predicted to bind to MHC.

* On the basis of mass alone this peak could also have been assigned to the peptide 32-50, however proteasomal removal of just the N-terminal amino acid is unlikely. N-terminal sequencing (below) verifies the assignment to 31-49.

** On the basis of mass this fragment might also represent 33-68. N-terminal sequencing below is consistent with the assignment to 31-65.

Table 5. Predicted HLA binding by proteasomally generated fragments

<u>SEQ ID NO.</u>	<u>PEPTIDE</u>	<u>HLA</u>	<u>SYFPEITHI</u>	<u>NIH</u>
11	FSKEEWKEM	B*3501	NP†	90
12	KMKASEKIF	B*08	17	<5
13 & (14)	(K) MKASEKIFY	A1	19 (19)	<5
15 & (16)	(M) KASEKIFYV	A*0201	22 (16)	1017
		B*08	17	<5
		B*5101	22 (13)	60
		B*5102	NP	133
		B*5103	NP	121
17 & (18)	(K) ASEKIFYVY	A1	34 (19)	14
19 & (20)	(K) RKYEAMTKL	A*0201	15	<5
		A26	15	NP
		B14	NP	45 (60)
		B*2705	21	15
		B*2709	16	NP
21	KYEAMTKLGF	B*5101	15	<5
		A1	16	<5
		A24	NP	300
		B*4403	NP	80
22	YEAMTKLGF	B*08	22	<5
23	EAMTKLGF			

†No prediction

5

As seen in Table 5, N-terminal addition of authentic sequence to epitopes can generate epitopes for the same or different MHC restriction elements. Note in particular the pairing of (K)RKYEAMTKL (SEQ ID NOS 19 and (20)) with HLA-B14, where the 10-mer has a longer predicted halftime of dissociation than the co-C-terminal 9-mer. Also note the case of the 10-mer KYEAMTKLGF (SEQ ID NO. 21) which can be used as a vaccine useful with several MHC types by relying on N-terminal trimming to create the epitopes for HLA-B*4403 and -B*08.

10

6. HLA-A0201 binding assay:

15

Binding of the candidate epitope KASEKIFYV, SSX-2₄₁₋₄₉, (SEQ ID NO. 15) to HLA-A2.1 was assayed using a modification of the method of Stauss et al., (Proc Natl Acad Sci USA 89(17):7871-5 (1992)). Specifically, T2 cells, which express empty or unstable MHC molecules on their surface, were washed twice with Iscove's modified Dulbecco's medium (IMDM) and cultured overnight in serum-free AIM-V medium (Life Technologies, Inc., Rockville, MD) supplemented with human β 2-microglobulin at 3 μ g/ml (Sigma, St. Louis, MO) and added peptide,

at 800, 400, 200, 100, 50, 25, 12.5, and 6.25 $\mu\text{g/ml}$ in a 96-well flat-bottom plate at 3×10^5 cells/200 μl /well. Peptide was mixed with the cells by repipeting before distributing to the plate (alternatively peptide can be added to individual wells), and the plate was rocked gently for 2 minutes. Incubation was in a 5% CO_2 incubator at 37°C . The next day the unbound peptide was removed by washing twice with serum free RPMI medium and a saturating amount of anti-class I HLA monoclonal antibody, fluorescein isothiocyanate (FITC)-conjugated anti-HLA A2, A28 (One Lambda, Canoga Park, CA) was added. After incubation for 30 minutes at 4°C , cells were washed 3 times with PBS supplemented with 0.5% BSA, 0.05% (w/v) sodium azide, pH 7.4-7.6 (staining buffer). (Alternatively W6/32 (Sigma) can be used as the anti-class I HLA monoclonal antibody the cells washed with staining buffer and then incubated with fluorescein isothiocyanate (FITC)-conjugated goat F(ab') antimouse-IgG (Sigma) for 30 min at 4°C and washed 3 times as before.) The cells were resuspended in 0.5 ml staining buffer. The analysis of surface HLA-A2.1 molecules stabilized by peptide binding was performed by flow cytometry using a FACScan (Becton Dickinson, San Jose, CA). If flow cytometry is not to be performed immediately the cells can be fixed by adding a quarter volume of 2% paraformaldehyde and storing in the dark at 4°C .

The results of the experiment are shown in Figure 5. SSX-2₄₁₋₄₉ (SEQ ID NO. 15) was found to bind HLA-A2.1 to a similar extent as the known A2.1 binder FLPSDYFPSV (HBV₁₈₋₂₇; SEQ ID NO: 24) used as a positive control. An HLA-B44 binding peptide, AEMGKYSFY (SEQ ID NO: 25), was used as a negative control. The fluorescence obtained from the negative control was similar to the signal obtained when no peptide was used in the assay. Positive and negative control peptides were chosen from Table 18.3.1 in *Current Protocols in Immunology* p. 18.3.2, John Wiley and Sons, New York, 1998.

7. Immunogenicity:

A. In vivo immunization of mice.

HHD1 transgenic A*0201 mice (Pascolo, S., et al. *J. Exp. Med.* 185:2043-2051, 1997) were anesthetized and injected subcutaneously at the base of the tail, avoiding lateral tail veins, using 100 μl containing 100 nmol of SSX-2₄₁₋₄₉ (SEQ ID NO. 15) and 20 μg of HTL epitope peptide in PBS emulsified with 50 μl of IFA (incomplete Freund's adjuvant).

B. Preparation of stimulating cells (LPS blasts).

Using spleens from 2 naive mice for each group of immunized mice, un-immunized mice were sacrificed and the carcasses were placed in alcohol. Using sterile instruments, the top dermal layer of skin on the mouse's left side (lower mid-section) was cut through, exposing the peritoneum. The peritoneum was saturated with alcohol, and the spleen was aseptically extracted. The spleen was placed in a petri dish with serum-free media. Splenocytes were isolated by using sterile plungers from 3 ml syringes to mash the spleens. Cells were collected in a 50 ml conical tubes in serum-free media, rinsing dish well. Cells were centrifuged (12000 rpm, 7 min) and

washed one time with RPMI. Fresh spleen cells were resuspended to a concentration of 1×10^6 cells per ml in RPMI-10%FCS (fetal calf serum). 25g/ml lipopolysaccharide and 7 μ g/ml Dextran Sulfate were added. Cell were incubated for 3 days in T-75 flasks at 37°C, with 5% CO₂. Splenic blasts were collected in 50 ml tubes pelleted (12000 rpm, 7 min) and resuspended to 3×10^7 /ml in RPMI. The blasts were pulsed with the priming peptide at 50 μ g/ml, RT 4hr. mitomycin C-treated at 25 μ g/ml, 37°C, 20 min and washed three times with DMEM.

C. In vitro stimulation.

3 days after LPS stimulation of the blast cells and the same day as peptide loading, the primed mice were sacrificed (at 14 days post immunization) to remove spleens as above. 3×10^6 splenocytes were co-cultured with 1×10^6 LPS blasts/well in 24-well plates at 37°C, with 5% CO₂ in DMEM media supplemented with 10% FCS, 5×10^{-5} M β -mercaptoethanol, 100 μ g/ml streptomycin and 100 IU/ml penicillin. Cultures were fed 5% (vol/vol) ConA supernatant on day 3 and assayed for cytolytic activity on day 7 in a ⁵¹Cr-release assay.

D. Chromium-release assay measuring CTL activity.

To assess peptide specific lysis, 2×10^6 T2 cells were incubated with 100 μ Ci sodium chromate together with 50 μ g/ml peptide at 37 C for 1 hour. During incubation they were gently shaken every 15 minutes. After labeling and loading, cells were washed three times with 10 ml of DMEM-10% FCS, wiping each tube with a fresh Kimwipe after pouring off the supernatant. Target cells were resuspended in DMEM-10% FBS 1×10^5 /ml. Effector cells were adjusted to 1×10^7 /ml in DMEM-10% FCS and 100 μ l serial 3-fold dilutions of effectors were prepared in U-bottom 96-well plates. 100 μ l of target cells were added per well. In order to determine spontaneous release and maximum release, six additional wells containing 100 μ l of target cells were prepared for each target. Spontaneous release was revealed by incubating the target cells with 100 μ l medium; maximum release was revealed by incubating the target cells with 100 μ l of 2% SDS. Plates were then centrifuged for 5 min at 600 rpm and incubated for 4 hours at 37°C in 5% CO₂ and 80% humidity. After the incubation, plates were then centrifuged for 5 min at 1200 rpm. Supernatants were harvested and counted using a gamma counter. Specific lysis was determined as follows: % specific release = [(experimental release - spontaneous release)/(maximum release - spontaneous release)] x 100.

Results of the chromium release assay demonstrating specific lysis of peptide pulsed target cells are shown in figure 6.

8. Cross-reactivity with other SSX proteins:

SSX-2₄₁₋₄₉ (SEQ ID NO. 15) shares a high degree of sequence identity with the same region of the other SSX proteins. The surrounding regions have also been generally well conserved. Thus the housekeeping proteasome can cleave following V₄₉ in all five sequences. Moreover, SSX₄₁₋₄₉ is

predicted to bind HLA-A*0201 (see Table 6). CTL generated by immunization with SSX-2₄₁₋₄₉ cross-react with tumor cells expressing other SSX proteins.

Table 6. SSX₄₁₋₄₉ – A*0201 Predicted Binding

SEQ ID NO.	Family Member	Sequence	SYFPEITHI Score	NIH Score
15	SSX-2	KASEKIFYV	22	1017
26	SSX-1	KYSEKISYV	18	1.7
27	SSX-3	KVSEKIVYV	24	1105
28	SSX-4	KSSEKIVYV	20	82
29	SSX-5	KASEKIYV	22	175

5 Example 4

Cluster Analysis (PSMA₁₆₃₋₁₉₂).

A peptide, AFSPQGMPEGDLVYVNYARTEDFFKLERDM, PSMA₁₆₃₋₁₉₂, (SEQ ID NO. 30), containing an A1 epitope cluster from prostate specific membrane antigen, PSMA₁₆₈₋₁₉₀ (SEQ ID NO. 31) was synthesized using standard solid-phase F-moc chemistry on a 433A ABI Peptide synthesizer. After side chain deprotection and cleavage from the resin, peptide first dissolved in formic acid and then diluted into 30% Acetic acid, was run on a reverse-phase preparative HPLC C4 column at following conditions: linear AB gradient (5% B/min) at a flow rate of 4 ml/min, where eluent A is 0.1% aqueous TFA and eluent B is 0.1% TFA in acetonitrile. A fraction at time 16.642 min containing the expected peptide, as judged by mass spectrometry, was pooled and lyophilized. The peptide was then subjected to proteasome digestion and mass spectrum analysis essentially as described above. Prominent peaks from the mass spectra are summarized in Table 7.

Table 7. PSMA₁₆₃₋₁₉₂ Mass Peak Identification.

PEPTIDE	SEQUENCE	CALCULATED MASS (MH ⁺)
163-177	AFSPQGMPEGDLVYV	1610.0
178-189	NYARTEDFFKLE	1533.68
170-189	PEGDLVYVNYARTEDFFKLE	2406.66
178-191	NYARTEDFFKLERD	1804.95
170-191	PEGDLVYVNYARTEDFFKLERD	2677.93
178-192	NYARTEDFFKLERDM	1936.17
163-176	AFSPQGMPEGDLVY	1511.70
177-192	VNYARTEDFFKLERDM	2035.30
163-179	AFSPQGMPEGDLVYVNY	1888.12

180-192	ARTEDFFKLERDM	1658.89
163-183	AFSPQGMPEGDLVYVNYARTE	2345.61
184-192	DFFKLERDM	1201.40
176-192	YVNYARTEDFFKLERDM	2198.48
167-185	QGMPEGDLVYVNYARTEDF	2205.41
178-186	NYARTEDF	1163.22

Boldface sequences correspond to peptides predicted to bind to MHC, see Table 8.

N-terminal Pool Sequence Analysis

One aliquot at one hour of the proteasomal digestion (see Example 3 part 3 above) was subjected to N-terminal amino acid sequence analysis by an ABI 473A Protein Sequencer (Applied Biosystems, Foster City, CA). Determination of the sites and efficiencies of cleavage was based on consideration of the sequence cycle, the repetitive yield of the protein sequencer, and the relative yields of amino acids unique in the analyzed sequence. That is if the unique (in the analyzed sequence) residue X appears only in the nth cycle a cleavage site exists n-1 residues before it in the N-terminal direction. In addition to helping resolve any ambiguity in the assignment of mass to sequences, these data also provide a more reliable indication of the relative yield of the various fragments than does mass spectrometry.

For PSMA₁₆₃₋₁₉₂ (SEQ ID NO. 30) this pool sequencing supports a single major cleavage site after V₁₇₇ and several minor cleavage sites, particularly one after Y₁₇₉. Reviewing the results presented in figures 7A-C reveals the following:

- S at the 3rd cycle indicating presence of the N-terminus of the substrate.
- Q at the 5th cycle indicating presence of the N-terminus of the substrate.
- N at the 1st cycle indicating cleavage after V₁₇₇.
- N at the 3rd cycle indicating cleavage after V₁₇₅. Note the fragment 176-192 in Table 7.
- T at the 5th cycle indicating cleavage after V₁₇₇.
- T at the 1st–3rd cycles, indicating increasingly common cleavages after R₁₈₁, A₁₈₀ and Y₁₇₉.
Only the last of these correspond to peaks detected by mass spectrometry; 163-179 and 180-192, see Table 7. The absence of the others can indicate that they are on fragments smaller than were examined in the mass spectrum.
- K at the 4th, 8th, and 10th cycles indicating cleavages after E₁₈₃, Y₁₇₉, and V₁₇₇, respectively, all of which correspond to fragments observed by mass spectroscopy. See Table 7.
- A at the 1st and 3rd cycles indicating presence of the N-terminus of the substrate and cleavage after V₁₇₇, respectively.
- P at the 4th and 8th cycles indicating presence of the N-terminus of the substrate.

G at the 6th and 10th cycles indicating presence of the N-terminus of the substrate.

M at the 7th cycle indicating presence of the N-terminus of the substrate and/or cleavage after F₁₈₅.

M at the 15th cycle indicating cleavage after V₁₇₇.

5 The 1st cycle can indicate cleavage after D₁₉₁, see Table 7.

R at the 4th and 13th cycle indicating cleavage after V₁₇₇.

R at the 2nd and 11th cycle indicating cleavage after Y₁₇₉.

V at the 2nd, 6th, and 13th cycle indicating cleavage after V₁₇₅, M₁₆₉ and presence of the N-terminus of the substrate, respectively. Note fragments beginning at 176 and 170 in
10 Table 7.

Y at the 1st, 2nd, and 14th cycles indicating cleavage after V₁₇₅, V₁₇₇, and presence of the N-terminus of the substrate, respectively.

L at the 11th and 12th cycles indicating cleavage after V₁₇₇, and presence of the N-terminus of the substrate, respectively, is the interpretation most consistent with the other data.

15 Comparing to the mass spectrometry results we see that L at the 2nd, 5th, and 9th cycles is consistent with cleavage after F₁₈₆, E₁₈₃ or M₁₆₉, and Y₁₇₉, respectively. See Table 7.

Epitope Identification

20 Fragments co-C-terminal with 8-10 amino acid long sequences predicted to bind HLA by the SYFPEITHI or NIH algorithms were chosen for further analysis. The digestion and prediction steps of the procedure can be usefully practiced in any order. Although the substrate peptide used in proteasomal digest described here was specifically designed to include a predicted HLA-A1 binding sequence, the actual products of digestion can be checked after the fact for actual or predicted binding to other MHC molecules. Selected results are shown in Table 8.

Table 8. Predicted HLA binding by proteasomally generated fragments

SEQ ID NO	PEPTIDE	HLA	SYFPEITHI	NIH
32 & (33)	(G)MPEGDLVY V	A*0201	17 (27)	(2605)
		B*0702	20	<5
		B*5101	22	314
34 & (35)	(Q)GMPEGDLV Y	A1	24 (26)	<5
		A3	16 (18)	36
		B*2705	17	25
	MPEGDLVY	B*5101	15	NP†
36	(P)EGDLVYVN Y	A1	27 (15)	12
		A26	23 (17)	NP
39	LVYVNYARTE	A3	21	<5
40 & (41)	(Y)VNYARTED F	A26	(20)	NP
		B*08	15	<5
		B*2705	12	50
42	NYARTEDFF	A24	NP†	100
		Cw*0401	NP	120
43	YARTEDFF	B*08	16	<5
44	RTEDFFKLE	A1	21	<5
		A26	15	NP

†No prediction

5 **HLA-A*0201 binding assay:**

HLA-A*0201 binding studies were performed with PSMA₁₆₈₋₁₇₇, GMPEGDLVYV, (SEQ ID NO. 33) essentially as described in Example 3 above. As seen in figure 8, this epitope exhibits significant binding at even lower concentrations than the positive control peptides. The Melan-A peptide used as a control in this assay (and throughout this disclosure), ELAIGILTV, is actually a variant of the natural sequence (EAAGIGILTV) and exhibits a high affinity in this assay.

10

Example 5**Cluster Analysis (PSMA₂₈₁₋₃₁₀).**

Another peptide, RGIAEAVGLPSIPVHPIGYDDAQKLEKMG, PSMA₂₈₁₋₃₁₀, (SEQ ID NO. 45), containing an A1 epitope cluster from prostate specific membrane antigen, PSMA₂₈₃₋₃₀₇ (SEQ ID NO. 46), was synthesized using standard solid-phase F-moc chemistry on a 433A ABI Peptide synthesizer. After side chain deprotection and cleavage from the resin, peptide in ddH₂O was run on a reverse-phase preparative HPLC C18 column at following conditions: linear AB gradient (5% B/min) at a flow rate of 4 ml/min, where eluent A is 0.1% aqueous TFA and eluent B is 0.1% TFA in acetonitrile. A fraction at time 17.061 min containing the expected peptide as judged by mass spectrometry, was pooled and lyophilized. The peptide was then subjected to proteasome digestion and mass spectrum analysis essentially as described above. Prominent peaks from the mass spectra are summarized in Table 9.

Table 9. PSMA₂₈₁₋₃₁₀ Mass Peak Identification.

PEPTIDE	SEQUENCE	CALCULATE D MASS (MH ⁺)
281-297	RGIAEAVGLPSIPVHPI*	1727.07
286-297	AVGLPSIPVHPI**	1200.46
287-297	VGLPSIPVHPI	1129.38
288-297	GLPSIPVHPI†	1030.25
298-310	GYDDAQKLEKMG‡	1516.5
298-305	GYDDAQKLS	958.05
281-305	RGIAEAVGLPSIPVHPIGYDDAQKL	2666.12
281-307	RGIAEAVGLPSIPVHPIGYDDAQKLE	2908.39
286-307	AVGLPSIPVHPIGYDDAQKLE¶	2381.78
287-307	VGLPSIPVHPIGYDDAQKLE	2310.70
288-307	GLPSIPVHPIGYDDAQKLE#	2211.57
281-299	RGIAEAVGLPSIPVHPIGY	1947
286-299	AVGLPSIPVHPIGY	1420.69
287-299	VGLPSIPVHPIGY	1349.61
288-299	GLPSIPVHPIGY	1250.48
287-310	VGLPSIPVHPIGYDDAQKLEKMG	2627.14
288-310	GLPSIPVHPIGYDDAQKLEKMG	2528.01

Boldface sequences correspond to peptides predicted to bind to MHC, see Table 10.

*By mass alone this peak could also have been 296-310 or 288-303.

**By mass alone this peak could also have been 298-307. Combination of HPLC and mass spectrometry show that at some later time points this peak is a mixture of both species.

† By mass alone this peak could also have been 289-298.

By mass alone this peak could also have been 281-295 or 294-306.

§ By mass alone this peak could also have been 297-303.

¶ By mass alone this peak could also have been 285-306.

By mass alone this peak could also have been 288-303.

5 None of these alternate assignments are supported N-terminal pool sequence analysis.

N-terminal Pool Sequence Analysis

One aliquot at one hour of the proteasomal digestion (see Example 3 part 3 above) was subjected to N-terminal amino acid sequence analysis by an ABI 473A Protein Sequencer (Applied Biosystems, Foster City, CA). Determination of the sites and efficiencies of cleavage was based on consideration of the sequence cycle, the repetitive yield of the protein sequencer, and the relative yields of amino acids unique in the analyzed sequence. That is if the unique (in the analyzed sequence) residue X appears only in the nth cycle a cleavage site exists n-1 residues before it in the N-terminal direction. In addition to helping resolve any ambiguity in the assignment of mass to sequences, these data also provide a more reliable indication of the relative yield of the various fragments than does mass spectrometry.

For PSMA₂₈₁₋₃₁₀ (SEQ ID NO. 45) this pool sequencing supports two major cleavage sites after V₂₈₇ and I₂₉₇ among other minor cleavage sites. Reviewing the results presented in Fig. 9 reveals the following:

20 S at the 4th and 11th cycles indicating cleavage after V₂₈₇ and presence of the N-terminus of the substrate, respectively.

H at the 8th cycle indicating cleavage after V₂₈₇. The lack of decay in peak height at positions 9 and 10 versus the drop in height present going from 10 to 11 can suggest cleavage after A₂₈₆ and E₂₈₅ as well, rather than the peaks representing latency in the sequencing reaction.

25 D at the 2nd, 4th, and 7th cycles indicating cleavages after Y₂₉₉, I₂₉₇, and V₂₉₄, respectively.

This last cleavage is not observed in any of the fragments in Table 10 or in the alternate assignments in the notes below.

Q at the 6th cycle indicating cleavage after I₂₉₇.

30 M at the 10th and 12th cycle indicating cleavages after Y₂₉₉ and I₂₉₇, respectively.

Epitope Identification

Fragments co-C-terminal with 8-10 amino acid long sequences predicted to bind HLA by the SYFPEITHI or NIH algorithms were chosen for further study. The digestion and prediction steps of the procedure can be usefully practiced in any order. Although the substrate peptide used in proteasomal digest described here was specifically designed to include a predicted HLA-A1 binding sequence, the actual products of digestion can be checked after the fact for actual or predicted binding to other MHC molecules. Selected results are shown in Table 10.

Table 10.

Predicted HLA binding by proteasomally generated fragments: PSMA₂₈₁₋₃₁₀

SEQ ID NO.	PEPTIDE	HLA	SYFPEITHI	NIH
47 & (48)	(G)LPSIPVH PI	A*0201	16 (24)	(24)
		B*0702/B7	23	12
		B*5101	24	572
		Cw*0401	NP†	20
49 & (50)	(P)IGYYDAQ KL	A*0201	(16)	<5
		A26	(20)	NP
		B*2705	16	25
		B*2709	15	NP
		B*5101	21	57
		Cw*0301	NP	24
51 & (52)	(P)SIPVHPI GY	A1	21 (27)	<5
		A26	22	NP
		A3	16	<5
		B*5101	16	NP
53	IPVHPIGY			
54	YYDAQKLE	A1	22	<5

†No prediction

5

As seen in Table 10, N-terminal addition of authentic sequence to epitopes can often generate still useful, even better epitopes, for the same or different MHC restriction elements. Note for example the pairing of (G)LPSIPVHPI with HLA-A*0201, where the 10-mer can be used as a vaccine useful with several MHC types by relying on N-terminal trimming to create the epitopes for HLA-B7, -B*5101, and Cw*0401.

10

HLA-A*0201 binding assay:

HLA-A*0201 binding studies were performed with PSMA₂₈₈₋₂₉₇, GLPSIPVHPI, (SEQ ID NO. 48) essentially as described in Examples 3 and 4 above. As seen in figure 8, this epitope exhibits significant binding at even lower concentrations than the positive control peptides.

Example 6**Cluster Analysis (PSMA₄₅₄₋₄₈₁).**

Another peptide, SSIEGNYTLRV DCTPLMYSLVHLTKEL, PSMA₄₅₄₋₄₈₁, (SEQ ID NO. 55) containing an epitope cluster from prostate specific membrane antigen, was synthesized by MPS (purity >95%) and subjected to proteasome digestion and mass spectrum analysis as described above. Prominent peaks from the mass spectra are summarized in Table 11.

Table 11. PSMA₄₅₄₋₄₈₁ Mass Peak Identification.

MS PEAK (measured)	PEPTIDE	SEQUENCE	CALCULATED MASS (MH ⁺)
1238.5	454-464	SSIEGNYTLRV	1239.78
1768.38±0.60	454-469	SSIEGNYTLRV DCTPL	1768.99
1899.8	454-470	SSIEGNYTLRV DCTPLM	1900.19
1097.63±0.91	463-471	RVDCTPLMY	1098.32
2062.87±0.68	454-471*	SSIEGNYTLRV DCTPLMY	2063.36
1153	472-481**	SLVHNLTKE L	1154.36
1449.93±1.79	470-481	MYSLVHNLTKE L	1448.73

Boldface sequence correspond to peptides predicted to bind to MHC, see Table 12.

* On the basis of mass alone this peak could equally well be assigned to the peptide 455-472 however proteasomal removal of just the N-terminal amino acid is considered unlikely. If the issue were important it could be resolved by N-terminal sequencing.

**On the basis of mass this fragment might also represent 455-464.

Epitope Identification

Fragments co-C-terminal with 8-10 amino acid long sequences predicted to bind HLA by the SYFPEITHI or NIH algorithms were chosen for further study. The digestion and prediction steps of the procedure can be usefully practiced in any order. Although the substrate peptide used in proteasomal digest described here was specifically designed to include predicted HLA-A2.1 binding sequences, the actual products of digestion can be checked after the fact for actual or predicted binding to other MHC molecules. Selected results are shown in Table 12.

Table 12. Predicted HLA binding by proteasomally generated fragments

SEQ ID NO	PEPTIDE	HLA	SYFPEITHI	NIH
56 & (57)	(S) IEGNYTLRV	A1	(19)	<5
58	EGNYTLRV	A*0201	16 (22)	<5
		B*5101	15	NP†
59 & (60)	(Y) TLRVDCTPL	A*0201	20 (18)	(5)
		A26	16 (18)	NP
		B7	14	40
		B8	23	<5
		B*2705	12	30
		Cw*0301	NP	(30)
61	LRVDCTPLM	B*2705	20	600
		B*2709	20	NP
62 & (63)	(L) RVDCTPLMY	A1	32 (22)	125 (13.5)
		A3	25	<5
		A26	22	NP
		B*2702	NP	(200)
		B*2705	13 (NP)	(1000)

†No prediction

- 5 As seen in Table 12, N-terminal addition of authentic sequence to epitopes can often generate still useful, even better epitopes, for the same or different MHC restriction elements. Note for example the pairing of (L)RVDCTPLMY (SEQ ID NOS 62 and (63)) with HLA-B*2702/5, where the 10-mer has substantial predicted halftimes of dissociation and the co-C-terminal 9-mer does not. Also note the case of SIEGNYTLRV (SEQ ID NO 57) a predicted HLA-A*0201 epitope which can be used as a vaccine useful with HLA-B*5101 by relying on N-terminal trimming to create the epitope.

HLA-A*0201 binding assay

- 15 HLA-A*0201 binding studies were preformed, essentially as described in Example 3 above, with PSMA₄₆₀₋₄₆₉, TLRVDCTPL, (SEQ ID NO. 60). As seen in figure 10, this epitope was found to bind HLA-A2.1 to a similar extent as the known A2.1 binder FLPSDYFPSV (HBV₁₈₋₂₇; SEQ ID NO: 24) used as a positive control. Additionally, PSMA₄₆₁₋₄₆₉, (SEQ ID NO. 59) binds nearly as well.

ELISPOT analysis: PSMA₄₆₃₋₄₇₁ (SEQ ID NO. 62)

- 20 The wells of a nitrocellulose-backed microtiter plate were coated with capture antibody by incubating overnight at 4°C using 50 µl/well of 4µg/ml murine anti-human γ-IFN monoclonal

antibody in coating buffer (35 mM sodium bicarbonate, 15 mM sodium carbonate, pH 9.5). Unbound antibody was removed by washing 4 times 5 min. with PBS. Unbound sites on the membrane then were blocked by adding 200µl/well of RPMI medium with 10% serum and incubating 1 hr. at room temperature. Antigen stimulated CD8⁺ T cells, in 1:3 serial dilutions, were seeded into the wells of the microtiter plate using 100µl/well, starting at 2x10⁵ cells/well. (Prior antigen stimulation was essentially as described in Scheibenbogen, C. et al. *Int. J. Cancer* 71:932-936, 1997. PSMA₄₆₂₋₄₇₁ (SEQ ID NO. 62) was added to a final concentration of 10µg/ml and IL-2 to 100 U/ml and the cells cultured at 37°C in a 5% CO₂, water-saturated atmosphere for 40 hrs. Following this incubation the plates were washed with 6 times 200 µl/well of PBS containing 0.05% Tween-20 (PBS-Tween). Detection antibody, 50µl/well of 2g/ml biotinylated murine anti-human γ-IFN monoclonal antibody in PBS+10% fetal calf serum, was added and the plate incubated at room temperature for 2 hrs. Unbound detection antibody was removed by washing with 4 times 200 µl of PBS-Tween. 100µl of avidin-conjugated horseradish peroxidase (Pharmingen, San Diego, CA) was added to each well and incubated at room temperature for 1 hr. Unbound enzyme was removed by washing with 6 times 200 µl of PBS-Tween. Substrate was prepared by dissolving a 20 mg tablet of 3-amino 9-ethylcarbazole in 2.5 ml of N, N-dimethylformamide and adding that solution to 47.5 ml of 0.05 M phosphate-citrate buffer (pH 5.0). 25 µl of 30% H₂O₂ was added to the substrate solution immediately before distributing substrate at 100 µl/well and incubating the plate at room temperature. After color development (generally 15-30 min.), the reaction was stopped by washing the plate with water. The plate was air dried and the spots counted using a stereomicroscope.

Figure 11 shows the detection of PSMA₄₆₃₋₄₇₁ (SEQ ID NO. 62)-reactive HLA-A1⁺ CD8⁺ T cells previously generated in cultures of HLA-A1⁺ CD8⁺ T cells with autologous dendritic cells plus the peptide. No reactivity is detected from cultures without peptide (data not shown). In this case it can be seen that the peptide reactive T cells are present in the culture at a frequency between 1 in 2.2x10⁴ and 1 in 6.7x10⁴. That this is truly an HLA-A1-restricted response is demonstrated by the ability of anti-HLA-A1 monoclonal antibody to block γ-IFN production; see figure 12.

Example 7

Cluster Analysis (PSMA₆₅₃₋₆₈₇).

Another peptide, FDKSNPIVLRMMNDQLMFLERAFIDPLGLPDRPFY PSMA₆₅₃₋₆₈₇, (SEQ ID NO. 64) containing an A2 epitope cluster from prostate specific membrane antigen, PSMA₆₆₀₋₆₈₁ (SEQ ID NO 65), was synthesized by MPS (purity >95%) and subjected to proteasome digestion and mass spectrum analysis as described above. Prominent peaks from the mass spectra are summarized in Table 13.

Table 13. PSMA₆₅₃₋₆₈₇ Mass Peak Identification.

MS PEAK (measured)	PEPTIDE	SEQUENCE	CALCULATED MASS (MH ⁺)
906.17±0.65	681-687**	LPDRPFY	908.05
1287.73±0.76	677-687**	DPLGLPDRPFY	1290.47
1400.3±1.79	676-687	IDPLGLPDRPFY	1403.63
1548.0±1.37	675-687	FIDPLGLPDRPFY	1550.80
1619.5±1.51	674-687**	AFIDPLGLPDRPFY	1621.88
1775.48±1.32	673-687*	RAFIDPLGLPDRPFY	1778.07
2440.2±1.3	653-672	FDKSNPIVLRMMNDQLMFLE	2442.932
1904.63±1.56	672-687*	ERAFIDPLGLPDRPFY	1907.19
2310.6±2.5	653-671	FDKSNPIVLRMMNDQLMFL	2313.82
2017.4±1.94	671-687	LERAFIDPLGLPDRPFY	2020.35
2197.43±1.78	653-670	FDKSNPIVLRMMNDQLMF	2200.66

Boldface sequence correspond to peptides predicted to bind to MHC, see Table 13.

* On the basis of mass alone this peak could equally well be assigned to a peptide beginning at 654, however proteasomal removal of just the N-terminal amino acid is considered unlikely. If the issue were important it could be resolved by N-terminal sequencing.

** On the basis of mass alone these peaks could have been assigned to internal fragments, but given the overall pattern of digestion it was considered unlikely.

Epitope Identification

Fragments co-C-terminal with 8-10 amino acid long sequences predicted to bind HLA by the SYFPEITHI or NIH algorithms were chosen for further study. The digestion and prediction steps of the procedure can be usefully practiced in any order. Although the substrate peptide used in proteasomal digest described here was specifically designed to include predicted HLA-A2.1 binding sequences, the actual products of digestion can be checked after the fact for actual or predicted binding to other MHC molecules. Selected results are shown in Table 14.

Table 14. Predicted HLA binding by proteasomally generated fragments

SEQ ID NO	PEPTIDE	HLA	SYFPEITHI	NIH
66 & (67)	(R)MMNDQLMF L	A*0201	24 (23)	1360 (722)
		A*0205	NP†	71 (42)
		A26	15	NP
		B*2705	12	50
68	RMMNDQLMF	B*2705	17	75

†No prediction

5

As seen in Table 14, N-terminal addition of authentic sequence to epitopes can generate still useful, even better epitopes, for the same or different MHC restriction elements. Note for example the pairing of (R)MMNDQLMFL (SEQ ID NOS. 66 and (67)) with HLA-A*02, where the 10-mer retains substantial predicted binding potential.

10

HLA-A*0201 binding assay

HLA-A*0201 binding studies were performed, essentially as described in Example 3 above, with PSMA₆₆₃₋₆₇₁, (SEQ ID NO. 66) and PSMA₆₆₂₋₆₇₁, RMMNDQLMFL (SEQ NO. 67). As seen in figures 10, 13 and 14, this epitope exhibits significant binding at even lower concentrations than the positive control peptide (FLPSDYFPSV (HBV₁₈₋₂₇); SEQ ID NO: 24). Though not run in parallel, comparison to the controls suggests that PSMA₆₆₂₋₆₇₁ (which approaches the Melan A peptide in affinity) has the superior binding activity of these two PSMA peptides.

15

Example 8**Vaccinating with epitope vaccines.**

20

1. Vaccination with peptide vaccines:A. Intranodal delivery

A formulation containing peptide in aqueous buffer with an antimicrobial agent, an antioxidant, and an immunomodulating cytokine, was injected continuously over several days into the inguinal lymph node using a miniature pumping system developed for insulin delivery (MiniMed; Northridge, CA). This infusion cycle was selected in order to mimic the kinetics of antigen presentation during a natural infection.

25

B. Controlled release

A peptide formulation is delivered using controlled PLGA microspheres as is known in the art, which alter the pharmacokinetics of the peptide and improve immunogenicity. This formulation is injected or taken orally.

30

C. Gene gun delivery

A peptide formulation is prepared wherein the peptide is adhered to gold microparticles as is known in the art. The particles are delivered in a gene gun, being accelerated at high speed so as to penetrate the skin, carrying the particles into dermal tissues that contain pAPCs.

5 D. Aerosol delivery

A peptide formulation is inhaled as an aerosol as is known in the art, for uptake into appropriate vascular or lymphatic tissue in the lungs.

2. Vaccination with nucleic acid vaccines:

10 A nucleic acid vaccine is injected into a lymph node using a miniature pumping system, such as the MiniMed insulin pump. A nucleic acid construct formulated in an aqueous buffered solution containing an antimicrobial agent, an antioxidant, and an immunomodulating cytokine, is delivered over a several day infusion cycle in order to mimic the kinetics of antigen presentation during a natural infection.

Optionally, the nucleic acid construct is delivered using controlled release substances, such as PLGA microspheres or other biodegradable substances. These substances are injected or taken orally. Nucleic acid vaccines are given using oral delivery, priming the immune response through uptake into GALT tissues. Alternatively, the nucleic acid vaccines are delivered using a gene gun, wherein the nucleic acid vaccine is adhered to minute gold particles. Nucleic acid constructs can also be inhaled as an aerosol, for uptake into appropriate vascular or lymphatic tissue in the lungs.

20 **Example 9**

Assays for the effectiveness of epitope vaccines.

1. Tetramer analysis:

Class I tetramer analysis is used to determine T cell frequency in an animal before and after administration of a housekeeping epitope. Clonal expansion of T cells in response to an epitope indicates that the epitope is presented to T cells by pAPCs. The specific T cell frequency is measured against the housekeeping epitope before and after administration of the epitope to an animal, to determine if the epitope is present on pAPCs. An increase in frequency of T cells specific to the epitope after administration indicates that the epitope was presented on pAPC.

2. Proliferation assay:

30 Approximately 24 hours after vaccination of an animal with housekeeping epitope, pAPCs are harvested from PBMCs, splenocytes, or lymph node cells, using monoclonal antibodies against specific markers present on pAPCs, fixed to magnetic beads for affinity purification. Crude blood or splenocyte preparation is enriched for pAPCs using this technique. The enriched pAPCs are then used in a proliferation assay against a T cell clone that has been generated and is specific for the housekeeping epitope of interest. The pAPCs are coincubated with the T cell clone and the T cells are monitored for proliferation activity by measuring the incorporation of radiolabeled

thymidine by T cells. Proliferation indicates that T cells specific for the housekeeping epitope are being stimulated by that epitope on the pAPCs.

3. Chromium release assay:

A human patient, or non-human animal genetically engineered to express human class I MHC, is immunized using a housekeeping epitope. T cells from the immunized subject are used in a standard chromium release assay using human tumor targets or targets engineered to express the same class I MHC. T cell killing of the targets indicates that stimulation of T cells in a patient would be effective at killing a tumor expressing a similar TuAA.

Example 10

Induction of CTL response with naked DNA is efficient by Intra-lymph node immunization.

In order to quantitatively compare the CD8⁺ CTL responses induced by different routes of immunization a plasmid DNA vaccine (pEGFPL33A) containing a well-characterized immunodominant CTL epitope from the LCMV-glycoprotein (G) (gp33; amino acids 33-41) (Oehen, S., et al., *Immunology* 99, 163-169 2000) was used, as this system allows a comprehensive assessment of antiviral CTL responses. Groups of 2 C57BL/6 mice were immunized once with titrated doses (200-0.02μg) of pEGFPL33A DNA or of control plasmid pEGFP-N3, administered i.m. (intramuscular), i.d. (intradermal), i.spl. (intrasplenic), or i.ln. (intra-lymph node). Positive control mice received 500 pfu LCMV i.v. (intravenous). Ten days after immunization spleen cells were isolated and gp33-specific CTL activity was determined after secondary *in vitro* restimulation. As shown in Fig. 15, i.m. or i.d. immunization induced weakly detectable CTL responses when high doses of pEGFPL33A DNA (200μg) were administered. In contrast, potent gp33-specific CTL responses were elicited by immunization with only 2μg pEGFPL33A DNA i.spl. and with as little as 0.2μg pEGFPL33A DNA given i.ln. (figure 15; symbols represent individual mice and one of three similar experiments is shown). Immunization with the control pEGFP-N3 DNA did not elicit any detectable gp33-specific CTL responses (data not shown).

Example 11

Intra-lymph node DNA immunization elicits anti-tumor immunity.

To examine whether the potent CTL responses elicited following i.ln. immunization were able to confer protection against peripheral tumors, groups of 6 C57BL/6mice were immunized three times at 6-day intervals with 10μg of pEGFPL33A DNA or control pEGFP-N3 DNA. Five days after the last immunization small pieces of solid tumors expressing the gp33 epitope (EL4-33) were transplanted s.c. into both flanks and tumor growth was measured every 3-4d. Although the EL4-33 tumors grew well in mice that had been repetitively immunized with control pEGFP-N3 DNA (figure 16), mice which were immunized with pEGFPL33A DNA i.ln. rapidly eradicated the peripheral EL4-33 tumors (figure 16).

Example 12

Differences in lymph node DNA content mirrors differences in CTL response following intra-lymph node and intramuscular injection.

pEFGPL33A DNA was injected i.ln. or i.m. and plasmid content of the injected or draining lymph node was assessed by real time PCR after 6, 12, 24, 48 hours, and 4 and 30 days. At 6, 12, and 24 hours the plasmid DNA content of the injected lymph nodes was approximately three orders of magnitude greater than that of the draining lymph nodes following i.m. injection. No plasmid DNA was detectable in the draining lymph node at subsequent time points (Fig. 17). This is consonant with the three orders of magnitude greater dose needed using i.m. as compared to i.ln. injections to achieve a similar levels of CTL activity. CD8⁺ knockout mice, which do not develop a CTL response to this epitope, were also injected i.ln. showing clearance of DNA from the lymph node is not due to CD8⁺ CTL killing of cells in the lymph node. This observation also supports the conclusion that i.ln. administration will not provoke immunopathological damage to the lymph node.

Example 13

Administration of a DNA plasmid formulation of a therapeutic vaccine for melanoma to humans.

SYNCHROTOPE TA2M, a melanoma vaccine, encoding the HLA-A2-restricted tyrosinase epitope SEQ ID NO. 1 and epitope cluster SEQ ID NO. 69, was formulated in 1% Benzyl alcohol, 1% ethyl alcohol, 0.5mM EDTA, citrate-phosphate, pH 7.6. Aliquots of 80, 160, and 320 µg DNA/ml were prepared for loading into MINIMED 407C infusion pumps. The catheter of a SILHOUETTE infusion set was placed into an inguinal lymph node visualized by ultrasound imaging. The assembly of pump and infusion set was originally designed for the delivery of insulin to diabetics and the usual 17mm catheter was substituted with a 31mm catheter for this application. The infusion set was kept patent for 4 days (approximately 96 hours) with an infusion rate of about 25 µl/hour resulting in a total infused volume of approximately 2.4 ml. Thus the total administered dose per infusion was approximately 200, and 400 µg; and can be 800 µg, respectively, for the three concentrations described above. Following an infusion subjects were given a 10 day rest period before starting a subsequent infusion. Given the continued residency of plasmid DNA in the lymph node after administration (as in example 12) and the usual kinetics of CTL response following disappearance of antigen, this schedule will be sufficient to maintain the immunologic CTL response.

Example 14**Additional Epitopes.**

The methodologies described above, and in particular in examples 3-7, have been applied to additional synthetic peptide substrates, leading to the identification of further epitopes as set for
5 the in tables 15-36 below. The substrates used here were designed to identify products of housekeeping proteasomal processing that give rise to HLA-A*0201 binding epitopes, but additional MHC-binding reactivities can be predicted, as discussed above. Many such reactivities are disclosed, however, these listings are meant to be exemplary, not exhaustive or limiting. As also discussed above, individual components of the analyses can be used in varying combinations and orders. The digests of the NY-ESO-1 substrates 136-163 and 150-177 (SEQ ID NOS. 254 and
10 255, respectively) yielded fragments that did not fly well in MALDI-TOF mass spectrometry. However, they were quite amenable to N-terminal peptide pool sequencing, thereby allowing identification of cleavage sites. Not all of the substrates necessarily meet the formal definition of an epitope cluster as referenced in example 3. Some clusters are so large, e.g. NY-ESO-1₈₆₋₁₇₁, that
15 it was more convenient to use substrates spanning only a portion of this cluster. In other cases, substrates were extended beyond clusters meeting the formal definition to include neighboring predicted epitopes. In some instances, actual binding activity may have dictated what substrate was made, as with for example the MAGE epitopes reported here, where HLA binding activity was determined for a selection of peptides with predicted affinity, before synthetic substrates were
20 designed.

Table 15
GP100: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

†Scores are given from the two binding prediction programs referenced above (see example 3).

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI /NIH)†					Comments
				A*0201	A1	A3	B7	B8	
609-644	630-638*	LPHSSSHWL	88				20/80	16/<5	*The digestion of 609-644 and 622-650 have generated the same epitopes.
	629-638*	QLPHSSSHWL	89	21/117					
	614-622	LYRRRLMK	90			32/20			
	613-622	SLYRRRLMK	91	14/<5		29/60			
	615-622	IYRRRLMK	92					15/<5	
622-650	630-638*	LPHSSSHWL	93				20/80	16/<5	
	629-638*	QLPHSSSHWL	94	21/117					

Table 16A
MAGE-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI /NIH)†						
				A*0201	A1	A3	B7	B8	Other	
86-109	95-102	ESLFRVI	95					16/<5		
	93-102	ILESFRVI	96	21/<5		20/<5				
	93-101	ILESFRV	97	23/<5						
	92-101	CILESFRV	98	23/55						
	92-100	CILESFR	99	20/138						
263-292	263-271	EFLWGPRAL	100						A26 (R 21), A24 (NIH 30)	
	264-271	FLWGPRAL	101					17/<5		
	264-273	FLWGPRALAE	102	16/<5		19/<5				
	265-274	LWGPRALAEI	103	16/<5						
	268-276	PRALAEISY	104	15/<5						
	267-276	GPRALAEISY	105	15/<5			<15/<5		B4403 (NIH 7); B3501 (NIH 120)	
	269-277	RALAEISYV	106	18/20						
	271-279	LAETSYVKV	107	19/<5						
	270-279	ALAETSYVKV	108	30/427		19/<5<5				
	272-280	AETSYVKVL	109	15/<5					B4403 (NIH 36)	
	271-280	LAETSYVKVL	110	18/<5			<15/<5			
	274-282	TSYVKVLEY	111		26/<5				B4403 (NIH 14)	
	273-282	ETSYVKVLEY	112		28/6				A26 (R 31), B4403 (NIH 14)	
278-286	KVLEYVKV		113	26/743		16/<5				

Table 16B
MAGE-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI /NIH) [†]					
				A*0201	A1	A3	B7	B8	Other
168-193	168-177	SYVLVTCLGL	114						A24 (NIH 300)
	169-177	YVLVTCLGL	115	20/32		15/<5	<15/20		
	170-177	VLVTCLGL	116					17/<5	
229-258	240-248	TQDLVQEKY	117		29/<5				
	239-248	LTQDLVQEKY	118		23/<5				A26 (R 22)
	232-240	YGPRKLLT	119		24/11				
	243-251	LVQEKYLEY	120		21/<5	21/<5			A26 (R 28)
	242-251	DLVQEKYLEY	121		22/<5	19/<5			A26 (R 30)
	230-238	SAYGEPRKL	122	21/<5					B5101 (25/121)
	278-286	KVLEYVVKV	123	26/743		16/<5			
272-297	277-286	VKVLEYVVKV	124	17/<5					
	276-284	YVKVLEYVI	125	15/<5		15/<5		17/<5	
	274-282	TSYVKVLEY	126		26/<5				
	273-282	ETSYVKVLEY	127		28/6				
	283-291	VKVSARVR	128			20/<5			
	282-291	YVIKVSARVR	129			24/<5			

[†]Scores are given from the two binding prediction programs referenced above (see example 3). R indicates a SYFPEITHI score.

Table 17A

MAGE-2: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID		HLA Binding Predictions (SYFPEITHI /NIH) [†]							
			NO	A*0201	A1	A3	B7	B8	Other			
107-126	115-122	ELVHFLLL	130					18/<5				
	113-122	MVELVHFLLL	131		21/<5				A26 (R 22)			
	109-116	ISRKMVEL	132					17/<5				
	108-116	AI SRKMVEL	133	25/7		19/<5	16/12	26/<5				
	107-116	A AI SRKMVEL	134	22/<5			14/36	n.p./16				
	112-120	KMVELVHFL	135	27/2800								
	109-117	ISRKMVELV	136	16/<5								
	108-117	AI SRKMVELV	137	24/11								
	116-124	LVHFLLLKY	138		23/<5	19/<5			A26 (R 26)			
	115-124	ELVHFLLLKY	139		24/<5	19/5			A26 (R 29)			
	111-119	RKMVELVHF	140									
	158-166	LQLVFGIEV	141	17/168								
	157-166	YLQLVFGIEV	142	24/1215								
	159-167	QLVFGIEVV	143	25/32		18/<5						
145-175	158-167	LQLVFGIEVV	144	18/20								
	164-172	IEVVEVVPI	145	16/<5								
	163-172	GIEVVEVVPI	146	22/<5								
	162-170	FGIEVVEVV	147	19/<5					B5101(24/69.212)			
	154-162	ASEYLQLVF	148		22/68							
	153-162	KASEYLQLVF	149			15/<5						

[†]Scores are given from the two binding prediction programs referenced above (see example 3). R indicates a SYFPEITHI score.

Table 17B
MAGE-2: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence		HLA Binding Predictions (SYFPEITHI/NIH) [†]					
				A*0201	A1	A3	B7	B8	Other
213-233	218-225	EEKIWEEL	150					22/<5	
	216-225	APEEKIWEEL	151	15/<5			22/72		
	216-223	APEEKIWE	152					18/<5	
	220-228	KIWEEL SML	153	26/804		16/<5		16/<5	A26 (R 26)
	219-228	EKIWEEL SML	154						A26 (R 22)
271-291	271-278	FLWGPRAL	155					17/<5	
	271-279	FLWGPRALJ	156	25/398		16/7			
	278-286	LIETSYVKV	157	23/<5					
	277-286	ALIETSYVKV	158	30/427		21/<5			
	276-284	RALIETSYV	159	18/19					B5101 (20/55)
	279-287	IETSYVKVL	160	15/<5					
	278-287	LIETSYVKVL	161	22/<5					A26 (R 22)

[†]Scores are given from the two binding prediction programs referenced above (see example 3). R indicates a SYFPEITHI score.

Table 18
MAGE-3: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI/NIH) [†]					
				A*0201	A1	A3	B7	B8	Other
267-286	271-278	FLWGPRAL	162					17/<5	
	270-278	EFLWGPRAL	163						A26 (R 21); A24 (NIH 30)
	271-279	FLWGPRALV	164	27/2655		16/<5			
	276-284	RALVETSYV	165	18/19					B5101 (20/55)
	272-280	LWGPRALVE	166			15/<5			
	271-280	FLWGPRALVE	167	15/<5		22/<5			
	272-281	LWGPRALVET	168	16/<5					

[†]Scores are given from the two binding prediction programs referenced above (see example 3). R indicates a SYFPEITHI score.

Table 19A
NY-ESO-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI/NIH) [†]						
				A*0201	A1	A3	B7	B8	Other	
81-113	82-90	GPESRLLEF	169		16/11		18/<5	22/<5		
	83-91	PESRLLEFY	170		15/<5				B4403 (NIH 18)	
	82-91	GPESRLLEFY	171		25/11					
	84-92	ESRLLEFY	172					19/8		
	86-94	RLLEFYLAM	173	21/430		21/<5				
101-133	88-96	LEFYLAMPF	174						B4403 (NIH 60)	
	87-96	LLEFYLAMPE	175		<15/45	18/<5				
	93-102	AMPFATPMEA	176	15/<5						
	94-102	MPFATPMEA	177				17/<5			
	115-123	PLPVPGVLL	178	20/<5		17/<5	16/<5	18/<5		
	114-123	PPLPVPGVLL	179				23/12			
	116-123*	LPVPGVLL	180					16/<5		
	103-112	ELARRSLAQD	181	15/<5		20/<5			*Evidence of the same epitope obtained from two digests.	
	118-126*	VPGVLLKEF	182				17/<5	16/<5		
	117-126*	PVPGVLLKEF	183			16/<5				
116-123*	LPVPGVLL	184					16/<5			
116-145	127-135	TVSGNILTI	185	21/<5		19/<5				
	126-135	FTVSGNILTI	186	20/<5						
	120-128	GVLKFTV	187	20/130		18/<5				
	121-130	VLLKEFTVSG	188	17/<5		18/<5				
	122-130	LLKEFTVSG	189	20/<5		18/<5				
	118-126*	VPGVLLKEF	190				17/<5	16/<5		
	117-126*	PVPGVLLKEF	191			16/<5				

[†]Scores are given from the two binding prediction programs referenced above (see example 3).

Table 19B
NY-ESO-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI/NIH) [†]					
				A*0201	A1	A3	B7	B8	Other
136-163 (SEQ ID NO 254)	139-147	AADHRQLQL	192	17/<5	17/<5			22/<5	
	148-156	SISSCLQQL	193	24/7					A26 (R 25)
	147-156	LSISSCLQQL	194	18/<5					
	138-147	TAADHRQLQL	195	18/<5					
	161-169	WTQCFLPV	196	18/84					
150-177 (SEQ ID NO 255)	157-165	SLLMWITQC	197	18/42		17/<5			
	150-158	SSCLQQLSL	198	15/<5					
	154-162	QQLSLLMWI	199	15/50					
	151-159	SCLQQLSLL	200	18/<5					
	150-159	SSCLQQLSLL	201	16/<5					
	163-171	TQCFLPVFL	202	<15/12					
	162-171	ITQCFLPVFL	203	18/<5					A26 (R 19)

[†]Scores are given from the two binding prediction programs referenced above (see example 3). R indicates a SYFPEITHI score

Table 20
PRAME: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI/NIH) [†]					
				A*0201	A1	A3	B7	B8	Other
211-245	219-227	PMQDIKMIL	204	16/<5				16/n.d.	A26 (R 20)
	218-227	MPMQDIKMIL	205				<15/240		
411-446	428-436	QHLIGLSNL	206	18/<5					
	427-436	LQHLIGLSNL	207	16/8					
	429-436	HLIGLSNL	208						
	431-439	IGLSNLTHV	209	18/7				17/<5	B15 (R 21)
	430-439	LIGLSNLTHV	210	24/37					B*5101 (R 22)

[†]Scores are given from the two binding prediction programs referenced above (see example 3). R indicates a SYFPEITHI score.

Table 21
PSA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI/NIH) [†]						
				A*0201	A1	A3	B7	B8	Other	
42-77	53-61	VLVHPQWVL	211	22/112			<15/6	17/<5		
	52-61	GVLVHPQWVL	212	17/21		16/<5	<15/30		A26 (R 18)	
	52-60	GVLVHPQWV	213	17/124						
	59-67	WVLTAAHCL	214	15/16						
	54-63	LVHPQWVLTA	215	19/<5		20/<5			A26 (R 16)	
	53-62	VLVHPQWVLT	216	17/22						
	54-62	LVHPQWVLT	217			17/n.d.				
55-95	66-73	CRNKSVI	218					26/20		
	65-73	HCIRNKSVI	219					<15/16		
	56-64	HPQWVLTA	220				18/<5			
	63-72	AAHCIRNKSV	221	17/<5						

[†]Scores are given from the two binding prediction programs referenced above (see example 3). R indicates a SYFPEITHI score.

Table 22
PSCA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI /NIH) [†]					
				A*0201	A1	A3	B7	B8	Other
93-123*	116-123	LLWGPQQL	222					16/<5	
	115-123	LLWGPQQL	223	<15/18					
	114-123	GLLLWGPQQL	224	<15/10					
	99-107	ALQPAAAIL	225	26/9		22/<5	<15/12	16/<5	A26 (R 19)
	98-107	HALQPAAAIL	226	18/<5			<15/12		

*L123 is the C-terminus of the natural protein.

[†]Scores are given from the two binding prediction programs referenced above (see example 3).

Table 23
Tyrosinase: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI/NIB) [†]					
				A*0201	A1	A3	B7	B8	Other
128-157	128-137	APEKDKFFAY	227		29/6		15/<5		B4403 (NIH 14)
	129-137	PEKDKFFAY	228		18/<5			21/<5	
	130-138	EKDKFFAYL	229				15/<5		
	131-138	KDKFFAYL	230					20/<5	
197-228	205-213	PAFLPWHRL	231					15/<5	
	204-213	APAFLPWHRL	232				23/360		
	207-216	FLPWHRLFL	1	25/1310				<15/8	
	208-216	LPWHRLFL	9	17/26			20/80	24/16	
191-211	214-223	FLLRWEQEIQ	233			15/<5			
	212-220	RLFLLRWEQ	234			16/<5			
	191-200	GSEIWRDIDF	235		18/68				
	192-200	SEIWRDIDF	236					16/<5	B4403 (NIH 400)
207-230	207-215	FLWHRLFL	8	22/540			<15/6	17/<5	
466-484	473-481	RIWSWLLGA	237	19/13		15/<5			
476-497	476-484	SWLLGAAMV	238	18/<5					
	477-486	WLLGAAMVGA	239	21/194		18/<5			
	478-486	LIIGAAMVGA	240	19/19		16/<5			

[†]Scores are given from the two binding prediction programs referenced above (see example 3).

Table 24
PSMA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI /NIH) [†]					
				A*0201	A1	A3	B7	B8	Other
1-30	4-12	LLHEITDSAV	241	25/485		15/<5			
	13-21	ATARRPRWL	242	18/<5				18/<5	A26 (R 19)
53-80	53-61	TPKHNMKAF	243					24/<5	
	64-73	ELKAENIKKF	244			17/<5			A26 (R 30)
	69-77	NIKKFLH [†] NF	245						A26 (R 27)
	68-77	ENIKKFLH [†] NF	246						A26 (R 24)
215-244	220-228	AGAKGVILY	247		25/<5				
457-489	468-477	PLMYSLVHNL	248	22/<5					
	469-477	LMYSLVHNL	249	27/193		<15/9			
	463-471	RVDCITPLMY	250		32/125	25/<5			A26 (R 22)
	465-473	DCTPLMYSL	251						A26 (R 22)
503-533	507-515	SGMPRISKL	252	21/<5				21/<5	
	506-515	FSGMPRISKL	253	17/<5					

[†]This H was reported as Y in the SWISSPROT database.

[†]Scores are given from the two binding prediction programs referenced above (see example 3).

Table 25A
MAGE-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
Mage-1 119-146	125-132	KAEMLESV	256	B5101	19	n.a.
	124-132	TKAEMLESV	257	A0201	20	<5
	123-132	VTKAEMLESV	258	A0201	20	<5
				A1	28	45
	128-136	MLESVIKNY	259	A26	24	n.a.
				A3	17	5
	127-136	EMLESVIKNY	260	A1	15	<1.0
				A26	23	<1.0
	125-133	KAEMLESVI	261	B5101	23	100
				A24	N.A.	4
Mage-1 143-170	146-153	KASESLQL	262	B08	16	<1.0
				B5101	17	N.A.
	145-153	GKASESLQL	263	B2705	17	1
	147-155	ASESLQLVF	264	B2709	16	N.A.
	153-161	LVFGIDVKE	265	A1	22	68
				A26	16	N.A.
				A3	16	<1.0

Table 25B

MAGE-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
Mage-1 99-125	114-121	LLKYRARE	266	B8	25	<1.0
	106-113	VADLVGFL	267	B8	16	<1.0
				B5101	21	N.A.
				A0201	23	44
				A26	25	N.A.
	105-113	KVADLVGFL	268	A3	16	<5
				B0702	14	20
				B2705	14	30
				A0201	17	<5
	107-115	ADLVGFLLL	269	B0702	15	<5
				B2705	16	1
	106-115	VADLVGFLLL	270	A0201	16	<5
				A1	22	3
	114-123	LLKYRAREPV	271	A0201	20	2

Table 26
MAGE-3: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
Mage-3 267-295	271-278	FLWGPRAL	162	B08	17	<5
	270-278	EFLWGPRAL	163	A26	21	N.A.
				A24	N.A.	30
				B1510	16	N.A.
	271-279	FLWGPRALV	164	A0201	27	2655
				A3	16	2
	278-286	LVETSYVKV	272	A0201	19	<1.0
				A26	17	N.A.
				A0201	28	428
	277-286	ALVETSYVKV	273	A26	16	<5
				A3	18	<5
				A0201	19	27
	285-293	KVLHMMVKI	274	A3	19	<5
				A0201	18	20
	276-284	RALVETSYV	165	A0201	17	<1.0
	283-291	YVKVLHMMV	275	A0201	17	<1.0
	275-283	PRALVETSY	276	A1	17	<1.0
	274-283	GPRALVETSY	277	A1	15	<1.0
	278-287	LVETSYVKVL	278	A0201	18	<1.0
	272-281	LWGPRALVET	168	A0201	16	<1.0
	271-280	FLWGPRALVE	167	A3	22	<5

Table 27A
Fibronectin ED-B: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
ED-B 14'-21*	4'-5**	<i>TIIP</i> EV [†] PQL	279	A0201	27	7
				A26	28	N.A.
				A3	17	<5
				B8	15	<5
				B1510	15	N.A.
				B2705	17	10
				B2709	15	N.A.
	5'-5**	<i>DTIIP</i> EV [†] PQL	280	A0201	20	<5
				A26	32	N.A.
				A26	29	N.A.
	1-10	EVPQLTDLSF	281			

*This substrate contains the 14 amino acids from fibronectin flanking ED-B to the N-terminal side.

**These peptides span the junction between the N-terminus of the ED-B domain and the rest of fibronectin.

† The *italicized* lettering indicates sequence outside the ED-B domain.

Table 27B
Fibronectin ED-B: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
ED-B 8-35	23-30	TPLNSSTI	282	B5101	22	N.A.
	18-25	IGLRWTPL	283	B5101	18	N.A.
	17-25	SIGLRWTPL	284	A0201	20	5
				A26	18	N.A.
				B08	25	<5
	25-33	LNSSTIIGY	285	A1	19	<5
				A26	16	<5
	24-33	PLNSSTIIGY	286	A1	20	<5
				A26	24	N.A.
				A3	16	<5
	23-31	TPLNSSTII	287	B0702	17	8
				B5101	25	440

Table 27C
Fibronectin ED-B: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Blinding Prediction		
				HLA type	SYFPEITHI	NIH
ED-B 20-49	31-38	IGYRITVV	288	B5101	25	N.A.
	30-38	IIGYRITVV	289	A0201	23	15
				A3	17	<1.0
				B08	15	<1.0
				B5101	15	3
	29-38	TIIGYRITVV	290	A0201	26	9
				A26	18	N.A.
				A3	18	<5
	23-30	TPLNSSTI	282	B5101	22	N.A.
	25-33	LNSSTIIGY	285	A1	19	<5
				A26	16	N.A.
	24-33	PLNSSTIIGY	286	A26	24	N.A.
				A3	16	<5
	31-39	IIGYRITVVA	291	A3	17	<5
	30-39	IIGYRITVVA	292	A0201	15	<5
				A3	18	<5
23-31	TPLNSSTII		287	B0702	17	8
				B5101	25	440

Table 28A

CEA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
CEA 176-202	184-191	SLPVSPRL	293	B08	19	<5
	183-191	QSLPVSPRL	294	A0201	15	<5
				B1510	15	
				B2705	18	10
				B2709	15	
	186-193	PVSPRLQL	295	B08	18	<5
	185-193	LPVSPRLQL	296	B0702	26	180
				B08	16	<5
				B5101	19	130
	184-193	SLPVSPRLQL	297	A0201	23	21
				A26	18	N.A.
				A3	18	<5
				B5101	17	N.A.
	185-192	LPVSPRLQ	298	A0201	21	4
	192-200	QLSNGNRTL	299	A26	16	N.A.
				A3	19	<5
				B08	17	<5
				B1510	15	
	191-200	LQLSNGNRTL	300	A0201	16	3
	179-187	WVNNQSLPV	301	A0201	16	28
	186-194	PVSPRLQLS	302	A26	17	N.A.
				A3	15	<5

Table 28B
CEA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
CEA 354-380	362-369	SLPVSPRL	303	B08	19	<1.0
	361-369	QSLPVSPRL	304	A0201	15	<1.0
				B2705	18	10
				B2709	15	
	364-371	PVSPRLQL	305	B08	18	<1.0
	363-371	LPVSPRLQL	306	B0702	26	180
				B08	16	<1.0
				B5101	19	130
	362-371	SLPVSPRLQL	307	A0201	23	21
				A26	18	N.A.
				A24	N.A.	6
				A3	18	<5
				B5101	17	N.A.
	363-370	LPVSPRLQ	308	A0201	22	4
	370-378	QLSNDNRTL	309	A26	16	N.A.
				A3	17	<1.0
				B08	17	<1.0
	369-378	LQLSNDNRTL	310	A0201	16	3
	357-365	WVNNQSLPV	311	A0201	16	28
	360-368	NQSLPVSPR	312	B2705	14	100

Table 28C CEA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion						
Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
CEA 532-558	540-547	SLPVSPRL	313	B08	19	<5
	539-547	QSLPVSPRL	314	A0201	15	<5
				B1510	15	<5
				B2705	18	10
				B2709	15	
	542-549	PVSPRLQL	315	B08	18	<5
	541-549	LPVSPRLQL	316	B0702	26	180
				B08	16	<1.0
				B5101	19	130
	540-549	SLPVSPRLQL	317	A0201	23	21
				A26	18	N.A.
				A3	18	<5
	541-548	LPVSPRLQ	318	B5101	17	N.A.
	548-556	QLSNGNRTL	319	A0201	24	4
				A26	16	N.A.
				A3	19	<1.0
				B08	17	<1.0
	547-556	LQLSNGNRTL	320	B1510	15	
	535-543	WVNGQSLPV	321	A0201	16	3
				A0201	18	28
	533-541	LWVVGQSL	322	A3	15	<1.0
				A0201	15	<5

Table 28D
CEA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
CEA 532-558 (continued)	532-541	YLWWVNGQSL	323	A0201	25	816
				A26	18	N.A.
	538-546	GQSLPVSPR	324	B2705	17	100

Table 29A
HER2/NEU: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
Her-2 25-52	30-37	DMKLRIPA	325	B08	19	8
	28-37	GTDMKLRIPA	326	A1	23	6
	42-49	HLDMLRHL	327	B08	17	<5
	41-49	THLDMLRHL	328	A0201	17	<5
				B1510	24	N.A.
	40-49	ETHLDMLRHL	329	A26	29	N.A.
	36-43	PASPETHL	330	B5101	17	N.A.
				A0201	15	<5
	35-43	LPASPETHL	331	B5101	20	130
				B5102	N.A.	100
	34-43	RLPASPETHL	332	A0201	20	21
				A0201	15	<5
	38-46	SPETHLDML	333	B0702	20	24
				B08	18	<5
	37-46	ASPETHLDML	334	B5101	18	110
				A0201	18	<5
	42-50	HLDMLRHLY	335	A1	29	25
				A26	20	N.A.
	41-50	THLDMLRHLY	336	A3	17	4
				A1	18	<1.0

Table 29B
HER2/NEU: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
Her-2 705-732	719-726	ELRKVKVL	337	B08	24	16
	718-726	TELKVKVL	338	A0201	16	1
				B08	22	<5
				B5101	16	<5
	717-726	ETELRKVKVL	339	A1	18	2
				A26	28	6
				A0201	17	<5
	715-723	LKETELRKV	340	B5101	15	<5
				A0201	29	8
	714-723	ILKETELRKV	341	A0201	15	<5
				B08	22	<5
				B2705	27	2000
	712-720	MRILKETEL	342	B2709	21	N.A.
				A0201	20	2
				B0702	13	40
	711-720	QMRILKETEL	343	A1	18	5
	717-725	ETELRKVKV	344	A26	18	N.A.
				A0201	16	19
	716-725	KFTELKVKV	345	B0702	16	8
	706-714	MPNQAQMRI	346	B5101	22	629
	705-714	AMPNQAQMRI	347	A0201	18	8
	706-715	MPNQAQMRIL	348	B0702	20	80

Table 29C
HER2/NEU: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
Her-2 954-982	966-973	RPRFREL V	349	B08	20	24
				B5101	18	N.A.
	965-973	CRPRFREL V	350	B2709	18	
				A26	25	N.A.
				A24	N.A.	32
	968-976	RFREL VSEF	351	A3	15	<5
				B08	16	<5
				B2705	19	
	967-976	PRFREL VSEF	352	A26	18	N.A.
				A26	21	N.A.
				A24	N.A.	6
	964-972	ECRPRFREL	353	B0702	15	40
				B8	27	640
				B1510	16	<5

Table 30
NY-ESO-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
NY-ESO-1 51-77	67-75	GAASGLNGC	354	A0201	15	<5
	52-60	RASGPGGGA	355	B0702	15	<5
	64-72	PHGGAASGL	356	B1510	21	N.A.
	63-72	GPHGGAASGL	357	B0702	22	80
	60-69	APRGPHGGAA	358	B0702	23	60

Table 31A
PRAME: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
PRAME 103-135	112-119	VRPRRWKL	359	B08	19	
	111-119	EVRPRRWKL	360	A26	27	N.A.
				A24	N.A.	5
				A3	19	N.A.
				B0702	15	(B7) 300.00
	113-121	RPRRWKLQV	361	B08	26	160
				B0702	21	(B7) 40.00
	114-122	PRRWKLQVL	362	B5101	19	110
				B08	26	<5
				B2705	23	200
PRAME 161-187	113-122	RPRRWKLQVL	363	B0702	24	(B7) 800.00
				B8	N.A.	160
				B5101	N.A.	61
				B5102	N.A.	61
	116-124	RWKLQVLDL	364	A24	N.A.	10
				B08	22	<5
				B2705	17	3
				A0201	16	<5
	115-124	RRWKLQVLDL	365			
	174-182	PVEVLVDLF	366	A26	25	N.A.

Table 31B
PRAME: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
PRAME 185-215	199-206	VKRKKNVL	367	B08	27	8
	198-206	KVKKKKNVL	368	A0201	16	<1.0
				A26	20	N.A.
				A3	22	<1.0
				B08	30	40
	197-206	EKVKRKKNVL	369	B2705	16	
				A26	15	N.A.
				B08	20	6
				B08	20	<5
	200-208	KRKKNVRL	372	A0201	15	<1.0
				A26	15	N.A.
				B0702	15	<1.0
				B08	21	<1.0
	199-208	VKRKKNVRL	373	B2705	28	
				B2709	25	
				A0201	16	<1.0
				B0702	16	4
	189-196	DELFYLI	374	B5101	15	N.A.
				A0201	22	3
	205-213	VLRLOCKKL	375	A26	17	N.A.
				B08	25	8
	204-213	NVLRLOCKKL	376	A0201	17	7
				A26	19	N.A.

Table 31C

PRAME: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
PRAME 185-215 (continued)	194-202	YLIEKVKRK	377	A0201	20	<1.0
				A26	18	N.A.
				A3	25	68
				B08	20	<1.0
				B2705	17	
PRAME 71-98	74-81	QAWPFTCL	378	B5101	17	n.a.
	73-81	VQAWPFTCL	379	A0201	14	7
				A24	n.a.	5
				B0702	16	6
	72-81	MVQAWPFTCL	380	A26	22	n.a.
				A24	n.a.	7
				B0702	13	30
	81-88	LPLGVLMK	381	B5101	18	n.a.
	80-88	CLPLGVLMK	382	A0201	17	<1.0
				A3	27	120
	79-88	TCLPLGVLMK	383	A1	12	10
				A3	19	3
	84-92	GVLMKGQHL	384	A0201	18	7
				A26	21	n.a.
				B08	21	4
	81-89	LPLGVLMKG	385	B5101	20	2
	80-89	CLPLGVLMKG	386	A0201	16	<1.0
	76-85	WPFTCLPLGV	387	B0702	18	4

Table 31D
PRAME: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
PRAME 39-65	51-59	ELFPPLFMA	388	A0201	19	18
				A26	23	N.A.
	49-57	PRELFPPLF	389	B2705	22	
				B2709	19	
	48-57	LPRELFPPLF	390	B0702	19	4
	50-58	RELFPPLFM	391	B2705	16	
				B2705	15	
	49-58	PRELFPPLFM	392	A1	16	<1.0

Table 32
PSA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
PSA 232-258	239-246	RPSLYTKV	393	B5101	21	N.A.
	238-246	ERPSLYTKV	394	B2705	15	60
	236-243	LPERPSLY	395	B5101	18	N.A.
	235-243	ALPERPSLY	396	A1	19	<1.0
				A26	22	N.A.
				A3	26	6
				B08	16	<1.0
				B2705	11	15
				B2709	19	N.A.
				A0201	20	<1.0
	241-249	SLYTKVVHY	397	A1	19	<1.0
				A26	25	N.A.
				A3	26	60
				B08	20	<1.0
				B2705	13	75
	240-249	PSLYTKVVHY	398	A1	20	<1.0
	239-247	RPSLYTKVV	399	A26	16	N.A.
				B0702	21	4
				B5101	23	110

Table 33A

PSMA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
PSMA 202-228	211-218	GNKVKNQAQ	400	B08	22	<5
	202-209	IARYGKVF	401	B08	18	<5
	217-225	AQLAGAKGV	402	A0201	16	26
	207-215	KVFRGNKVK	403	A3	32	15
	211-219	GNKVKNQAQL	404	B8	33	80
PSMA 255-282				B2705	17	20
	269-277	TPGYPANEY	405	A1	16	<5
	268-277	LTPGYPANEY	406	A1	21	1
				A26	24	N.A.
	271-279	GYPANEYAY	407	A1	15	<5
	270-279	PGYPANEYAY	408	A1	19	<5
	266-274	DPLTPGYPA	409	B0702	21	3
PSMA 483-509				B5101	17	20
				A0201	17	<5
	492-500	SLYESWTKK	410	A3	27	150
				B2705	18	150
	491-500	KSLYESWTKK	411	A3	16	<5
				A1	19	<5
	486-494	EGFEGKSLY	412	A26	21	N.A.
				B2705	16	<5
	485-494	DEGFEGKSLY	413	A1	17	<5
				A26	17	N.A.
	498-506	TKKSPSPEF	414	B08	17	<5

Table 33B

PSMA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
PSMA 483-509 (continued)	497-506	WTKKSPSEF	415	A26	24	N.A.
	492-501	SLYESWTKKS	416	A0201	16	<5
				A3	16	<5
PSMA 721-749	725-732	WGEVKRQI	417	B08	17	<5
	724-732	AWGEVKRQI	418	B5101	17	N.A.
				B5101	15	6
	723-732	KAWGEVKRQI	419	A0201	16	<1.0
	723-730	KAWGEVKR	420	B5101	15	N.A.
	722-730	SKAWGEVKR	421	B2705	15	<5
	731-739	QIYVAAFTV	422	A0201	21	177
				A3	21	<1.0
				B5101	15	5
	733-741	YVAAFTVQA	423	A0201	17	6
PSMA 721-749	725-733	WGEVKRQIY	424	A3	20	<1.0
				A1	26	11
	727-735	EVKRQIYVA	425	A26	22	N.A.
				A3	18	<1.0
	738-746	TVQAAAETL	426	A26	18	N.A.
				A3	19	<1.0
	737-746	FTVQAAAETL	427	A0201	17	<1.0
				A26	19	N.A.

Table 33C
PSMA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
PSMA 721-749 (continued)	729-737	KRQIYVAAF	428	A26	16	N.A.
				B2705	24	3000
				B2709	21	N.A.
	721-729	PSKAWGEVK	429	A3	20	<1.0
PSMA 95-122	723-731	KAWGEVKRQ	430	B5101	16	<1.0
	100-108	WKEFGSDV	431	A0201	16	<5
	99-108	QWKEFGSDV	432	A0201	17	<5
	102-111	EFGLDSVELA	433	A26	16	N.A.

Table 34A
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 117-143	126-134	ELRQKESKL	434	A0201	20	<5
				A26	26	N.A.
				A3	17	<5
				B0702	13	(B7) 40.00
				B8	34	320
SCP-1 281-308	125-134	AELRQKESKL	435	A0201	16	<5
	133-141	KIQENRKII	436	A0201	20	61
	298-305	QLEEKTKL	437	B08	28	2
	297-305	NQLEEKTKL	438	A0201	16	33
				B2705	19	200
	288-296	LLEESRDKV	439	A0201	25	15
	287-296	FLLEESRDKV	440	B5101	15	3
				A0201	27	2378
	291-299	ESRDKVNQL	441	A26	21	N.A.
	290-299	EESRDKVNQL	442	B08	29	240
SCP-1 471-498				A26	19	N.A.
	475-483	EKEVHDLEY	443	A1	31	11
				A26	17	N.A.
	474-483	REKEYHDLEY	444	A1	21	<1.0
				A1	26	45
	480-488	DLEYSYCHY	445	A26	30	N.A.
				A3	16	<5
	477-485	EVHDLEYSY	446	A1	15	1

Table 34B
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 471-498 (continued)	477-485	EVHDL EYSY		A26	29	N.A.
	477-486	EVHDL EYSYC	447	A3	19	<1.0
	502-509	KLSSKREL	448	A26	22	N.A.
	508-515	ELKNTEYF	449	B08	26	4
	507-515	RELKNTEYF	450	B08	24	<1.0
SCP-1 493-520	496-503	RELKNTEYF	450	B2705	18	45
	494-503	KRGQRPKL	451	B4403	N.A.	120
		LPKRGQRPKL	452	B08	18	<1.0
	509-517	LKNTEYFTL	453	B0702	22	120
	508-517	ELKNTEYFTL	454	B8	N.A.	16
	506-514	KRELKNTEY	455	B5101	N.A.	130
	502-510	KLSSKRELK	456	B3501	N.A.	60
	498-506	GQRPKLSSK	457	A0201	15	<5
	497-506	RGQRPKLSSK	458	A0201	18	<1.0
	500-508	RPKLSSKRE	459	A26	27	N.A.
				A3	16	<1.0
				A1	26	2
				B2705	26	3000
				A3	25	60
				A3	22	4
				B2705	18	200
				A3	22	<1.0
				B08	18	<1.0

Table 34C
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 570-596	573-580	LEYVREEL	460	B08	19	<5
	572-580	ELEYVREEL	461	A0201	17	<1.0
				A26	23	N.A.
				A24	N.A.	9
				B08	20	N.A.
	571-580	N ELEYVREEL	462	A0201	16	4
	579-587	ELKQKRDEV	463	A0201	19	<1.0
				A26	18	N.A.
				B08	29	48
				A26	17	N.A.
SCP-1 618-645	575-583	YVREELKQK	464	A3	27	2
	632-640	QLNVYEKV	465	A0201	24	70
	630-638	SKQLNVYEI	466	A0201	17	<5
	628-636	AESKQLNVY	467	A1	19	<5
				A26	16	N.A.
	627-636	TAESKQLNVY	468	A1	26	45
				A26	15	N.A.

Table 34D
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 633-660	638-645	IKVNKLEL	469	B08	21	<1.0
	637-645	EIKVNKLEL	470	A0201	17	<1.0
				A26	26	N.A.
				B08	28	8
				B1510	15	N.A.
	636-645	YEIKVNKLEL	471	A0201	17	2
	642-650	KLELEESA	472	A0201	20	1
				A3	16	<1.0
				A0201	18	<1.0
	635-643	VYEIKVNKL	473	A24	N.A.	396
				B08	22	<1.0
				A0201	24	56
SCP-1 640-668	634-643	NVYEIKVNKL	474	A26	25	N.A.
				A24	N.A.	6
				A3	15	<5
				B0702	11	(B7) 20
				B08	N.A.	6
				A26	27	N.A.
	642-650	KLELEESA	476	A0201	20	1
				A3	16	<1.0
	646-654	ELESAKQKF	477	A26	27	N.A.

Table 34E
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYRPEITHI	NIH
SCP-1 768-796	771-778	KEKLRREA	478	B08	21	<5
				A0201	18	<5
				A26	18	N.A.
				A24	N.A.	5
	777-785	EAKENTATL	479	B0702	13	12
				B08	28	48
				B5101	20	121
	776-785	REAKENTATL	480	A0201	16	<5
	773-782	KLKREAKENT	481	A3	17	<5
	112-119	EAEKIKKW	482	B5101	17	N.A.
SCP-1 92-125				A0201	23	32
				A26	22	N.A.
	101-109	GLSRVYSKL	483	A24	N.A.	6
				A3	17	3
				B08	17	<1.0
				A26	21	N.A.
	100-109	EGLSRVYSKL	484	A24	N.A.	9
				A0201	22	57
	108-116	KLYKEAEKI	485	A3	20	9
				B5101	18	5
	98-106	NSEGLSRVY	486	A1	31	68
	97-106	ENSEGLSRVY	487	A26	18	N.A.
	102-110	LSRVYSKLY	488	A1	22	<1.0

Table 34F

SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 92-125 (continued)	101-110	GLSRVYSKLY	489	A1	18	<1.0
				A26	18	N.A.
				A3	19	18
	96-105	LENSEGLSRV	490	A0201	17	5
SCP-1 931-958	108-117	KLYKEAEKIK	491	A3	27	150
	949-956	REDRWAVI	492	B5101	15	N.A.
				B2705	18	600
	948-956	MREDRWAVI	493	B2709	18	N.A.
				B5101	15	1
	947-956	KMREDRWAVI	494	A0201	21	6
				B08	N.A.	15
	947-955	KMREDRWAV	495	A0201	22	411
	934-942	TTPGSTLKF	496	A26	25	N.A.
	933-942	LTPGSTLKF	497	A26	23	N.A.
SCP-1 232-259	937-945	GSTLKFGAI	498	B08	19	1
	945-953	IRKMREDRW	499	B08	19	<5
	236-243	RLEMHFKL	500	B08	16	<5
	235-243	SRLEMHFKL	501	A0201	18	<5
				B2705	25	2000
	242-250	KLKEDYFKI	502	B2709	22	
				A0201	22	4

Table 34G

SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 232-259 (continued)				A26	16	N.A.
				A3	15	3
				B08	24	<5
				B5101	14	2
				A1	15	<5
	249-257	KIQHLEQY	503	A26	23	N.A.
				A3	17	<5
	248-257	EKIQHLEQY	504	A1	15	<5
				A26	21	N.A.
	233-242	ENSRLEMHF	505	A26	19	N.A.
SCP-1 310-340	236-245	RIEMHFKLKE	506	A1	19	<5
				A3	17	<5
	324-331	LEDIKVSL	507	B08	20	<1.0
				A0201	21	<1.0
	323-331	ELEDIKVSL	508	A26	25	N.A.
				A24	N.A.	10
				A3	17	<1.0
				B08	19	<1.0
				B1510	16	N.A.
	322-331	KELEDIKVSL	509	A0201	19	22
	320-327	LTKELEDI	500	B08	18	<5
	319-327	HLTKLEEDI	511	A0201	21	<1.0
	330-338	SLQRSVSTQ	512	A0201	18	<1.0

Table 34H

SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 310-340 (continued)	321-329	TKELEDKV	513	A1	16	<1.0
	320-329	LTKLEDKV	514	A0201	19	<1.0
	326-335	DIKVSQRSV	515	A26	18	N.A.
	281-288	KMKDLTFL	516	B08	20	3
	280-288	NKMKDLTFL	517	A0201	15	1
SCP-1 272-305	279-288	ENKMKDLTFL	518	A26	19	N.A.
	288-296	LLEESRDV	519	A0201	25	15
	287-296	FLLEESRDV	520	B5101	15	3
	291-299	ESRDKVNQL	521	A0201	27	2378
	290-299	EESRDKVNQL	522	A26	21	N.A.
	277-285	EKENKMKDL	523	B08	29	240
	276-285	TEKENKMKDL	524	A26	19	N.A.
	279-287	ENKMKDLTF	525	A26	19	N.A.
	218-225	IEKMITAF	526	B08	28	4
	217-225	NIEKMITAF	527	B08	17	<5
SCP-1 211-239	216-225	SNIEKMITAF	528	A26	26	N.A.
	223-230	TAFEELRV	529	A26	19	N.A.
	222-230	ITAFEELRV	530	B5101	23	N.A.
	221-230	MITAFEELRV	531	A0201	18	2
				A0201	18	16

Table 34I
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 211-239 (continued)	220-228	KMTAFEEL	532	A0201	23	50
				A26	15	N.A.
				A24	N.A.	16
	219-228	EKMTAFEEL	533	A26	19	N.A.
	227-235	ELRVQAENS	534	A3	16	<1.0
				B08	15	<1.0
	213-222	DLNSNIEKMI	535	A0201	17	<1.0
				A26	16	N.A.
	837-844	WTSAKNIL	536	B08	20	4
				A0201	18	2
SCP-1 836-863	846-854	TPLPKAYTV	537	B0702	17	4
				B08	16	2
				B5101	25	220
	845-854	STPLPKAYTV	538	A0201	19	<5
	844-852	LSTPLPKAY	539	A1	23	8
				A1	16	<1.0
	843-852	TLSTPLPKAY	540	A26	19	N.A.
				A3	18	2
	842-850	NTLSTPLPK	541	A3	16	3
	841-850	KNLSTPLPK	542	A3	18	<1.0

Table 34J
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 819-845	828-835	ISKDKRDY	543	B08	21	3
				A26	21	N.A.
	826-835	HGISKDKRDY	544	A1	15	<5
	832-840	KRDYLTSA	545	B2705	16	600
	829-838	SKDKRDYLT	546	A1	18	<5
	279-286	ENKMKDLT	547	B08	22	8
				A0201	17	3
	260-268	EINDKEKQV	548	A26	19	N.A.
				B08	17	<5
				A0201	17	3
SCP-1 260-288	274-282	QITEKENKM	549	A26	22	N.A.
				B08	16	<5
				A0201	16	<1.0
	269-277	SLLLIQITE	550	A3	18	<1.0
				B08	21	<1.0
SCP-1 437-464	453-460	FEKIAEEL	551	B2705	15	
	452-460	QFEKIAEEL	552	A0201	16	56
	451-460	KQFEKIAEEL	553	B08	16	2
	449-456	DNKQFEKI	554	B5101	16	N.A.
	448-456	YDNKQFEKI	555	B5101	16	1
	447-456	LYDNKQFEKI	556	A1	15	<1.0

Table 34K
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 437-464 (continued)	440-447	LGEKETLL	557	B5101	16	N.A.
	439-447	VLGEKETLL	558	A0201	24	149
				A26	19	N.A.
				B08	29	12
	438-447	KVLGEKETLL	559	A0201	19	24
				A26	20	N.A.
				A24	N.A.	12
				A3	18	<1.0
				B0702	14	20
				A0201	22	3
SCP-1 383-412	390-398	LLRTEQQRL	560	A26	18	N.A.
				B08	22	1.6
				B2705	15	30
				A0201	19	6
	389-398	ELLRTEQQRL	561	A26	24	N.A.
				A3	15	<1.0
				A1	15	<5
	393-401	TEQQRLNY	562	A26	16	N.A.
	392-401	RTEQQRLNY	563	A1	31	113
				A26	26	N.A.
	402-410	EDQLILTM	564	A26	18	N.A.
	397-406	RLNYEDQLI	565	A0201	17	<1.0
				A3	15	<1.0

Table 34K
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 366-394	368-375	KARAAHSF	566	B08	16	<1.0
	376-384	VVTEFETTV	567	A0201	19	161
				A3	16	<1.0
	375-384	FVVTEFETTV	568	A0201	17	106
	377-385	VTEFETTV	569	A1	18	2
SCP-1 331-357	376-385	VVTEFETTV	570	A3	16	<5
	344-352	DLQIATNTI	571	A0201	22	<5
				A3	15	<1.0
				B5101	17	11
	347-355	IATNTICQL	572	A0201	19	1
				B08	16	<1.0
				B5101	20	79
	346-355	QIATNTICQL	573	A0201	24	7
				A26	24	N.A.

Table 35
SSX-4: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SSX4 45-76	57-65	VMTKLGFKV	574	A0201	21	495
	53-61	LNVEVMTKL	575	A0201	17	7
				A0201	23	172
				A26	21	N.A.
	52-61	KLNYEVMTKL	576	A24	N.A.	18
SSX4 98-124				A3	14	4
				B7	N.A.	4
	66-74	TLPPFMRSK	577	A26	16	N.A.
				A3	25	14
	110-118	KMPKKPAE	578	A0201	15	<5
SSX4 98-124				A26	15	N.A.
				A3	16	<5
	103-112	SLQRIFPKIM	579	A0201	15	8
				A26	16	N.A.
				A3	15	<5

Tyrosinase: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion				Binding Prediction		
Substrate	Epitope	Sequence	Seq. ID No.	HLA type	SYFPEITHI	NIH
Tyr 445-474	463-471	YKSYLEQA	580	A0201	18	<5
				A26	17	N.A.
	459-467	SFQDYKSY	581	A1	18	<5
				A26	22	N.A.
	458-467	DSFQDYKSY	582	A1	19	<5
				A26	24	N.A.
	507-514	LPEEKQPL	583	B08	28	5
				B5101	18	N.A.
				A0201	22	88
	506-514	QLPEEKQPL	584	A26	20	N.A.
				A24	N.A.	9
Tyr 490-518				B08	18	<5
	505-514	KQLPEEKQPL	585	A0201	15	28
				A24	N.A.	17
				A0201	15	<5
	507-515	LPEEKQPLL	586	B0702	21	24
				B08	28	5
				B5101	21	157
	506-515	QLPEEKQPLL	587	A0201	23	88
				A26	20	N.A.
	497-505	SLLCRHKRK	588	A24	N.A.	7
				A3	25	15

Example 15**Evaluating Likelihood of Epitope Cross-reactivity on Non-target Tissues.**

As noted above PSA is a member of the kallikrein family of proteases, which is itself a subset of the serine protease family. While the members of this family sharing the greatest degree of sequence identity with PSA also share similar expression profiles, it remains possible that individual epitope sequences might be shared with proteins having distinctly different expression profiles. A first step in evaluating the likelihood of undesirable cross-reactivity is the identification of shared sequences. One way to accomplish this is to conduct a BLAST search of an epitope sequence against the SWISSPROT or Entrez non-redundant peptide sequence databases using the "Search for short nearly exact matches" option; hypertext transfer protocol accessible on the world wide web (<http://www.ncbi.nlm.nih.gov/blast/index.html>). Thus searching SEQ ID NO. 214, WVLTAAHCI, against SWISSPROT (limited to entries for homo sapiens) one finds four exact matches, including PSA. The other three are from kallikrein 1 (tissue kallikrein), and elastase 2A and 2B. While these nine amino acid segments are identical, the flanking sequences are quite distinct, particularly on the C-terminal side, suggesting that processing may proceed differently and that thus the same epitope may not be liberated from these other proteins. (Please note that kallikrein naming is confused. Thus the kallikrein 1 [accession number P06870] is a different protein than the one [accession number AAD13817] mentioned in the paragraph on PSA above in the section on tumor-associated antigens).

It is possible to test this possibility in several ways. Synthetic peptides containing the epitope sequence embedded in the context of each of these proteins can be subjected to *in vitro* proteasomal digestion and analysis as described above. Alternatively, cells expressing these other proteins, whether by natural or recombinant expression, can be used as targets in a cytotoxicity (or similar) assay using CD8⁺ T cells that recognize the epitope, in order to determine if the epitope is processed and presented.

Example 16**Epitope Clusters.**

Known and predicted epitopes are generally not evenly distributed across the sequences of protein antigens. As referred to above, we have defined segments of sequence containing a higher than average density of (known or predicted) epitopes as epitope clusters. Among the uses of epitope clusters is the incorporation of their sequence into substrate peptides used in proteasomal digestion analysis as described herein. Epitope clusters can also be useful as vaccine components. A fuller discussion of the definition and uses of epitope clusters is found in U.S. Patent Application No. 09/561,571 entitled EPITOPE CLUSTERS, previously incorporated by reference.

The following tables (37-60) present 9-mer epitopes predicted for HLA-A2 binding using both the SYFPEITHI and NIH algorithms and the epitope density of regions of overlapping

epitopes, and of epitopes in the whole protein, and the ratio of these two densities. (The ratio must exceed one for there to be a cluster by the above definition; requiring higher values of this ratio reflect preferred embodiments). Individual 9-mers are ranked by score and identified by the position of their first amino in the complete protein sequence. Each potential cluster from a protein is numbered. The range of amino acid positions within the complete sequence that the cluster covers is indicated as are the rankings of the individual predicted epitopes it is made up of.

Table 37
BIMAS-NIH/Parker algorithm Results for gp100

Rank	Start	Score	Rank	Start	Score
1	619	1493	21	416	19
2	602	413	22	25	18
3	162	226	23	566	17
4	18	118	24	603	15
5	178	118	25	384	14
6	273	117	26	13	14
7	601	81	27	290	12
8	243	63	28	637	10
9	606	60	29	639	9
10	373	50	30	485	9
11	544	36	31	453	8
12	291	29	32	102	8
13	592	29	33	399	8
14	268	29	34	456	7
15	47	27	35	113	7
16	585	26	36	622	7
17	576	21	37	69	7
18	465	21	38	604	6
19	570	20	39	350	6
20	9	19	40	583	5

Table 38
SYFPEITHI (Rammensee algorithm) Results for gp100

Rank	Start	Score	Rank	Start	Score	Rank	Start	Score
1	606	30	37	291	20	73	60	18
2	162	29	38	269	20	74	17	18
3	456	28	39	2	20	75	613	17
4	18	28	40	610	19	76	599	17
5	602	27	41	594	19	77	572	17
6	598	27	42	591	19	78	557	17
7	601	26	43	583	19	79	556	17
8	597	26	44	570	19	80	512	17
9	13	26	45	488	19	81	406	17
10	585	25	46	446	19	82	324	17
11	449	25	47	322	19	83	290	17
12	4	25	48	267	19	84	101	17
13	603	24	49	250	19	85	95	17
14	576	24	50	205	19	86	635	16
15	453	24	51	180	19	87	588	16
16	178	24	52	169	19	88	584	16
17	171	24	53	88	19	89	577	16
18	11	24	54	47	19	90	559	16
19	619	23	55	10	19	91	539	16
20	280	23	56	648	18	92	494	16
21	268	23	57	605	18	93	482	16
22	592	22	58	604	18	94	468	16
23	544	22	59	595	18	95	442	16
24	465	22	60	571	18	96	413	16
25	399	22	61	569	18	97	408	16
26	373	22	62	450	18	98	402	16
27	273	22	63	409	18	99	286	16
28	243	22	64	400	18	100	234	16
29	566	21	65	371	18	101	217	16
30	563	21	66	343	18	102	211	16
31	485	21	67	298	18	103	176	16
32	384	21	68	209	18	104	107	16
33	350	21	69	102	18	105	96	16
34	9	21	70	97	18	106	80	16
35	463	20	71	76	18	107	16	16
36	397	20	72	69	18	108	14	16
						109	7	16

Table 39

Prediction of clusters for gp100

Total AAs: 661

Total 9-mers: 653

SYFPEITHI 16: 109 9-mers

NIH 5: 40 9-mers

				Epitopes/AA		
	Cluster #	AAs	Epitopes (by Rank)	Cluster	Whole Pr	Ratio
SYFPEITHI	1	2 to 26	39, 12, 109, 34, 55, 11, 9, 108, 107, 74, 4	0.440	0.165	2.668
	2	69-115	72, 71, 106, 53, 85, 105, 70, 84, 69, 104	0.213	0.165	1.290
	3	95-115	85, 105, 70, 84, 69	0.238	0.165	1.444
	4	162-188	2, 52, 17, 103, 16, 51	0.222	0.165	1.348
	5	205-225	50, 68, 102, 101	0.190	0.165	1.155
	6	243-258	28, 49	0.125	0.165	0.758
	7	267-306	48, 21, 38, 27, 20, 99, 83, 37, 67	0.225	0.165	1.364
	8	322-332	47, 82	0.182	0.165	1.103
	9	343-358	66, 33	0.125	0.165	0.758
	10	371-381	65, 26	0.182	0.165	1.103
	11	397-421	36, 25, 64, 98, 81, 97, 63, 96	0.320	0.165	1.941
	12	442-476	95, 46, 11, 62, 15, 3, 35, 24, 94	0.257	0.165	1.559
	13	482-502	93, 31, 45, 93	0.190	0.165	1.155
	14	539-552	91, 23	0.143	0.165	0.866
	15	556-627	79, 78, 90, 30, 29, 61, 44, 60, 77, 14, 89, 43, 88, 10, 87, 42, 22, 41, 59, 8, 6, 76, 7, 5, 13, 58, 57, 1, 40, 75, 19	0.431	0.165	2.611
NIH	1	9 to 33	20, 26, 4, 22	0.160	0.061	2.644
	2	268-281	14, 6	0.143	0.061	2.361
	3	290-299	27, 12	0.200	0.061	3.305
	4*	102-121	32, 35	0.100	0.061	1.653
	5*	373-392	10, 25	0.100	0.061	1.653
	6	453-473	31, 34, 18	0.143	0.061	2.361
	7	566-600	23, 19, 17, 40, 16, 13	0.171	0.061	2.833
	8	601-614	7, 2, 24, 38, 9	0.357	0.061	5.902
	9	619-630	1, 36	0.17	0.061	2.754
	10	637-647	28, 29	0.18	0.061	3.005

*Nearby but not overlapping epitopes

Table 40
BIMAS-NIH/Parker algorithm Results for PSMA

Rank	Start	Score
1	663	1360
2	711	1055
3	4	485
4	27	400
5	26	375
6	668	261
7	707	251
8	469	193
9	731	177
10	35	67
11	33	64
12	554	59
13	427	50
14	115	47
15	20	40
16	217	26
17	583	24
18	415	19
19	193	14
20	240	12
21	627	11
22	260	10
23	130	10
24	741	9
25	3	9
26	733	8
27	726	7
28	286	6
29	174	5
30	700	5

Table 41
SYFPEITHI (Rammensee algorithm) Results for PSMA

Rank	Start	Score	Rank	Start	Score	Rank	Start	Score
1	469	27	31	26	20	61	305	17
2	27	27	32	3	20	62	304	17
3	741	26	33	583	19	63	286	17
4	711	26	34	579	19	64	282	17
5	354	25	35	554	19	65	169	17
6	4	25	36	550	19	66	142	17
7	663	24	37	547	19	67	122	17
8	130	24	38	390	19	68	738	16
9	57	24	39	219	19	69	634	16
10	707	23	40	193	19	70	631	16
11	260	23	41	700	18	71	515	16
12	20	23	42	472	18	72	456	16
13	603	22	43	364	18	73	440	16
14	218	22	44	317	18	74	385	16
15	109	22	45	253	18	75	373	16
16	731	21	46	91	18	76	365	16
17	668	21	47	61	18	77	361	16
18	660	21	48	13	18	78	289	16
19	507	21	49	733	17	79	278	16
20	454	21	50	673	17	80	258	16
21	427	21	51	671	17	81	247	16
22	358	21	52	642	17	82	217	16
23	284	21	53	571	17	83	107	16
24	115	21	54	492	17	84	100	16
25	33	21	55	442	17	85	75	16
26	606	20	56	441	17	86	37	16
27	568	20	57	397	17	87	30	16
28	473	20	58	391	17	88	21	16
29	461	20	59	357	17			
30	200	20	60	344	17			

Table 42

Prediction of clusters for prostate-specific membrane antigen (PSMA)

Total AAs: 750

Total 9-mers: 742

SYFPEITHI 16: 88 9-mers

NIH 5: 30 9-mers

	Cluster #	Aas	Eptopes (by rank)	Eptopes/AA		
				Cluster	Whole Pr	Ratio
SYFPEITHI	1	3 to 12	32, 6	0.200	0.117	1.705
	2	13-45	13, 12, 88, 31, 2, 87, 25, 86	0.242	0.117	2.066
	3	57-69	9, 47	0.154	0.117	1.311
	4	100-138	84, 83, 15, 24, 67, 8	0.154	0.117	1.311
	5	193-208	40, 30	0.111	0.117	0.947
	6	217-227	82, 14, 39	0.273	0.117	2.324
	7	247-268	81, 45, 80, 11	0.182	0.117	1.550
	8	278-297	79, 64, 23, 63, 78	0.250	0.117	2.131
	9	354-381	5, 59, 22, 77, 43, 76, 75	0.250	0.117	2.131
	10	385-405	74, 38, 58, 57	0.190	0.117	1.623
	11	440-450	73, 56, 55	0.273	0.117	2.324
	12	454-481	20, 72, 29, 1, 42, 28	0.214	0.117	1.826
	13	507-523	17, 71	0.118	0.117	1.003
	14	547-562	37, 36, 35	0.188	0.117	1.598
	15	568-591	27, 53, 34, 33	0.167	0.117	1.420
	16	603-614	13, 26	0.167	0.117	1.420
	17	631-650	70, 69, 52	0.150	0.117	1.278
	18	660-681	18, 7, 17, 51, 50	0.227	0.117	1.937
	19	700-719	41, 10, 4	0.150	0.117	1.278
	20	731-749	16, 49, 68, 3	0.211	0.117	1.794
NIH	1	3 to 12	25, 3	0.200	0.040	5.000
	2	20-43	15, 5, 4, 11, 10	0.208	0.040	5.208
	3*	415-435	18, 13	0.095	0.040	2.381
	4	663-676	1, 6	0.143	0.040	3.571
	5	700-715	30, 7, 3	0.188	0.040	4.688
	6	726-749	27, 9, 26, 24	0.167	0.040	4.167

*Nearby but not overlapping epitopes

Table 43
BIMAS-NIH/Parker algorithm Results for PSA

Rank	Start	Score
1	7	607
2	170	243
3	52	124
4	53	112
5	195	101
6	165	23
7	72	18
8	245	18
9	2	16
10	59	16
11	122	15
12	125	15
13	191	13
14	9	8
15	14	6
16	175	5
17	130	5

Table 44
SYFPEITHI (Rammensee algorithm) Results for PSA

Rank	Start	Score
1	72	26
2	170	22
3	53	22
4	7	22
5	234	21
6	166	21
7	140	21
8	66	21
9	241	20
10	175	20
11	12	20
12	41	19
13	20	19
14	14	19
15	130	18
16	124	18
17	121	18
18	47	18
19	17	18
20	218	17
21	133	17
22	125	17
23	122	17
24	118	17
25	110	17
26	67	17
27	52	17
28	21	17
29	16	17
30	2	17
31	184	16
32	179	16
33	158	16
34	79	16
35	73	16
36	4	16

Table 45

Prediction of clusters for prostate specific antigen (PSA)

Total AAs: 261

Total 9-mers: 253

SYFPEITHI 16: 36 9-mers

NIH 5: 17 9-mers

	Cluster #	AAs	Epitopes (by rank)	Epitopes/AA		
				Cluster	Whole Pr	Ratio
SYFPEITHI	1	2 to 29	30, 36, 4, 11, 14, 29, 19, 13, 28	0.321	0.138	2.330
	2	41-61	12, 18, 27, 3	0.190	0.138	1.381
	3	66-87	8, 26, 1, 35, 34	0.227	0.138	1.648
	4	110-148	25, 24, 17, 23, 16, 22, 15, 21, 7	0.184	0.138	1.332
	5	158-192	33, 6, 2, 10, 32, 31	0.171	0.138	1.243
	6	234-249	5, 9	0.125	0.138	0.906
	7*	118-133	24, 17, 23, 16, 22	0.313	0.138	2.266
	8*	118-138	24, 17, 23, 16, 22, 15	0.286	0.138	2.071
NIH	1	2-22	9, 1, 14, 15	0.190	0.065	2.924
	2	52-67	3, 4, 10	0.188	0.065	2.879
	3	122-138	11, 12, 17	0.176	0.065	2.709
	4	165-183	6, 2, 16	0.158	0.065	2.424
	5	191-203	13, 5	0.154	0.065	2.362
	6**	52-80	3, 4, 10, 7	0.138	0.065	2.118

*These clusters are internal to the less preferred cluster #4.

**Includes a nearby but not overlapping epitope.

Table 46
BIMAS-NIH/Parker algorithm Results for PSCA

Rank	Start	Score
1	43	153
2	5	84
3	7	79
4	109	36
5	105	105
6	108	24
7	14	21
8	20	18
9	115	17
10	42	15
11	36	15
12	99	9
13	58	8

20

Table 47
SYFPEITHI (Rammensee algorithm) Results for PSCA

25

Rank	Start	Score	Rank	Start	Score
1	108	30	17	54	19
2	14	30	18	12	19
3	105	29	19	4	19
4	5	28	20	1	19
5	115	26	21	112	18
6	99	26	22	101	18
7	7	26	23	98	18
8	109	24	24	51	18
9	53	23	25	43	18
10	107	21	26	106	17
11	20	21	27	104	17
12	8	21	28	83	17
13	13	20	29	63	17
14	102	19	30	50	17
15	60	19	31	3	17
16	57	19	32	9	16
			33	92	16

Table 48

Prediction of clusters for prostate stem cell antigen (PSCA)

Total AAs: 123

Total 9-mers: 115

SYFPEITHI 16: 33;

SYFPEITHI 20: 13

NIH 5: 13

	Cluster #	AAs	Epitopes (by rank)	Epitopes/AA		
				Cluster	Whole Pr.	Ratio
SYFPEITHI >16	1	1 to 28	20, 31, 19, 4, 7, 12, 33, 18, 13, 2, 11	0.393	0.268	1.464
	2	43-71	25, 30, 24, 9, 17, 16, 15, 29	0.276	0.268	1.028
	3	92-123	32, 23, 6, 27, 14, 22, 3, 26, 10, 1, 8, 21, 5	0.406	0.268	1.514
SYFPEITHI >20	1	5 to 28	4, 7, 12, 13, 2, 11	0.250	0.106	2.365
	2	99-123	6, 3, 10, 1, 8, 5	0.240	0.106	2.271
NIH	1	5 to 28	2, 3, 7, 8	0.167	0.106	1.577
	2	36-51	11, 10, 1	0.188	0.106	1.774
	3	99-123	12, 5, 6, 4, 9	0.200	0.106	1.892
	4*	105-116	5, 6, 4	0.250	0.106	2.365

*This cluster is internal to the less preferred cluster #3.

In tables 49-60 epitope prediction and cluster analysis data for each algorithm are presented together in a single table.

Table 49

Prediction of clusters for MAGE-1 (NIH algorithm)

Total AAs: 309

Total 9-mers: 301

NIH 5:19 9-mers

Cluster #	AAs	Epitope Rank	Start Position	NIH Score	Cluster	Epitopes/AA Whole Pr.	Ratio
1	18-32	16 19	18 24	9 7	0.133	0.063	2.112
2	101-113	14 7	101 105	11 44	0.154	0.063	2.442
3	146-159	9 3	146 151	32 169	0.143	0.063	2.263
4	169-202	10 13 18 17 6 5	169 174 181 187 188 194	32 16 8 8 74 110	0.176	0.063	2.796
5	264-277	2 12	264 269	190 20	0.143	0.063	2.263
6	278-290	1 11	278 282	743 28	0.154	0.063	2.437

Table 50

Prediction of clusters for MAGE-1 (SYFPEITHI algorithm)

Total AAs: 309

Total 9-mers: 301

SYFPEITHI 16: 46 9-mers

Cluster #	Aas	Epitope Rank	Start Position	SYFPEITHI Score	Epitopes/AA		
					Cluster	Whole	Ratio
1	7-49	22	7	19	0.233	0.153	1.522
		9	15	22			
		27	18	18			
		16	20	20			
		28	22	18			
		29	24	18			
		33	31	17			
		30	35	18			
		2	38	26			
		17	41	20			
2	89-132	10	89	22	0.273	0.153	1.783
		18	92	20			
		7	93	23			
		23	96	19			
		43	98	16			
		4	101	25			
		8	105	23			
		34	107	17			
		35	108	17			
		36	113	17			
		37	118	17			
		19	124	20			
3	167-203	44	167	16	0.270	0.153	1.766
		20	169	20			
		12	174	21			
		24	181	19			
		6	187	24			
		31	188	18			
		25	191	19			
		38	192	17			
		1	194	27			
		13	195	21			
4	230-246	14	230	21	0.118	0.153	0.769
		39	238	17			
5	264-297	15	264	21	0.235	0.153	1.538
		32	269	18			
		40	270	17			
		26	271	19			
		46	275	16			
		3	278	26			
		21	282	20			
		41	289	17			

Table 51

Prediction of clusters for MAGE-2 (NIH algorithm)

Total AAs: 314

Total 9-mers: 308

NIH \geq 5: 20 9-mers

Cluster #	AAs	Epitope Rank	Start Position	NIH Score	Cluster	Epitope/AA Whole Pr.	Ratio
1	101-120	18	101	5.373	0.150	0.065	2.310
		16	108	6.756			
		1	112	2800.697			
2	153-167	8	153	31.883	0.200	0.065	3.080
		4	158	168.552			
		7	159	32.138			
3	169-211	14	169	8.535	0.209	0.065	3.223
		19	174	5.346			
		6	176	49.993			
		11	181	15.701			
		15	188	7.536			
		12	195	12.809			
		5	200	88.783			
		10	201	16.725			
4	271-284	3	271	398.324	0.143	0.065	2.200
		9	276	19.658			

Table 52

Prediction of clusters for MAGE-2 (SYFPEITHI algorithm)

Total AAs: 314

Total 9-mers: 308

SYFPEITHI 16: 52 9-mers

Cluster #	AAs	Enitone Rank	Start Position	SYFPEITHI Score	Cluster	Enitones/AA Whole Pr.	Ratio
1	15-32	13	15	21	0.278	0.169	1.645
		29	18	18			
		43	20	16			
		30	22	18			
		21	24	19			
2	37-56	31	37	18	0.250	0.169	1.481
		16	40	20			
		44	44	16			
		14	45	21			
		22	48	19			
3	96-133	36	96	17	0.211	0.169	1.247
		46	101	16			
		6	108	25			
		47	109	16			
		2	112	27			
		37	120	17			
		38	125	17			
		17	131	20			
4	153-216	12	153	22	0.344	0.169	2.036
		39	158	17			
		7	159	25			
		23	161	19			
		24	162	19			
		48	164	16			
		49	167	16			
		32	170	18			
		50	171	16			
		4	174	26			
		9	176	24			
		51	177	16			
		15	181	21			
		25	188	19			
		18	194	20			
		33	195	18			
		19	198	20			
		3	200	27			
		1	201	28			
		40	202	17			
		10	203	23			
		52	208	16			
5	237-254	26	237	19	0.167	0.169	0.987
		27	245	19			
		34	246	18			
6	271-299	8	271	25	0.241	0.169	1.430
		35	276	18			
		41	277	17			
		11	278	23			
		28	283	19			
		20	285	20			
		42	291	17			

Table 53

Prediction of clusters for MAGE-3 (NIH algorithm)

Total AAs: 314

Total 9-mers: 308

NIH 5: 22 9-mers

Cluster #	AAs	Epitope Rank	Start Position	NIH Score	Cluster	Epitopes/AA Whole Pr.	Ratio
1	101-120	15	101	11.002	0.200	0.071	2.800
		21	105	6.488			
		8	108	49.134			
		2	112	339.313			
2	153-167	18	153	7.776	0.200	0.071	2.800
		6	158	51.77			
		22	159	5.599			
3	174-209	17	174	8.832	0.194	0.071	2.722
		7	176	49.993			
		13	181	15.701			
		19	188	7.536			
		14	195	12.809			
		5	200	88.783			
4	237-251	12	201	16.725	0.200	0.071	2.800
		16	237	10.868			
		4	238	148.896			
5	271-284	20	243	6.88	0.143	0.071	2.000
		1	271	2655.495			
		11	276	19.658			

Table 54

Prediction of clusters for MAGE-3 (SYFPEITHI algorithm)

Total AAs: 314

Total 9-mers: 308

SYFPEITHI 16: 47 9-mers

Cluster #	AAs	Epitope Rank	Start Position	SYFPEITHI Score	Cluster	Epitones/AA Whole Pr.	Ratio
1	15-32	12	15	21	0.278	0.153	1.820
		26	18	18			
		37	20	16			
		27	22	18			
		18	24	19			
2	38-56	38	38	16	0.263	0.153	1.725
		15	40	20			
		39	44	16			
		13	45	21			
3	101-142	19	48	19	0.190	0.153	1.248
		28	101	18			
		40	105	16			
		1	108	31			
		6	112	25			
		31	120	17			
		32	125	17			
		16	131	20			
4	153-216	41	134	16	0.313	0.153	2.048
		20	153	19			
		29	156	18			
		33	158	17			
		21	159	19			
		34	161	17			
		42	164	16			
		43	167	16			
		10	174	22			
		8	176	23			
		14	181	21			
		22	188	19			
		44	193	16			
		11	194	22			
		23	195	19			
		45	197	16			
		17	198	20			
		3	200	27			
		2	201	28			
		35	202	17			
		46	208	16			
5	220-230	5	220	26	0.182	0.153	1.191
		47	222	16			
6	237-246	7	237	25	0.200	0.153	1.311
		9	238	23			
7	271-293	4	271	27	0.217	0.153	1.425
		30	276	18			
		24	278	19			
		36	283	17			
		25	285	19			

Table 55

Prediction of clusters for PRAME (NIH algorithm)

Total AAs: 509

Total 9-mers: 501

NIH 5: 40 9-mers

Cluster #	AAs	Epitope Rank	Start Position	NIH Score	Cluster	Epitopes/AA Whole Pr.	Ratio
1	33-47	20	33	18	0.133	0.080	1.670
		17	39	21			
2	71-81	9	71	50	0.2	0.07984	2.505
		32	73	7			
3	99-108	23	100	15	0.2	0.07984	2.505
		24	99	13			
4	126-135	38	126	5	0.2	0.07984	2.505
		35	127	6			
5	224-246	5	224	124	0.130	0.080	1.634
		8	230	63			
		39	238	5			
6	290-303	18	290	18	0.214	0.080	2.684
		14	292	23			
		7	295	66			
7	305-324	28	305	10	0.200	0.080	2.505
		30	308	8			
		25	312	13			
		36	316	6			
8	394-409	2	394	182	0.188	0.080	2.348
		12	397	42			
		31	401	7			
9	422-443	10	422	49	0.227	0.080	2.847
		3	425	182			
		34	431	7			
		29	432	9			
		4	435	160			
10	459-487	15	459	21	0.172	0.080	2.159
		11	462	45			
		22	466	15			
		40	472	5			
		37	479	6			

Table 56

Prediction of clusters for PRAME (SYFPEITHI algorithm)

Total AAs: 509

Total 9-mers: 501

SYFPEITHI 17: 80 9-mers

Cluster #	AAs	Epitope Rank	Start Position	SYFPEITHI Score	Cluster	Epitopes/AA Whole Pr.	Ratio
1	18-59	65	18	17	0.238	0.160	1.491
		50	21	18			
		66	26	17			
		35	33	20			
		22	34	22			
		51	37	18			
		5	39	27			
		23	40	22			
		13	44	24			
		46	51	19			
2	78-115	36	78	20	0.263	0.160	1.648
		67	80	17			
		52	84	18			
		24	86	22			
		53	91	18			
		25	93	22			
		9	99	25			
		8	100	26			
		54	103	18			
		55	107	18			
3	191-202	56	191	18	0.167	0.160	1.044
		38	194	20			
4	205-215	26	205	22	0.182	0.160	1.139
		27	207	22			
5	222-238	47	222	19	0.235	0.160	1.474
		14	224	24			
		69	227	17			
		57	230	18			
6	241-273	70	241	17	0.212	0.160	1.328
		15	248	24			
		71	255	17			
		30	258	21			
		39	259	20			
		58	261	18			
7	290-342	40	265	20	0.208	0.160	1.300
		72	290	17			
		48	293	19			
		31	298	21			
		73	301	17			
		18	305	23			
		6	308	27			
		10	312	25			
		19	316	23			
		28	319	22			

Prediction of clusters for PRAME (SYFPEITHI algorithm)

Total AAs: 509

Total 9-mers: 501

SYFPEITHI 17: 80 9-mers

Cluster #	AAs	Epitope Rank	Start Position	SYFPEITHI Score	Cluster	Epitopes/AA Whole Pr.	Ratio
8	343-363	41	326	20	0.238	0.160	1.491
		74	334	17			
		59	343	18			
		60	348	18			
		75	351	17			
		20	353	23			
9	364-447	76	355	17	0.250	0.160	1.566
		49	364	19			
		32	371	21			
		11	372	25			
		61	375	18			
		77	382	17			
		21	390	23			
		78	391	17			
		1	394	30			
		42	397	20			
		62	403	18			
		33	410	21			
		43	418	20			
		34	419	21			
		7	422	27			
		2	425	29			
		79	426	17			
		63	428	18			
		64	431	18			
		12	432	25			
10	455-474	16	435	24	0.200	0.160	1.253
		80	439	17			
		29	455	22			
		17	459	24			
		4	462	28			
		3	466	29			

Table 57
Predication of clusters for CEA (NIH algorithm)

Total AAs:702

Total 9-mers: 694

NIH 5: 30 9-mers

Cluster #	AA	Peptides Rank	Start Position	Score	Cluster	Peptides/AAs Whole Pr.	Ratio
1	17-32	5	17	79.041	0.188	0.043	4.388
		7	18	46.873			
		20	24	12.668			
2	113-129	2	113	167.991	0.118	0.043	2.753
		15	121	21.362			
3	172-187	25	172	9.165	0.125	0.043	2.925
		14	179	27.995			
4	278-291	30	278	5.818	0.143	0.043	3.343
		17	283	19.301			
5	350-365	9	350	43.075	0.125	0.043	2.925
		12	357	27.995			
6	528-543	8	528	43.075	0.125	0.043	2.925
		13	535	27.995			
7	631-645	23	631	9.563	0.200	0.043	4.680
		19	634	13.381			
		24	637	9.245			
8	691-702	1	691	196.407	0.167	0.043	3.900
		27	694	7.769			

Table 58
Predication of clusters for CEA (SYFPEITHI algorithm)

Total AAs: 702

Total 9-mers: 694

SYFPEITHI 16: 81 9-mers

Cluster #	AA	Peptides Rank	Start Position	Score	Cluster	Peptides/AAs Whole Pr.	Ratio
1	5-36	67	5	16	0.250	0.117	2.140
		23	12	19			
		24	16	19			
		9	17	22			
		25	18	19			
		32	19	18			
		68	23	16			
		33	28	18			
2	37-62	41	37	17	0.269	0.117	2.305
		20	44	20			
		26	45	19			
		42	46	17			
		27	50	19			
		43	53	17			
3	99-115	44	54	17	0.235	0.117	2.014
		14	99	21			
		5	100	23			
		45	104	17			
4	116-129	34	107	18	0.143	0.117	1.223
		69	116	16			
5	172-187	21	121	20	0.125	0.117	1.070
		46	172	17			
6	192-202	70	179	16	0.182	0.117	1.557
		3	192	24			
7	226-241	47	194	17	0.188	0.117	1.605
		48	226	17			
		49	229	17			
8	307-318	15	233	21	0.250	0.117	2.140
		11	307	22			
		71	308	16			
9	319-349	51	310	17	0.129	0.117	1.105
		52	319	17			
		53	327	17			
		72	335	16			
10	370-388	35	341	18	0.211	0.117	1.802
		12	370	22			
		54	372	17			
		74	375	16			
11	403-419	6	380	23	0.235	0.117	2.014
		56	403	17			
		57	404	17			
		58	407	17			
		28	411	19			

12	427-442	59	427	17	0.188	0.117	1.605
		75	432	16			
		76	434	16			
13	450-462	77	450	16	0.154	0.117	1.317
		13	454	22			
14	488-505	36	488	18	0.167	0.117	1.427
		18	492	21			
		60	497	17			
15	548-558	4	548	24	0.182	0.117	1.557
		61	550	17			
16	565-577	62	565	17	0.154	0.117	1.317
		19	569	21			
17	579-597	78	579	16	0.143	0.117	1.223
		79	582	16			
		7	589	23			
18	605-618	2	605	25	0.143	0.117	1.223
		38	610	18			
19	631-669	29	631	19	0.154	0.117	1.317
		63	637	17			
		80	644	16			
		64	652	17			
		39	660	18			
		81	661	16			
20	675-702	22	675	20	0.286	0.117	2.446
		30	683	19			
		31	687	19			
		40	688	18			
		65	690	17			
		1	691	31			
		66	692	17			
		8	694	23			

Table 59
Predication of clusters for SCP-1 (NIH algorithm)

Total AAs: 976
 Total 9-mers: 968
 NIH 5: 37 9-mers

Cluster #	AA	Peptides Rank	Start Position	Score	Cluster	Peptides/AAs Whole Pr.	Ratio
1	101-116	15	101	40.589	0.125	0.038	3.270
		13	108	57.255			
2*	281-305	14	281	44.944	0.12	0.038	3.139
		24	288	15.203			
		17	297	32.857			
3	431-447	8	431	80.217	0.073	0.038	1.914
		26	438	11.861			
		4	439	148.896			
4	557-579	11	557	64.335	0.174	0.038	4.550
		19	560	24.937			
		6	564	87.586			
		18	571	32.765			
5	635-650	10	635	69.552	0.125	0.038	3.270
		34	642	6.542			
6	755-767	36	755	5.599	0.154	0.038	4.025
		35	759	5.928			
7	838-854	2	838	284.517	0.118	0.038	3.078
		28	846	11.426			

Table 60
Predication of clusters for SCP-1

Total AAs: 976

Total 9-mers: 968

Rammensee 16: 118 9-mers

Cluster #	AA	Peptides Rank	Start Position	Score	Cluster	Peptides/AAs Whole Pr.	Ratio
1	8-28	99	8	16	0.143	0.121	1.182
		77	15	17			
		100	20	16			
2	63-80	78	63	17	0.222	0.121	1.838
		50	66	19			
		102	69	16			
		60	72	18			
3	94-123	79	94	17	0.133	0.121	1.103
		12	101	23			
		17	108	22			
		103	115	16			
4	126-158	35	126	20	0.182	0.121	1.504
		36	133	20			
		51	139	19			
		80	140	17			
		61	143	18			
		37	150	20			
5	161-189	38	161	20	0.207	0.121	1.711
		52	165	19			
		81	171	17			
		82	177	17			
		62	178	18			
		39	181	20			
6	213-230	40	213	20	0.167	0.121	1.379
		13	220	23			
		28	222	21			
7	235-250	63	235	18	0.125	0.121	1.034
		18	242	22			
8	260-296	83	260	17	0.243	0.121	2.012
		105	262	16			
		84	267	17			
		106	269	16			
		41	270	20			
		64	271	18			
		85	274	17			
		19	281	22			
		3	288	25			
9	312-338	108	312	16	0.148	0.121	1.225
		29	319	21			
		30	323	21			
		65	330	18			
10	339-355	66	339	18	0.235	0.121	1.946
		31	340	21			
		42	344	20			
		53	347	19			

11	376-447	54	376	19	0.194	0.121	1.608
		43	382	20			
		44	386	20			
		20	390	22			
		55	397	19			
		6	404	24			
		86	407	17			
		45	411	20			
		67	417	18			
		21	425	22			
		46	431	20			
		68	432	18			
		32	438	21			
		7	439	24			
12	455-488	33	455	21	0.235	0.121	1.946
		47	459	20			
		56	462	19			
		87	463	17			
		88	466	17			
		14	470	23			
		109	473	16			
		34	480	21			
13	515-530	57	515	19	0.125	0.121	1.034
		22	522	22			
14	557-590	8	557	24	0.147	0.121	1.216
		23	564	22			
		9	571	24			
		90	575	17			
		58	582	19			
15	610-625	69	610	18	0.125	0.121	1.034
		91	617	17			
16	633-668	92	633	17	0.222		
		10	635	24			
		70	638	18			
		93	640	17			
		48	642	20			
		49	645	20			
		111	652	16			
		112	660	16			
17	674-685	71	674	18	0.167	0.121	1.379
		11	677	24			
18	687-702	1	687	26	0.125	0.121	1.034
		94	694	17			
19	744-767	113	744	16	0.250	0.121	2.068
		95	745	17			
		4	745	25			
		24	752	22			
		2	755	26			
		72	759	18			
20	812-827	97	812	17	0.125	0.121	1.034
		115	819	16			
21	838-857	116	838	16	0.150	0.121	1.241
		25	846	22			
		74	849	18			

22	896-913	117	896	16	0.222	0.121	1.838
		98	899	17			
		26	902	22			
		76	905	18			

1 MLLAVLYCLL WSFQTSAGHF PRACVSSKNL MEKECCPPWS GDRSPCGQLS
 GRGSCQNILL
 5 61 SNAPLGPQFP FTGVDDRESW PSVFYNRTCQ CSGNFMGFNC GNCKFGFWGP
 NCTERRLLVR
 121 RNIFDLSAPE KDKFFAYLTL AKHTISSDYV IPIGTYGQMK NGSTPMFNDI
 NIYDLFVWMH
 181 YYVSMDALLG GSEIWRDIDF AHEAPAFLPW HRLFLLRWEQ EIQLTGDEN
 10 FTIPYWDWRD
 241 AEKCDICTDE YMGGQHPTNP NLLSPASFFS SWQIVCSRLE EYNHQSLCN
 GTPEGPLRRN
 301 PGNHDKSRTP RLPSSADVEF CLSLTQYESG SMDKAANFSF RNTLEGFASP
 LTGIADASQS
 15 361 SMHNALHIYM NGTMSQVQGS ANDPIFLLHH AFVDSIFEQW LRRHRPLQEV
 YPEANAPIGH
 421 NRESYMPFI PLYRNGDFFI SSKDLGYDYS YLQSDPDSF QDYIKSYLEQ
 ASRIWSWLLG
 481 AAMVGAVLTA LLAGLVSLLC RHKRKQLPEE KQPLLMEKED YHSLYQSHL
 20 TYROSINASE PROTEIN

25

30 1 MNGDDAFARR PTVGAQIPEK IQKAFDDIAK YFSKEEWEKM KASEKIFYVY
 MKRKYEAMTK
 61 LGFKATLPPF MCNKRAEDFQ GNDLDNDPNR GNQVERPQMT FGRLQGISPK
 IMPKKPAEEG
 121 NDSEEVPEAS GPQNDGKELC PPGKPTTSEK IHERSGPKRG EHAWTHRLRE
 35 RKQLVIYEEI
 181 SDPEEDE

SSX-2 PROTEIN

40

1 MWNLLHETDS AVATARRPRW LCAGALVLAG GFFLLGFLFG WFIKSSNEAT
 NITPKHNMKA
 61 FLDELKAENI KKFLYNFTQI PHLAGTEQNF QLAKQIQSQW KEFGLDSEVL
 5 AHYDVLLSYP
 121 NKTHPNYISI INEDGNEIFN TSLFEPPLPG YENVSDIVPP FSAFSPQGMF
 EGDLVVYNYA
 181 RTEDFFKLER DMKINCSGKI VIARYGKVFR GNKVKNQALA GAKGVILYSD
 PADYFAPGVK
 10 241 SYPDGWNLPG GGVQRGNIIN LNGAGDPLTP GYPANEYAYR RGIAEAVGLP
 SIPVHPIGYI
 301 DAQKLEKMG GSAPPDSSWR GSKLVFYNVG PGFTGNFSTQ KVKMHIHSTN
 EVTRIYNVIG
 361 TLRGAVEPDR YVILGGHRDS WVFGGIDPOS GAAVVHEIVR SFGTLKKEGW
 15 RPRRTILFAS
 421 WDAEEFGLLG STEWAEENSR LLQERGVAI NADSSIEGNY TLRVDCTPLM
 YSLVHNLTKE
 481 LKSPDEGFEG KSLYESWTKK SPSPEFSGMP RISKLGSGND FEVFFQRLGI
 ASGRARYTKN
 20 541 WETNKFSGYP LYHSVYETYE LVEKFYDPMF KYHLTVAQVR GGMVFELANS
 IVLPFDCRDY
 601 AVVLRKYADK IYSISMKHPQ EMKTYSVSFD SLFSAVKNFT EIASKFSERL
 QDFDKSNPIV
 661 LRMMNDQLMF LERAFIDPLG LPDRPFYRHV IYAPSSHNKY AGESFPGIYD
 25 ALFDIESKVD
 721 PSKAWGEVKK QIYVAAFTVQ AAAETLSEVA

PSMA PROTEIN

30

1: Homo sapiens tyrosinase PubMed, Protei
 NM_00037 (oculocutaneous albinism IA) (TYR), Taxonomy, OMIM, LinkO
 2 Mrna
 LOCUS NM_000372 1964 bp mRNA PRI 31-
 OCT-2000
 DEFINITION Homo sapiens tyrosinase (oculocutaneous albinism IA)
 35 (TYR), mRNA.
 ACCESSION NM_000372
 VERSION NM_000372.1 GI:4507752
 KEYWORDS .
 SOURCE human.
 40 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
 Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
 Homo.
 45 REFERENCE 1 (bases 1 to 1964)
 AUTHORS Kwon BS, Haq AK, Pomerantz SH and Halaban R.
 TITLE Isolation and sequence of a cDNA clone for human
 tyrosinase that
 maps at the mouse c-albino locus
 50 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 84 (21), 7473-7477
 (1987)
 MEDLINE 88041128
 PUBMED 2823263
 REMARK Erratum: [[published erratum appears in Proc Natl Acad
 55 Sci U S A

- 1988 Sep;85(17):6352]]
2 (bases 1 to 1964)
AUTHORS Barton DE, Kwon BS and Francke U.
TITLE Human tyrosinase gene, mapped to chromosome 11 (q14---
5 -q21),
defines second region of homology with mouse
chromosome 7
JOURNAL Genomics 3 (1), 17-24 (1988)
MEDLINE 89122007
10 PUBMED 3146546
REFERENCE 3 (bases 181 to 1964)
AUTHORS Shibahara,S., Tomita,Y., Tagami,H., Muller,R.M. and
Cohen,T.
TITLE Molecular basis for the heterogeneity of human
15 tyrosinase
JOURNAL Tohoku J. Exp. Med. 156 (4), 403-414 (1988)
MEDLINE 89222868
REFERENCE 4 (bases 1 to 1964)
20 AUTHORS Bouchard B, Fuller BB, Vijayasaradhi S and Houghton
AN.
TITLE Induction of pigmentation in mouse fibroblasts by
expression of
human tyrosinase cDNA
JOURNAL J. Exp. Med. 169 (6), 2029-2042 (1989)
25 MEDLINE 89279151
PUBMED 2499655
REFERENCE 5 (bases 1 to 1964)
AUTHORS Takeda,A., Tomita,Y., Okinaga,S., Tagami,H. and
Shibahara,S.
30 TITLE Functional analysis of the cDNA encoding human
tyrosinase precursor
JOURNAL Biochem. Biophys. Res. Commun. 162 (3), 984-990 (1989)
MEDLINE 89351001
REFERENCE 6 (bases 1 to 1964)
35 AUTHORS Kikuchi H, Miura H, Yamamoto H, Takeuchi T, Dei T and
Watanabe M.
TITLE Characteristic sequences in the upstream region of the
human
tyrosinase gene
40 JOURNAL Biochim. Biophys. Acta 1009 (3), 283-286 (1989)
MEDLINE 90089403
PUBMED 2480811
REFERENCE 7 (bases 1 to 1964)
AUTHORS Giebel LB, Strunk KM and Spritz RA.
45 TITLE Organization and nucleotide sequences of the human
tyrosinase gene
and a truncated tyrosinase-related segment
JOURNAL Genomics 9 (3), 435-445 (1991)
50 MEDLINE 91236163
PUBMED 1903356
REFERENCE 8 (bases 1 to 1964)
AUTHORS Brichard V, Van Pel A, Wolfel T, Wolfel C, De Plaen B,
Lethe B,
Coulie P and Boon T.
55 TITLE The tyrosinase gene codes for an antigen recognized by
autologous
cytolytic T lymphocytes on HLA-A2 melanomas
JOURNAL J. Exp. Med. 178 (2), 489-495 (1993)

MEDLINE 93340625
PUBMED 8340755
COMMENT PROVISIONAL REFSEQ: This record has not yet been
subject to final NCBI review. The reference sequence was derived from
M27160.1.
5 FEATURES Location/Qualifiers
source 1..1964
/organism="Homo sapiens"
10 /db_xref="taxon:9606"
/chromosome="11"
/map="11q14-q21"
/cell_line="MeWo melanoma DNA, and cDNA to
mRNA"
15 /tissue_type="placenta"
gene 1..1964
/gene="TYR"
/note="OCAIA"
/db_xref="LocusID:7299"
20 /db_xref="MIM:203100"
CDS 83..1672
/gene="TYR"
/EC_number="1.14.18.1"
25 /note="Tyrosinase"
/codon_start=1
/db_xref="LocusID:7299"
/db_xref="MIM:203100"
/product="tyrosinase (oculocutaneous albinism
IA)"
30 /protein_id="NP_000363.1"
/db_xref="GI:4507753"
/translation="MLLAVLYCLWSFQTSAGHFPRACVSSKNLMEKECCPPWSGDRS
35 PCGQLSGRGSCQNILLSNAPLGPQFPFTGVDDRESWPSVFYNRTCQCSGNFMGFNCGN
CKFGFWGPNCTERRLLVRRNIFDLSAPEKDKFFAYLTLAKHTISSDYVIPIGTYGQMK
NGSTPMFNDINIYDLFVWMHYYSMDALLGGSEIWRDIDFAHEAPAFLPWHRLFLLRW
40 EQEIQKLTGDNFTIPYWDWRDAEKCDICTDEYMGGQHPTNPPLLSPASFFSSWQIVC
SRLEEYNShQSLCNGTPEGPLRRNPGNHDKSRTPLPSSADVEFCLSLTQYESGSMDK
AANFSFRNTLEGFASPLTGIADASQSSMHNALHIYMNGTMSQVQGSANDPIFLLHHAF
45 VDSIFEQWLRHRPLQEVYPEANAPIGHNRESYMVPFIPLYRNGDFFISSKDLGYDYS
YLQSDPDSFQDYIKSYLEQASRIWSWLLGAAMVGAVLTALLAGLVSLLCRHKRKLPL
50 EEKQPLLMEKEDYHSLYQSHL"
sig peptide 83..136
mat peptide 137..1669
/note="tyrosinase"
/product="tyrosinase (oculocutaneous albinism
55 IA)"
misc feature 560..1429
/note="tyrosinase; Region: Common central
domain of

tyrosinase"

	BASE COUNT	520 a	462 c	445 g	537 t	
	ORIGIN					
		1	atcactgtag	tagtagctgg	aaagagaaat	ctgtgactcc aattagccag
5			ttcctgcaga			
		61	ccttgtgagg	actagaggaa	gaatgctcct	ggctgttttg tactgcctgc
			tgtggagttt			
		121	ccagacctcc	gctggccatt	tccttagagc	ctgtgtctcc tctaagaacc
			tgatggagaa			
10		181	ggaatgctgt	ccaccgtgga	gcggggacag	gagtccttgt ggccagcttt
			caggcagagg			
		241	ttcctgtcag	aatatccttc	tgtccaatgc	accacttggg cctcaatttc
			ccttcacagg			
		301	ggtggatgac	cgggagtcgt	ggccttcctg	ottttataat aggacctgcc
15			agtgtcttgg			
		361	caacttcatg	ggattcaact	gtggaaactg	caagtttggc ttttggggac
			caaactgcac			
		421	agagagacga	ctcttgggtg	gaagaaacat	cttcgatttg agtgccccag
			agaaggacaa			
20		481	attttttggc	tacctcactt	tagcaaagca	taccatcagc tcagactatg
			tcaccccat			
		541	agggacctat	ggccaaatga	aaaatggatc	aacacccatg ttttaacgaca
			tcaatattta			
		601	tgacctcttt	gtctggatgc	attattatgt	gtcaatggat gcaactgcttg
25			ggggatctga			
		661	aatctggaga	gacattgatt	ttgcccatga	agcaccagct tttctgcctt
			ggcatagact			
		721	cttcttgttg	cggtgggaac	aagaaatcca	gaagctgaca ggagatgaaa
			acttcactat			
30		781	tcacatattg	gactggcggg	atgcagaaaa	gtgtgacatt tgcacagatg
			agtacatggg			
		841	aggtcagcac	cccacaaatc	ctaacttact	cagcccagca tcattcttct
			cctcttggca			
		901	gattgtctgt	agccgatttg	aggagtacaa	cagccatcag tctttatgca
35			atggaacgcc			
		961	cgagggacct	ttacggcgta	atcctggaaa	ccatgacaaa tccagaaccc
			caaggctccc			
		1021	ctcttcagct	gatgtagaat	tttgccctgag	tttgacccaa tatgaatctg
			gttccatgga			
40		1081	taaagctgcc	aatttcagct	ttagaaatac	actggaagga tttgctagtc
			cacttactgg			
		1141	gatagcggat	gcctctcaaa	gcagcatgca	caatgccttg cacatctata
			tgaatggaac			
		1201	aatgtcccag	gtacagggat	ctgccaacga	tcctatcttc cttcttcacc
45			atgcatttgt			
		1261	tgacagtatt	tttgagcagt	ggctccgaag	gcaccgtcct cttcaagaag
			tttatccaga			
		1321	agccaatgca	cccattggac	ataaccggga	atcctacatg gttcctttta
			taccactgta			
50		1381	cagaaatggt	gatttcttta	tttcatccaa	agatctgggc tatgactata
			gctatctaca			
		1441	agattcagac	ccagactctt	ttcaagacta	cattaagtcc tatttggaac
			aagcgagtgc			
		1501	gatctggtca	tggctccttg	gggcccgcgat	ggtagggggc gtccctcactg
55			ccctgctggc			
		1561	agggcttgtg	agcttgctgt	gtcgtcacaa	gagaaagcag cttcctgaag
			aaaagcagcc			

1621 actcctcatg gagaaagagg attaccacag cttgtatcag agccatttat
 aaaaggctta
 1681 ggcaatagag tagggccaaa aagcctgacc tcaactaac tcaaagtaat
 gtccaggttc
 5 1741 ccagagaata tctgctggta tttttctgta aagaccattt gcaaaattgt
 aacctaatac
 1801 aaagtgtagc ctctctccaa ctcaggtaga acacacctgt ctttgtcttg
 ctgttttcac
 1861 tcagcccttt taacattttc ccctaagccc atatgtctaa ggaaaggatg
 10 ctatttggtg
 1921 atgaggaact gttatttgta tgtgaattaa agtgctctta tttt

	1:	Homo sapiens synovial sarcoma, NM_003147 X breakpoint 2 (SSX2), mRNA	PubMed, Protein, Related Sequences, Taxonomy, OMIM LinkOut
15	LOCUS	NM_003147 766 bp mRNA	PRI 14-
	MAR-2001		
	DEFINITION	Homo sapiens synovial sarcoma, X breakpoint 2 (SSX2), mRNA.	
20	ACCESSION	NM_003147	
	VERSION	NM_003147.1 GI:10337582	
	KEYWORDS	.	
	SOURCE	human.	
	ORGANISM	Homo sapiens	
25		Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;	
	Euteleostomi;		
		Mammalia; Eutheria; Primates; Catarrhini; Hominidae;	
	Homo.		
	REFERENCE	1 (bases 1 to 766)	
30	AUTHORS	Shipley JM, Clark J, Crew AJ, Birdsall S, Rocques PJ, Gill S,	
		Chelly J, Monaco AP, Abe S, Gusterson BA and et al.	
	TITLE	The t(X;18)(p11.2;q11.2) translocation found in human	
	synovial		
35	chromosome	sarcomas involves two distinct loci on the X	
	JOURNAL	Oncogene 9 (5), 1447-1453 (1994)	
	MEDLINE	94203675	
	PUBMED	8152806	
	REFERENCE	2 (bases 1 to 766)	
40	AUTHORS	Crew,A.J., Clark,J., Fisher,C., Gill,S., Grimer,R., Chand,A.,	
		Shipley,J., Gusterson,B.A. and Cooper,C.S.	
	TITLE	Fusion of SYT to two genes, SSX1 and SSX2, encoding	
45	proteins with	homology to the Kruppel-associated box in human	
	synovial sarcoma		
	JOURNAL	EMBO J. 14 (10), 2333-2340 (1995)	
	MEDLINE	95292974	
	REFERENCE	3 (bases 1 to 766)	
50	AUTHORS	Tureci O, Sahin U, Schobert I, Koslowski M, Schmitt H, Schild HJ,	
		Stenner F, Seitz G, Rammensee HG and Pfreundschuh M.	
	TITLE	The SSX-2 gene, which is involved in the t(X;18)	
	translocation of		

synovial sarcomas, codes for the human tumor antigen

HOM-MEL-40
 JOURNAL Cancer Res. 56 (20), 4766-4772 (1996)
 MEDLINE 96438636
 PUBMED 8840996

5 COMMENT PROVISIONAL REFSEQ: This record has not yet been
 subject to final
 NCBI review. The reference sequence was derived from
 X86175.1.

10 FEATURES Location/Qualifiers
 source 1..766
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /chromosome="X"
 /map="Xp11.23-p11.22"
 /dev_stage="adult"
 gene 1..766
 /gene="SSX2"
 /note="HD21; HOM-MEL-40; SSX"
 /db_xref="LocusID:6757"
 /db_xref="MIM:300192"
 misc feature 20..61
 /note="Kruppel associated box homology"
 CDS 92..658
 /gene="SSX2"
 /codon_start=1
 /db_xref="LocusID:6757"
 /db_xref="MIM:300192"
 /product="synovial sarcoma, X breakpoint 2"
 /protein_id="NP_003138.1"
 /db_xref="GI:10337583"

25

30 /translation="MNGDDAFARRPTVGAQIPEKIQKAFDDIAKYFSKEWEKMKASE
 35 KIFYVYMKRKYEAMTKLGFKATLPPFMCNKRAEDFQGNLDNDPNRGNQVERPQMTFG
 RLQGISPKIMPKPAEEGNDSEEVPEASGPQNDGKELCPPGKPTTSEKIHERSGPKRG
 EHAWTHRLRERKQLVIYEEISDPPEEDDE"
 misc feature 161..337
 /note="KRAB; Region: krueppel associated box"

40 BASE COUNT 229 a 181 c 200 g 156 t
 ORIGIN
 1 ctctctttcg attcttccat actcagagta cgcaagggtct gattttctct
 ttggattctt
 45 61 ccaaaatcag agtcagactg ctcccgggtgc catgaacgga gacgacgcct
 ttgcaaggag
 121 acccacggtt ggtgctcaaa taccagagaa gatccaaaag gccttcgatg
 atattgccaa
 181 atacttctct aaggaagagt gggaaaagat gaaagcctcg gagaaaatct
 50 tctatgtgta
 241 tatgaagaga aagtatgagg ctatgactaa actaggtttc aaggccaccc
 tcccaccttt
 301 catgtgtaat aaacgggccc aagacttcca ggggaatgat ttggataatg
 accctaaccg
 55 361 tgggaatcag gttgaacgtc ctcatgatgac ttctggcagg ctccagggaa
 tctcccggaa
 421 gatcatgccc aagaagccag cagaggaagg aaatgattcg gaggaagtgc
 cagaagcatc

481 tggcccacaa aatgatggga aagagctgtg ccccccggga aaaccaacta
 cctctgagaa
 541 gattcacgag agatctggac ccaaaagggg ggaacatgcc tggaccaca
 gactgcgtga
 5 601 gagaaaacag ctggtgattt atgaagagat cagcgaccct gaggaagatg
 acgagtaact
 661 cccctcaggg atacgacaca tgcccatgat gagaagcaga acgtggtgac
 ctttcacgaa
 721 catgggcatg gctgcggacc cctcgtcac aggtgcatag caagtg
 10

1: **Homo sapiens folate hydrolase** PubMed, Protein, Related
NM_004 (prostate-specific membrane antigen) Sequences, Taxonomy, OMIM
476 1 (FOLH1), mRNA LinkOut
 LOCUS NM_004476 2653 bp mRNA PRI 01-
 NOV-2000
 15 DEFINITION Homo sapiens folate hydrolase (prostate-specific
 membrane antigen)
 1 (FOLH1), mRNA.
 ACCESSION NM_004476
 VERSION NM_004476.1 GI:4758397
 20 KEYWORDS .
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
 Euteleostomi;
 25 Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
 Homo.
 REFERENCE 1 (bases 1 to 2653)
 AUTHORS Israeli,R.S., Powell,C.T., Fair,W.R. and Heston,W.D.
 TITLE Molecular cloning of a complementary DNA encoding a
 30 prostate-specific membrane antigen
 JOURNAL Cancer Res. 53 (2), 227-230 (1993)
 MEDLINE 93113576
 REFERENCE 2 (bases 1 to 2653)
 35 AUTHORS Rinker-Schaeffer CW, Hawkins AL, Su SL, Israeli RS,
 Griffin CA,
 Isaacs JT and Heston WD.
 TITLE Localization and physical mapping of the prostate-
 specific membrane
 40 antigen (PSM) gene to human chromosome 11
 JOURNAL Genomics 30 (1), 105-108 (1995)
 MEDLINE 96129312
 PUBMED 8595888
 REFERENCE 3 (bases 1 to 2653)
 45 AUTHORS O'Keefe DS, Su SL, Bacich DJ, Horiguchi Y, Luo Y,
 Powell CT,
 Zandvliet D, Russell PJ, Molloy PL, Nowak NJ, Shows
 TB, Mullins C,
 Vonder Haar RA, Fair WR and Heston WD.
 TITLE Mapping, genomic organization and promoter analysis of
 50 the human
 prostate-specific membrane antigen gene
 JOURNAL Biochim. Biophys. Acta 1443 (1-2), 113-127 (1998)
 MEDLINE 99057588
 55 PUBMED 9838072
 REFERENCE 4 (bases 1 to 2653)

AUTHORS Maraj BH, Leek JP, Karayi M, Ali M, Lench NJ and
 Markham AF.
 TITLE Detailed genetic mapping around a putative prostate-
 specific
 5 membrane antigen locus on human chromosome 11p11.2
 JOURNAL Cytogenet. Cell Genet. 81 (1), 3-9 (1998)
 MEDLINE 98358137
 PUBMED 9691167
 COMMENT PROVISIONAL REFSEQ: This record has not yet been
 0 subject to final
 NCBI review. The reference sequence was derived from
 M99487.1.
 FEATURES Location/Qualifiers
 .5 source 1..2653
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /chromosome="11"
 /map="11p11.2"
 /sex="male"
 20 /cell_line="LNCaP-ATCC"
 /cell_type="prostate"
 /tissue_type="prostatic carcinoma metastatic
 lymph node"
 25 gene 1..2653
 /gene="FOLH1"
 /note="FOLH; PSM; PSMA"
 /db_xref="LocusID:2346"
 /db_xref="MIM:600934"
 30 CDS 262..2514
 /gene="FOLH1"
 /note="folate hydrolase 1 (prostate-specific
 membrane
 35 antigen)"
 /codon_start=1
 /db_xref="LocusID:2346"
 /db_xref="MIM:600934"
 /evidence=experimental
 /product="folate hydrolase (prostate-specific
 40 membrane
 antigen) 1"
 /protein_id="NP_004467.1"
 /db_xref="GI:4758398"
 45 /translation="MWNLLHETDSAVATARRPRWLCAGALVLGGFFLLGFLFGWFIK
 SSNEATNITPKHNMKAFLDELKAENIKKFLYNFTQIPHLAGTEQNFQLAKQIQSQWKE
 FGLDSVELAHYDVLLSYPNKTHPNYISIIINEDGNEIFNTSLFEPPPPGYENVSDIVPP
 50 FSAFSPQGMPEGDLVYVNYARTEDFFKLERDMKINCSGKIVARIYGVFRGNKVKNAQ
 LAGAKGVILYSDPADYFAPGVKSYPDGWNLPGGGVQQRGNILNLNGAGDPLTPGYANE
 55 YAYRRGIAEAVGLPSIPVHPIGYYDAQKLEKMGGSAPPDSSWRGSLKVPYNVGPGET
 GNFTSQKVKMHIHSTNEVTRIYNVIGTLRGAVEPDRYVILGGHRDSWVFGGIDPQSGA

AVVHEIVRSFGTLKKEGWRPRRTILFASWDAEEFGLLGSTEWAEENSRLQERGVAYI
 NADSSIEGNYTLRVDCTPLMYSLVHNLTKEKSPDEGFEGKSLYESWTKKSPSPFSG
 5 MPRISKLGSGNDFEVFFQRLGIASGRARYTKNWETNKFSGYPLYHSVYETVELVEKFY
 DPMFKYHLTVAQVRGGMVFELANSIVLPFDCRDYAVVLRKYADKIYSISMKHPQEMKT
 10 YSVSFDLSLFAVKNFTEIASKFSERLQDFDKSNPIVLRMMNDQLMFLERAFIDPLGLP
 DRPFYRHVIYAPSSSHNKYAGESFPGIYDALFDIESKVDPSKAWGEVVKRQIYVAAFTVQ
 AAAETLSEVA"
 misc feature 778..1029
 15 /note="PA; Region: PA domain"
 BASE COUNT 782 a 524 c 640 g 707 t
 ORIGIN
 1 ctcaaaagg ggcggatttc cttctcctgg aggcagatgt tgctctcttc
 tctcgctcgg
 20 61 attggttcag tgcactctag aaacactgct gtggtggaga aactggaccc
 caggctctgga
 121 gcgaattcca gcctgcaggg ctgataagcg aggcattagt gagattgaga
 gagactttac
 181 cccgccgtgg tgggttgagg gcgcgcagta gagcagcagc acaggcgagg
 25 gtcccgaggag
 241 gccggctctg ctgcgcgcga gatgtggaat ctcccttcacg aaaccgactc
 ggctgtggcc
 301 accgcgcgcc gccgcgctg gctgtgcgct ggggcgctgg tgctggcggg
 tggcttcttt
 361 ctccctcggct tcctcttcgg gtggtttata aaatcctcca atgaagctac
 30 taacattact
 421 ccaaagcata atatgaaagc atttttggat gaattgaaag ctgagaacat
 caagaagttc
 481 ttatataatt ttacacagat accacattta gcaggaacag aacaaaactt
 35 tcagcttgca
 541 aagcaaattc aatcccagtg gaaagaattt ggcttgatt ctgttgagct
 agcacattat
 601 gatgtcctgt tgcctaccc aaataagact catcccaact acatctcaat
 aattaatgaa
 40 661 gatggaaatg agattttcaa cacatcatta tttgaaccac ctccctcagg
 atatgaaaat
 721 gtttcggata ttgtaccacc tttcagtgtt ttctctctc aaggaatgcc
 agagggcgat
 781 ctagtgatat ttaactatgc acgaactgaa gacttcttta aattggaacg
 45 ggacatgaaa
 841 atcaattgct ctgggaaaat tgtaattgcc agatatggga aagttttcag
 aggaaataag
 901 gttaaaaatg ccagctggc aggggccaaa ggagtcattc tctactccga
 ccctgctgac
 50 961 tactttgctc ctggggtgaa gtccatocca gatggttgga atcttcctgg
 aggtggtgtc
 1021 cagcgtggaa ataccctaaa tctgaatggt gcaggagacc ctctcacacc
 aggttaccca
 1081 gcaaatgaat atgcttatag gcgtggaatt gcagaggctg ttggtcttcc
 55 aagtattcct
 1141 gttcatocaa ttggatacta tgatgcacag aagctcctag aaaaaatggg
 tggctcagca

1201 ccaccagata gcagctggag aggaagtctc aaagtgccct acaatgttgg
acctggcttt
1261 actggaaact tttctacaca aaaagtcaag atgcacatcc actctacca
tgaagtgaca
5 1321 agaatttaca atgtgatagg tactctcaga ggagcagtgg aaccagacag
atatgtcatt
1381 ctgggagggtc accgggactc atgggtgttt ggtggtattg accctcagag
tggagcagct
1441 gttgttcatg aaattgtgag gagctttgga aactgaaaa aggaaggggtg
10 gagacctaga
1501 agaacaattt tgtttgcaag ctgggatgca gaagaatttg gtcttcttgg
ttctactgag
1561 tgggcagagg agaattcaag actccttcaa gagcgtggcg tggcttatat
taatgctgac
15 1621 tcatctatag aaggaaacta cactctgaga gttgattgta caccgctgat
gtacagcttg
1681 gtacacaacc taacaaaaga gctgaaaagc cctgatgaag gctttgaagg
caaatctctt
1741 tatgaaagtt ggactaaaaa aagtccttcc ccagagttca gtggcatgcc
20 caggataagc
1801 aaattgggat ctggaaatga ttttgagggtg ttcttccaac gacttggaa
tgcttcaggc
1861 agagcacggt atactaaaaa ttgggaaaca aacaaattca gcggctatcc
actgtatcac
25 1921 agtgtctatg aaacatatga gttggtggaa aagttttatg atccaatgtt
taaataatcac
1981 ctactgtgg cccaggttcg aggagggatg gtgtttgagc tagccaattc
catagtgtc
2041 ctttttgatt gtcgagatta tgctgtagtt ttaagaaagt atgctgacaa
30 aatctacagt
2101 atttctatga aacatccaca ggaaatgaag acatacagtg tatcatttga
ttcaactttt
2161 tctgcagtaa agaattttac agaaattgct tccaagttca gtgagagact
ccaggacttt
35 2221 gacaaaagca acccaatagt attaagaatg atgaatgatc aactcatggt
tctggaaaga
2281 gcatttattg atccattagg gttaccagac aggccttttt ataggcatgt
catctatgct
2341 ccaagcagcc acaacaagta tgcaggggag tcattcccag gaatttatga
40 tgctctgttt
2401 gatattgaaa gcaaagtgga cccttccaag gcctggggag aagtgaagag
acagatttat
2461 gttgcagcct tcacagtgca ggcagctgca gagactttga gtgaagtagc
ctaagaggat
45 2521 tcttttagaga atccgtattg aatttgtgtg gtatgtcact cagaaagaat
cgtaatgggt
2581 atattgataa attttaaaat tggatatattt gaaataaagt tgaatattat
atataaaaaa
2641 aaaaaaaaaa aaa

U20093. Human melanocyte-...[gi:1142634]
LOCUS HSPMEL17S2 2817 bp DNA PRI 20-
NOV-1996
DEFINITION Human melanocyte-specific (pmel 17) gene, exons 2-5,
5 and complete
cds.
ACCESSION U20093
VERSION U20093.1 GI:1142634
KEYWORDS .
10 SEGMENT 2 of 2
SOURCE human.
ORGANISM Homo sapiens
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
15 Euteleostomi;
Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
Homo.
REFERENCE 1 (bases 1 to 2817)
AUTHORS Kim,K.K., Youn,B.S., Heng,H.H., Shi,X.M., Tsui,L.C.,
Lee,Z.H.,
20 Pickard,R.T. and Kwon,B.S.
TITLE Genomic organization and FISH mapping of human Pmel
17, the
putative silver locus
JOURNAL Pigment Cell Res. 9 (1), 42-48 (1996)
25 MEDLINE 96314705
REFERENCE 2 (bases 1 to 2817)
AUTHORS Kwon,B.S.
TITLE Direct Submission
JOURNAL Submitted (05-JAN-1995) Indiana University School of
30 Medicine,
Microbiology and Immunology, 635 Barnhill Drive,
Indianapolis, IN
46202, USA
FEATURES Location/Qualifiers
35 source 1..2817
/organism="Homo sapiens"
/db_xref="taxon:9606"
/chromosome="12"
40 /map="12pter-q21"
/clone_lib="lambda FIX II from Stratagene"
gene join(U19491.1:1111..3109,1..2722)
/gene="pmel 17"
CDS
45 join(U19491.1:1111..1186,67..459,853..1014,1266..2396,
2499..2722)
/gene="pmel 17"
/codon_start=1
/product="Pmel 17"
50 /protein_id="AAB19181.1"
/db_xref="GI:1142636"

/translation="MDLVLRCLLHLAVIGALLAVGATKVP RNQDWLGVS RQLRTKAWN RQLYPEW
TEAQR LDCWRGGQVSLKVSNDGPTLIGANASFSIALNFP GSQKVL PDGQVIWVNNTI INGSQVWGG
QPVP PQETDDACIFPDGGPCPSGWSQKR SFVYVWKTWGQY WQVLGGPVSGLSIGTGRAMLGTHM
55 EVTVYHRRGSR SYVPLAHSSSAFTITDQVPFSVSVSQLRALDGGNKHFLRNQPLTFALQLHDP SGY
LAEADLSYTWDFGDSSGTLISRAPVVTHTYLEPGPVTAQVVLQAAIPLTSCGSSPVP GTTDGHRPT
AEAPNTTAGQVPTTEVVGTT PGQAPTAEP SGTTTSVQVPTTEVISTAPVQMPTAESTGMTPEKVPVS
EVMGTTLAEMSTPEATGMTPAEVSIVVLSGTTAAQVTTTEWVETTARELP IPEPEGPDASSIMSTE

SITGSLGPLLDGTATLRLVKRQVPLDCVLYRYGSFSVTLDIVQGIESAEILQAVPSGEGDAFELTV
 SCQGGLPKACMEISSPGCQPPAQRLCQPVLPSPACQLVLHQILKGGSGTYCLNVSADTNSLAVV
 STQLIMPGQEAGLGQVPLIVGILLVLMVVLASLIYRRRLMKQDFSVLPQLPHSSSHWLRLPRIFCS
 CPIGENSPLLSGQQV"

5	exon	67..459				
		/gene="pmel 17"				
		/number=2				
	exon	853..1014				
10		/gene="pmel 17"				
		/number=3				
	exon	1266..2396				
		/gene="pmel 17"				
		/number=4				
	exon	2499..>2722				
15		/gene="pmel 17"				
		/number=5				
	BASE COUNT	627 a	772 c	739 g	679 t	-
	ORIGIN					
	1	gtgctaaaaa	gatgccttct	tcatttggtc	gtgataggtg	ctttgtggct
20	gtgggggcta					
	61	caaaagtacc	cagaaaccag	gactggcttg	gtgtctcaag	gcaactcaga
	accaaagcct					
	121	ggaacaggca	gctgtatcca	gagtggacag	aagcccagag	acttgactgc
	tggagagggtg					
25	181	gtcaagtgtc	cctcaaggtc	agtaatgatg	ggcctacact	gattgggtgca
	aatgcctcct					
	241	tctctattgc	cttgaacttc	cctggaagcc	aaaagggtatt	gccagatggg
	caggttatct					
	301	gggtcaacaa	taccatcatc	aatgggagcc	aggtgtgggg	aggacagcca
30	gtgtatcccc					
	361	aggaaactga	cgatgcctgc	atcttccttg	atggtggacc	ttgcccattc
	ggctcttggg					
	421	ctcagaagag	aagctttggt	tatgtctgga	agacctgggg	tgagggactc
	ccttctcagc					
35	481	ctatcatcca	cacttgtggt	tacttctttc	tacctgatca	cctttctttt
	ggccgccccct					
	541	tccaccttaa	cttctgtgat	tttctctaat	cttcattttc	ctcttagatc
	ttttctcttt					
	601	cttagcacct	agcccccttc	aagctctatc	ataattcttt	ctggcaactc
40	ttggcctcaa					
	661	ttgtagtcc	accccatgga	atgcctcatt	aggacctctt	ccctgtcccc
	ccatatcaca					
	721	gccttccaaa	caccctcaga	agtaatcata	cttcctgacc	tcccatctcc
	agtgcggttt					
45	781	cgaagcctgt	ccctcagtc	cctttgacca	gtaatctctt	cttccttgct
	tttcattcca					
	841	aaaatgcttc	aggccaatac	tggcaagttc	tagggggccc	agtgtctggg
	ctgagcattg					
	901	ggacaggcag	ggcaatgctg	ggcacacaca	ccatggaagt	gactgtctac
50	catcgccggg					
	961	gatcccggag	ctatgtgcct	cttgtctcatt	ccagctcagc	cttcaccatt
	actggtaagg					
	1021	gttcaggaag	ggcaaggcca	gttgtagggc	aaagagaagg	cagggaggct
	tggatggact					
55	1081	gcaaaggaga	aagggtgaaat	gctgtgcaaa	cttaaagtag	aagggccagg
	aagacctagg					
	1141	cagagaaatg	tgaggcttag	tgccagtgaa	gggccagcca	gtcagcttgg
	agttggaggg					

	1201	tgtggctgtg	aaaggagaag	ctgtggctca	ggcctggttc	tcaccttttc
		tggctccaat				
	1261	cccagaccag	gtgcctttct	ccgtgagcgt	gtcccagttg	cgggccttgg
		atggagggaa				
5	1321	caagcacttc	ctgagaaatc	agcctctgac	ctttgccctc	cagctccatg
		accccagtgg				
	1381	ctatctggct	gaagctgacc	tctcctacac	ctgggacttt	ggagacagta
		gtggaaccct				
10	1441	gatctctcgg	gcacctgtgg	tcactcatac	ttacctggag	cctggcccag
		tcactgcccc				
	1501	ggtggtcctg	caggctgcca	ttcctctcac	ctcctgtggc	tcctccccag
		ttccaggcac				
	1561	cacagatggg	cacaggccaa	ctgcagaggc	ccctaacacc	acagctggcc
		aagtgcctac				
15	1621	tacagaagtt	gtgggtacta	cacctgggtca	ggcgccaact	gcagagccct
		ctggaaccac				
	1681	atctgtgcag	gtgccaacca	ctgaagtcac	aagcactgca	cctgtgcaga
		tgccaactgc				
20	1741	agagagcaca	ggtatgacac	ctgagaagggt	gccagtttca	gaggtcatgg
		gtaccacact				
	1801	ggcagagatg	tcaactccag	aggctacagg	tatgacacct	gcagaggtat
		caattgtggg				
	1861	gctttctgga	accacagctg	cacaggtaac	aactacagag	tgggtggaga
		ccacagctag				
25	1921	agagctacct	atccctgagc	ctgaagggtcc	agatgccagc	tcaatcatgt
		ctacggaaaag				
	1981	tattacaggt	tccttgggcc	ccctgctgga	tggtagagcc	accttaaggc
		tggtagaagag				
	2041	acaagtcccc	ctggattgtg	ttctgtatcg	atatgggtcc	ttttccgtca
30		ccctggacat				
	2101	tgtccagggt	attgaaagtg	ccgagatcct	gcaggctgtg	ccgtccggtg
		agggggatgc				
	2161	atcttgagctg	actgtgtcct	gccaaggcgg	gctgcccag	gaagcctgca
		tggagatctc				
35	2221	atcgccaggg	tgccagcccc	ctgcccagcg	gctgtgccag	cctgtgctac
		ccagcccagc				
	2281	ctgccagctg	gttctgcacc	agatactgaa	gggtggctcg	gggacatact
		gcctcaatgt				
40	2341	gtctctggct	gataccaaca	gcctggcagt	ggtcagcacc	cagcttatca
		tgcctggtag				
	2401	gtccttggac	agagactaag	tgaggaggga	agtggataga	ggggacagct
		ggcaagcagc				
	2461	agacatgagt	gaagcagtgc	ctgggattct	tctcacaggt	caagaagcag
		gccttgggca				
45	2521	ggttccgctg	atcgtgggca	tcttgcctgg	gttgatggct	gtggtccttg
		catctctgat				
	2581	atataggcgc	agacttatga	agcaagactt	ctccgtaccc	cagttgccac
		atagcagcag				
50	2641	tcactggctg	cgtctacccc	gcattctctg	ctcttgctcc	attggtgaga
		atagccccct				
	2701	cctcagtggg	cagcaggtct	gagtactctc	atatgatgct	gtgattttcc
		tggagttgac				
	2761	agaaacacct	atatttcccc	cagtcttccc	tgggagacta	ctattaactg
55		aaataaa				
		//				

NM_001648. Homo sapiens kall...[gi:4502172]

LOCUS NM_001648 1466 bp mRNA PRI
31-OCT-2000
DEFINITION Homo sapiens kallikrein 3, (prostate specific antigen)
(KLK3),
5 mRNA.
ACCESSION NM_001648
VERSION NM_001648.1 GI:4502172
KEYWORDS .
SOURCE human.
10 ORGANISM Homo sapiens
Eukaryota; Metazoa; Chordata; Craniata;
Vertebrata; Euteleostomi;
Mammalia; Eutheria; Primates; Catarrhini;
Hominidae; Homo.
15 REFERENCE 1 (bases 1 to 1466)
AUTHORS Lundwall, A. and Lilja, H.
TITLE Molecular cloning of human prostate specific antigen
cDNA
JOURNAL FEBS Lett. 214 (2), 317-322 (1987)
20 MEDLINE 87190978
REFERENCE 2 (bases 1 to 1466)
AUTHORS Sutherland GR, Baker E, Hyland VJ, Callen DF, Close JA,
Tregear GW,
Evans BA and Richards RI.
25 TITLE Human prostate-specific antigen (APS) is a member of
the glandular
kallikrein gene family at 19q13
JOURNAL Cytogenet. Cell Genet. 48 (4), 205-207 (1988)
30 MEDLINE 89250658
PUBMED 2470553
REFERENCE 3 (bases 1 to 1466)
AUTHORS Riegman PH, Klaassen P, van der Korput JA, Romijn JC
and Trapman J.
TITLE Molecular cloning and characterization of novel
35 prostate antigen
cDNA's
JOURNAL Biochem. Biophys. Res. Commun. 155 (1), 181-188 (1988)
MEDLINE 88326297
PUBMED 2458104
40 REFERENCE 4 (bases 1 to 1466)
AUTHORS Schulz P, Stucka R, Feldmann H, Combriato G, Klobeck HG
and Fittler F.
TITLE Sequence of a cDNA clone encompassing the complete
mature human
45 prostate specific antigen (PSA) and an unspliced
leader sequence
JOURNAL Nucleic Acids Res. 16 (13), 6226 (1988)
MEDLINE 88289366
PUBMED 2456523
50 REFERENCE 5 (bases 1 to 1466)
AUTHORS Riegman PH, Vlietstra RJ, van der Korput JA, Romijn JC
and Trapman J.
TITLE Characterization of the prostate-specific antigen gene:
a novel
55 human kallikrein-like gene
JOURNAL Biochem. Biophys. Res. Commun. 159 (1), 95-102 (1989)
MEDLINE 89165891
PUBMED 2466464

REFERENCE 6 (bases 1 to 1466)
 AUTHORS Henttu P and Vihko P.
 TITLE cDNA coding for the entire human prostate specific
 antigen shows high homologies to the human tissue kallikrein
 genes
 JOURNAL Biochem. Biophys. Res. Commun. 160 (2), 903-910 (1989)
 MEDLINE 89246551
 PUBMED 2470373
 REFERENCE 7 (bases 1 to 1466)
 10 AUTHORS Klobeck HG, Combriato G, Schulz P, Arbusow V and
 Fittler F.
 TITLE Genomic sequence of human prostate specific antigen
 (PSA)
 15 JOURNAL Nucleic Acids Res. 17 (10), 3981 (1989)
 MEDLINE 89282407
 PUBMED 2471958
 REFERENCE 8 (bases 1 to 1466)
 20 AUTHORS Lundwall A.
 TITLE Characterization of the gene for prostate-specific
 antigen, a human glandular kallikrein
 JOURNAL Biochem. Biophys. Res. Commun. 161 (3), 1151-1159
 (1989)
 25 MEDLINE 89302090
 PUBMED 2472789
 COMMENT PROVISIONAL REFSEQ: This record has not yet been
 subject to final NCBI review. The reference sequence was derived
 30 from X05332.1.
 FEATURES Location/Qualifiers
 source 1..1466
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 35 /chromosome="19"
 /map="19q13"
 /clone="lambda HPSA-1"
 /tissue_type="prostate"
 /clone_lib="(lambda)gt11"
 40 gene 1..1466
 /gene="KLK3"
 /note="PSA; APS"
 /db_xref="LocusID:354"
 /db_xref="MIM:176820"
 45 sig_peptide 44..94
 CDS 44..829
 /gene="KLK3"
 /EC_number="3.4.21.77"
 /note="antigen, prostate specific"
 50 /codon_start=1
 /db_xref="LocusID:354"
 /db_xref="MIM:176820"
 /product="kallikrein 3, (prostate specific
 antigen)"
 55 /protein_id="NP_001639.1"
 /db_xref="GI:4502173"

```

/translation="MWVPVVFLLSVTWIGAAPLILSRIVGGWECEKHSQPWQVLVAS
RGRAVCGGVLVHPQWVLTAAHCI RNKSVILLGRHSLFHPEDTGQVFQVSHSFPHPPLYDMSLLKNRF
LRPGDDSSHDLMLLRLSEPAELTDAVKVMDLPTQEPALGTTTCYASGWSIEPPEFLT PPKLQCVDL
5 HVISNDVCAQVHPQKVTKFMLCAGRWTGGKSTCSGDSGGPLVCNGVLQGITSWGSEPCALPERPSL
YTKVVHYRKWIKDTIVANP"
    variation 91
                /allele="C"
                /allele="T"
10    variation 91
                /db_xref="dbSNP:1135765"
                /allele="C"
                /allele="T"
                /db_xref="dbSNP:11573"
15    misc_feature 95..115
                /note="propeptide (AA 1-7)"
    variation 97
                /allele="G"
                /allele="A"
20    misc_feature 113..802
                /db_xref="dbSNP:1135766"
                /note="Tryp_SPC; Region: Trypsin-like serine
protease"
    misc_feature 116..802
                /note="trypsin; Region: Trypsin"
25    mat_peptide 116..826
                /product="kallikrein 3, (prostate specific
antigen)"
    variation 280
                /allele="C"
                /allele="T"
                /db_xref="dbSNP:1058072"
    variation 280
                /allele="C"
                /allele="T"
35    variation 526
                /db_xref="dbSNP:12946"
                /allele="A"
                /allele="G"
40    variation 643
                /db_xref="dbSNP:1803130"
                /allele="A"
                /allele="G"
                /db_xref="dbSNP:1803129"
45    variation 788
                /note="WARNING: map location ambiguous"
                /allele="T"
                /allele="C"
                /db_xref="dbSNP:1802720"
50    variation 844
                /allele="T"
                /allele="C"
                /db_xref="dbSNP:1058205"
    variation 894
55    variation 906
                /allele="A"
                /allele="G"
                /db_xref="dbSNP:1058274"
    variation 906

```

		/allele="T"
		/allele="A"
		/db_xref="dbSNP:1803132"
5	variation	918
		/allele="T"
		/allele="C"
		/db_xref="dbSNP:1802722"
	variation	953
10		/allele="A"
		/allele="G"
		/db_xref="dbSNP:1802721"
	variation	1017
		/allele="T"
		/allele="C"
15		/db_xref="dbSNP:1803138"
	variation	1074
		/allele="C"
		/allele="T"
		/db_xref="dbSNP:12101"
20	variation	1107
		/allele="A"
		/allele="G"
		/db_xref="dbSNP:1803131"
	variation	1107
25		/allele="G"
		/allele="A"
		/db_xref="dbSNP:6998"
	variation	1211
		/allele="C"
30		/allele="T"
		/db_xref="dbSNP:1803136"
	variation	1229
		/allele="A"
		/allele="G"
35		/db_xref="dbSNP:1803135"
	variation	1249
		/allele="T"
		/allele="C"
		/db_xref="dbSNP:1803134"
40	variation	1273
		/allele="A"
		/allele="G"
		/db_xref="dbSNP:12040"
	variation	1326
45		/allele="T"
		/allele="C"
		/db_xref="dbSNP:1803133"
	variation	1344
		/allele="A"
50		/allele="G"
		/db_xref="dbSNP:12863"
	variation	1363
		/allele="A"
		/allele="G"
55		/db_xref="dbSNP:1803137"
	misc_feature	1445..1450
		/note="put.polyA signal"
	polyA_site	1466


```

                                /note="polyA site"
BASE COUNT      338 a      382 c      422 g      324 t
ORIGIN
5   1  agccccaagc  ttaccacctg  caccgcgaga  gctgtgtgtc  accatgtggg
   61  cttcctcacc  ctgtccgtga  cgtggattgg  tgctgcaccc  ctcactctgt
   121 gggaggctgg  gagtgcgaga  agcattccca  accctggcag  gtgcttgtgg
10  181 cagggcagtc  tgcggcggtg  ttctggtgca  ccccagtggt  gtccctcacag
   241 catcaggaac  aaaagcgtga  tcttgctggg  tcggcacage  ctgtttcatc
   301 aggccaggta  tttcagggtc  gccacagctt  cccacacccg  ctctacgata
15  361 gaagaatcga  ttcctcaggc  cagggtgatga  ctccagccac  gacctcatgc
   421 gtcagagcct  gccgagctca  cggatgctgt  gaaggctcatg  gacctgcca
   481 agcactgggg  accacctgct  acgcctcagg  ctggggcagc  attgaaccag
20  541 gaccccaaag  aaacttcagt  gtgtggacct  ccatgttatt  tccaatgacg
   601 agttcacctc  cagaagggtg  ccaagttcat  gctgtgtgtc  ggacgctgga
25  661 aagcacctgc  tcgggtgatt  ctgggggccc  acttgtctgt  aatggtgtgc
   721 cacgtcatgg  ggcagtgaac  catgtgcctt  gcccgaaagg  ccttccctgt
   781 ggtgcattac  cggaagtgga  tcaaggacac  catcgtggcc  aacccctgag
30  841 aacccctat  tgtagtaaac  ttggaacctt  ggaaatgacc  agccaagac
   901 ccagttctac  tgacctttgt  ccttaggtgt  gaggtccagg  gttgctagga
35  961 gcagacacag  gtgtagacca  gagtgtttct  taaatggtgt  aattttgtcc
   1021 ctggggaata  ctggccatgc  ctggagacat  atcactcaat  ttctctgagg
   1081 gatggggtgt  ctgtgttatt  tgtggggtac  agagatgaaa  gagggggtgg
40  1141 agagagtgga  gagtgcacat  tgctggacac  tgtccatgaa  gactgagca
   1201 gcacaacgca  ccagacactc  acagcaagga  tggagctgaa  aacataaccc
45  1261 ggaggcactg  ggaagcctag  agaaggctgt  gagccaagga  gggagggtct
   1321 tgggatgggg  atgaagtaag  gagagggact  ggacccctct  gaagctgatt
   1381 ggagggtgtat  tgaagtcctc  cagacaaccc  tcagatttga  tgatttccta
50  1441 cagaaataaa  gagctgttat  actgtg
   //

```

55

U87459. Human autoimmunog...[gi:1890098]

LOCUS

HSU87459

752 bp

mRNA

PRI

22-DEC-1999

DEFINITION Human autoimmunogenic cancer/testis antigen NY-ESO-1 mRNA, complete cds.
 ACCESSION U87459
 VERSION U87459.1 GI:1890098
 5 KEYWORDS .
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata;
 10 Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini;
 Hominidae; Homo.
 REFERENCE 1 (bases 1 to 752)
 AUTHORS Chen,Y.T., Scanlan,M.J., Sahin,U., Tureci,O.,
 15 Gure,A.O., Tsang,S.,
 Williamson,B., Stockert,E., Pfreundschuh,M. and
 Old,L.J.
 TITLE A testicular antigen aberrantly expressed in human
 cancers detected
 by autologous antibody screening
 20 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 94 (5), 1914-1918 (1997)
 MEDLINE 97203161
 PUBMED 9050879
 REFERENCE 2 (bases 1 to 752)
 AUTHORS Chen,Y.-T.
 25 TITLE Direct Submission
 JOURNAL Submitted (28-JAN-1997) Ludwig Institute for Cancer
 Research, New
 York Branch, 1275 York Avenue, New York, NY 10021,
 30 USA
 FEATURES Location/Qualifiers
 source 1..752
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /clone_lib="esophageal squamous cell
 35 carcinoma expression
 cDNA library"
 CDS 54..596
 /codon_start=1
 /product="autoimmunogenic cancer/testis
 40 antigen NY-ESO-1"
 /protein_id="AAB49693.1"
 /db_xref="GI:1890099"
 /translation="MQAEGRGTTGGSTGDADGPGGPGIPDGPGGNAGGPGEAG
 45 ATGGRGPRGAGAARASGPGGGAPRGPHGGAASGLNGCCRCGARGPESRLLEF
 YLAMPFATPMEABLARRSLAQDAPPLPVPGVLLKEFTVSGNILTIRLTAADH
 RQLQLSISSCLQQLSLLMWITQCFLPVFLAQPPSGQRR"
 BASE COUNT 126 a 230 c 256 g 140 t
 ORIGIN
 50 1 atcctcgtgg gccctgacct tctctctgag agccgggcag aggcctccgga
 gccatgcagg
 61 ccgaaggccg gggcacaggg gggtcgacgg gcgatgctga tggcccagga
 ggccctggca
 121 ttctgatgg cccagggggc aatgctggcg gcccaggaga ggcgggtgcc
 55 acgggcgggca
 181 gaggtccccg gggcgaggg gcagcaaggg cctcggggcc gggaggaggg
 gccccgcggg

```

      241 gtccgcatgg cggcgccggct tcagggctga atggatgctg cagatgcggg
gccaggggggc
      301 cggagagccg cctgcttgag ttctacctcg ccatgccttt cgcgacaccc
atggaagcag
5      361 agctggcccg caggagcctg gcccaggatg cccaccgct tcccggtcca
gggggtgcttc
      421 tgaaggagtt cactgtgtcc ggcaacatac tgactatccg actgactgct
gcagaccacc
      481 gccaaactgca gctctccatc agctcctgtc tccagcagct ttccctgttg
10 atgtggatca
      541 cgcagtgctt tctgcccggtg tttttggctc agcctccctc agggcagagg
cgctaagccc
      601 agcctggcgc cccttcctag gtcatgcctc ctcccctagg gaatggtccc
agcacgagtg
15      661 gccagttcat tgtggggggcc tgattgtttg tcgctggagg aggacggctt
acatgtttgt
      721 ttctgtagaa aataaaaactg agctacgaaa aa
//

20 CAA11116. LAGE-1a protein [...[gi:3255959]
LOCUS CAA11116 180 aa PRI
23-JUN-1998
DEFINITION LAGE-1a protein [Homo sapiens].
ACCESSION CAA11116
25 PID g3255959
VERSION CAA11116.1 GI:3255959
DBSOURCE embl locus HOS223093, accession AJ223093.1
KEYWORDS .
SOURCE human.
30 ORGANISM Homo sapiens
Eukaryota; Metazoa; Chordata; Craniata;
Vertebrata; Euteleostomi;
Mammalia; Eutheria; Primates; Catarrhini;
Hominidae; Homo.
35 REFERENCE 1 (residues 1 to 180)
AUTHORS Lethe,B., Lucas,S., Michaux,L., De Smet,C.,
Godelaine,D.,
Serrano,A., De Plaen,E. and Boon,T.
TITLE LAGE-1, a new gene with tumor specificity
40 JOURNAL Int. J. Cancer 76 (6), 903-908 (1998)
MEDLINE 98289662
REFERENCE 2 (residues 1 to 180)
AUTHORS Lethe,B.G.
TITLE Direct Submission
45 JOURNAL Submitted (08-JAN-1998) Lethe B.G., Brussels Branch,
Ludwig
Institute for Cancer Research, 74 Avenue
Hippocrate, B - 1200 -
Bruxelles, BELGIUM
50 COMMENT Related sequences: AJ223040, AJ223041 and AJ003149.
FEATURES Location/Qualifiers
source 1..180
/organism="Homo sapiens"
/isolate="individual LB33"
55 /db_xref="taxon:9606"
/chromosome="X"
/map="q28"
/clone="c2RB"

```

```

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040

```

```

                    /chromosome="X"
                    /map="q28"
                    /clone="c2RB"
                    /sex="Female"
5                 /cell_line="LB33-MELA"
                    /cell_type="melanoma"
                    /clone_lib="B6"
                Protein 1..210
                    /product="LAGE-1b protein"
10                CDS 1..210
                    /gene="LAGE-1"
                    /db_xref="SPTREMBL:O75638"
                    /coded_by="join(AJ223093.1:1231..1499,
15                AJ223093.1:2115..2478)"
                    /note="alternate splicing at exon 2 - intron
                2 boundary"
                ORIGIN
                    1 mqaegrgtgg stgdadgpgg pgipdgpggn aggpgeagat ggrgprgaga
                arasgprgga
20                61 prgphggaas aqdgrcpcga rrpdsrllel hitmpfsspm eaelvrrils
                rdaaplprpg
                    121 avlkdfvtvg nllfmsvwdq dregagrmrv vgwglgsasp egqkardlrt
                pkhkvsegrp
                    181 gtpgpppppeg aggdgcrgva fnvmfsaphi
25                //

30                M77481. Human antigen (MA...[gi:416114]
                LOCUS HUMMAG1A 2420 bp DNA PRI
                15-NOV-1993
                DEFINITION Human antigen (MAGE-1) gene, complete cds.
                ACCESSION M77481
35                VERSION M77481.1 GI:416114
                KEYWORDS antigen.
                SOURCE Homo sapiens (individual_isolate patient MZ2)
                melanoma abdominal
                    metastasis of melanoma DNA.
40                ORGANISM Homo sapiens
                    Eukaryota; Metazoa; Chordata; Craniata;
                Vertebrata; Euteleostomi;
                    Mammalia; Eutheria; Primates; Catarrhini;
                Hominidae; Homo.
45                REFERENCE 1 (bases 785 to 1286)
                AUTHORS van der Bruggen,P., Traversari,C., Chomez,P.,
                Lurquin,C., De
                    Plaen,E., Van den Eynde,B., Knuth,A. and Boon,T.
                TITLE A gene encoding an antigen recognized by cytolytic T
50                lymphocytes on
                    a human melanoma
                JOURNAL Science 254, 1643-1647 (1991)
                MEDLINE 92086861
                REFERENCE 2 (bases 1 to 2420)
55                AUTHORS van der Bruggen P.
                TITLE Direct Submission
                JOURNAL Submitted (05-FEB-1992) Pierre van der Bruggen, Ludwig
                Institute

```

for Cancer Research, Brussels Branch, Avenue
Hippocrate, 74, UCL
7459, Brussels, B-1200, Belgium

COMMENT On Nov 15, 1993 this sequence version replaced
gi:187294.

5 FEATURES Location/Qualifiers

source 1..2420
/organism="Homo sapiens"
/db_xref="taxon:9606"
10 /map="X"
/sex="female"
/cell_line="MZ2-MEL.43"

intron 1..412
15 /partial
/gene="MAGE-1"
/number=1

exon 413..485
/gene="MAGE-1"
/number=2

20 mRNA join(413..485,561..2111)
/partial
/gene="MAGE-1"

intron 486..560
25 /gene="MAGE-1"
/number=2

exon 561..2111
gene="MAGE-1"
/number=3

30 gene 626..1555
/gene="MAGE-1"

CDS 626..1555
35 /gene="MAGE-1"
/codon_start=1
/product="MAGE-1"
/protein_id="AAA03229.1"
/db_xref="GI:416115"

/translation="MSLEQRLHCKPEEALEAQQEALGLVCVQAATSSSSPL
40 VLGTLEEVPTAGSTDPPQSGASAFPTTINFTRQRPSEGSSSREEEGPST
SCILESIFRAVITKKVADLVGFLLKRYRAREPVTKAEMLESVIKYNKHCFP
IFGKASESLQLVFGIDVKEADPTGHSYVLVTCLGLSYDGLLDNQIMPKTGF
LIIVLVMIAEGGHAPEEEIWEELSVMEVYDGREHSAYGEPRKLLTQDLVQE
KYLEYRQVPDSDPARYEFLLWGPRLAETS YVKVLEYVIKVSARVRFFFP
EALREEEGV"

45 polyA_signal 2095..2100
/gene="MAGE-1"

intron 2112..2420
/partial
50 gene="MAGE-1"
/number=3

BASE COUNT 562 a 582 c 677 g 599 t
ORIGIN

1 ggatccaggc cctgccagga aaaatataag ggccctgcgt gagaacagag
gggggtcatcc

55 61 actgcatgag agtggggatg tcacagagtc cagcccaccc tcctggtagc
actgagaagc
121 cagggctgtg cttgcggtct gcaccctgag ggcccgtgga ttctctctcc
tggagctcca

	181	ggaaccaggc	agtgaggcct	tggtctgaga	cagtatcctc	aggtcacaga
	gcagaggatg					
	241	cacaggggtgt	gccagcagtg	aatgtttgcc	ctgaatgcac	accaagggcc
	ccacctgccca					
5	301	caggacacat	aggactccac	agagtctggc	ctcacctccc	tactgtcagt
	cctgtagaat					
	361	cgacctctgc	tgggcggctg	tacctgagt	accctctcac	ttcctccttc
	agggtttcag					
	421	gggacaggcc	aaccagagg	acaggattcc	ctggaggcca	cagaggagca
10	ccaaggagaa					
	481	gatctgtaag	taggcctttg	ttagagtctc	caaggttcag	ttctcagctg
	aggcctctca					
	541	cacactccct	ctctccccag	gcctgtgggt	cttcattgcc	cagctcctgc
	ccacactcct					
15	601	gcctgctgcc	ctgacgagag	tcatcatgtc	tcttgagcag	aggagtctgc
	actgcaagcc					
	661	tgaggaagcc	cttgaggccc	aacaagaggc	cctgggcctg	gtgtgtgtgc
	aggctgccac					
	721	ctcctcctcc	tctcctctgg	tctgggcac	cctggaggag	gtgccactg
20	ctgggtcaac					
	781	agatcctccc	cagagtcctc	agggagcctc	cgcccttccc	actaccatca
	acttcactcg					
	841	acagaggcaa	cccagtgagg	gttccagcag	ccgtgaagag	gaggggcca
	gcacctcttg					
25	901	tatcctggag	tccttggtcc	gagcagtaat	cactaagaag	gtggctgatt
	tggttggttt					
	961	tctgctcctc	aaatatcgag	ccaggagacc	agtcacaaag	gcagaaatgc
	tggagagtgt					
	1021	catcaaaaat	tacaagcact	gttttcctga	gatcttcggc	aaagcctctg
30	agtccttgca					
	1081	gctgggtcttt	ggcattgacg	tgaaggaagc	agacccacc	ggccactcct
	atgtccttgt					
	1141	cacctgccta	ggtctctcct	atgatggcct	gctgggtgat	aatcagatca
	tgcccaagac					
35	1201	aggcttcctg	ataattgtcc	tggtcatgat	tgcaatggag	ggcgcccatg
	ctcctgagga					
	1261	ggaaatctgg	gaggagctga	gtgtgatgga	ggtgtatgat	gggagggagc
	acagtgccta					
	1321	tggggagccc	aggaagctgc	tcacccaaga	tttggtgacg	gaaaagtacc
40	tggagtaccg					
	1381	gcagggtgccc	gacagtgate	ccgcacgcta	tgagttcctg	tgggggtccaa
	gggccttcgc					
	1441	tgaaaccagc	tatgtgaaag	tccttgagta	tgtgatcaag	gtcagtgcaa
	gagttcgctt					
45	1501	tttcttccca	tccttgctg	aagcagcttt	gagagaggag	gaagagggag
	tctgagcatg					
	1561	agttgcagcc	aaggccagtg	ggagggggac	tgggccagtg	caccttcag
	ggccgcgtcc					
	1621	agcagcttcc	cctgcctcgt	gtgacatgag	gccattctt	cactctgaag
50	agagcggtca					
	1681	gtgttctcag	tagtaggttt	ctgttctatt	gggtgacttg	gagatttatc
	tttgttctct					
	1741	tttggaattg	ttcaaagtgt	tttttttaag	ggatggttga	atgaacttca
	gcatccaagt					
55	1801	ttatgaatga	cagcagtcac	acagttctgt	gtatatagtt	taagggtaag
	agtcttgtgt					
	1861	tttattcaga	ttgggaaatc	cattctatct	tgtgaattgg	gataataaca
	gcagtggaaat					

```

1921 aagtacttag aaatgtgaaa aatgagcagt aaaatagatg agataaagaa
ctaaagaaat
1981 taagagatag tcaattcttg ccttatacct cagtctattc tgtaaaattt
ttaaagatat
5 2041 atgcatacct ggatttcctt ggcttctttg agaatgtaag agaaattaa
tctgaataaaa
2101 gaattcttcc tgttcactgg ctcttttctt ctccatgcac tgagcatctg
ctttttggaa
2161 ggccctgggt tagtagtgga gatgctaagg taagccagac tcataccac
10 ccatagggtc
2221 gtagagtcta ggagctgcag tcacgtaatc gaggtggcaa gatgtcctct
aaagatgtag
2281 ggaaaagtga gagaggggtg aggggtgtggg gctccgggtg agagtgggtg
15 agtgtcaatg
2341 ccctgagctg gggcattttg ggctttggga aactgcagtt ccttctgggg
gagctgattg
2401 taatgatctt ggggtggatcc
//

20 L18920. Human MAGE-2 gene...[gi:436180]
LOCUS HUMMAGE2X 4559 bp DNA PRI
20-APR-1994
DEFINITION Human MAGE-2 gene exons 1-4, complete cds.
ACCESSION L18920
25 VERSION L18920.1 GI:436180
KEYWORDS .
SOURCE Homo sapiens (human).
ORGANISM Homo sapiens
30 Eukaryota; Metazoa; Chordata; Craniata;
Vertebrata; Euteleostomi;
Mammalia; Eutheria; Primates; Catarrhini;
Hominidae; Homo.
REFERENCE 1 (bases 1 to 4559)
AUTHORS De Smet,C., Lurquin,C., van der Bruggen,P., De
35 Plaen,E.,
Brasseur,F. and Boon,T.
TITLE Sequence and expression pattern of the human MAGE2 gene
JOURNAL Immunogenetics 39 (2), 121-129 (1994)
MEDLINE 94102805
40 FEATURES Location/Qualifiers
source 1..4559
/organism="Homo sapiens"
/isolate="MZ-2"
45 /db_xref="taxon:9606"
/map="X"
/haplotype="A1 A29 B37 B44 C6"
/sex="female"
/cell_line="MZ2-MEL.43"
/cell_type="melanoma cell line"
50 gene 561..4496
/gene="MAGE-2"
exon 561..754
/gene="MAGE-2"
/number=1
55 intron 755..1850
/gene="MAGE-2"
/number=1
exon 1851..1969

```



```

                    /gene="MAGE-2"
                    /number=2
intron            1970..2788
                    /gene="MAGE-2"
5                /number=2
exon              2789..2854
                    /gene="MAGE-2"
                    /number=3
intron            2855..2934
10               /gene="MAGE-2"
                    /number=3
exon              2935..4496
                    /gene="MAGE-2"
                    /number=4
15               CDS            3000..3944
                    /gene="MAGE-2"
                    /codon_start=1
                    /protein_id="AAA17729.1"
                    /db_xref="GI:436181"
20
    /translation="MPLEQRSQHCKPEEGLEARGEALGLVGAQAPATEEQQTASSSSSTLVEVT
LGEVPAADSPSPPHSPQGASSFSTTINYTLWRQSDGSSNQEEGPRMFPDLE
SEFQA AISRKMVELVHFLLLKYRAREPVTKAEMLESVLRNCQDFFPVIFSKASEYLQLVFGIE
VVEVVPISHLYILVTCLGLSYDGLLDGNQVMPKTGLLIIVLAI IAEGDCAPEEKIWEELSML
25 EVFEGREDSVFAHPRKLLMQDLVQENYLEYRQVPGSDPACYEFLWGPRALIETSYVKVLHHTL
KIGGEPHISYPPLHERALREGEE"
polyA_signal      4479..4484
                    /gene="MAGE-2"
BASE COUNT        1049 a    1280 c    1283 g    947 t
30 ORIGIN
      1 attccttcat caaacagcca ggagtgagga agaggaccct cctgagtgag
gactgaggat
      61 ccaccctcac cacatagtgg gaccacagaa tccagctcag cccctcttgt
cagccctggg
35      121 acacactggc aatgatctca ccccgagcac acccctcccc ccaatgccac
ttcggggcga
      181 ctcagagtca gagacttggg ctgagggggag cagacacaat cggcagagga
tggcggtcca
      241 ggctcagtct ggcattcaag tcaggacctt gagggatgac caaaggcccc
40 tcccccccc
      301 aactcccccg accccaccag gatctacagc ctcaggatcc ccgtcccaat
ccctaccct
      361 acaccaacac catcttcatg cttaccccca cccccccatc cagatcccca
tccgggcaga
45      421 atccggttcc acccttgccg tgaaccagag gaagtcacgg gcccgatgt
gacgccactg
      481 acttgacatc tggagggtcag aggacagcga gattctcgcc ctgagcaacg
gcctgacgtc
      541 ggcggaggga agcaggcgca ggctccgtga ggaggcaagg taagacgccg
50 agggaggact
      601 gagggcgggc tcaccccgag cagagggccc ccaataatcc agcgctgcct
ctgctgccgg
      661 gcctggacca ccctgcaggg gaagacttct caggctcagt cggcaccacc
tcaccccgcc
55      721 accccccgcc gctttaaccg cagggaactc tggcgtaaga gctttgtgtg
accagggcag
      781 ggctgggttag aagtgtcag ggcccagact cagccaggaa tcaagggtcag
gacccaaga

```

	841	ggggactgag	ggcaaccac	cccctaccct	cactaccaat	cccatcccc
		aacaccaacc				
	901	ccacccccat	ccctcaaaca	ccaaccccac	ccccaacccc	cattccccatc
		tcctccccca				
5	961	ccaccatcct	ggcagaatcc	ggctttgccc	ctgcaatcaa	cccacggaag
		ctccgggaat				
	1021	ggcggccaag	cacgcggatc	ctgacgttca	catgtacggc	taagggaggg
		aaggggttgg				
10	1081	gtctcgtgag	tatggccttt	gggatgcaga	ggaagggccc	aggcctcctg
		gaagacagtg				
	1141	gagtccttag	gggacccagc	atgccaggac	agggggccca	ctgtaccctt
		gtctcaaact				
	1201	gagccacctt	ttcattcagc	cgagggaaac	ctagggatgc	agaccactt
		cagcaggggg				
15	1261	ttggggccca	gcctgcgagg	agtcaagggg	aggaagaaga	gggaggactg
		aggggacctt				
	1321	ggagtccaga	tcagtggcaa	ccttgggctg	ggggatcctg	ggcacagtgg
		ccgaatgtgc				
	1381	cccgtgctca	ttgcaccttc	agggtgacag	agagttgagg	gctgtggtct
20		gagggctggg				
	1441	acttcaggtc	agcagagggg	ggaatcccag	gatctgccgg	acccaagggtg
		tgcccccttc				
	1501	atgaggactg	gggatacccc	cggcccagaa	agaagggatg	ccacagagtc
		tggaagtccc				
25	1561	ttgttcttag	ctctggggga	acctgatcag	ggatggccct	aagtgacaat
		ctcatttgta				
	1621	ccacaggcag	gaggttgggg	aaccctcagg	gagataaggt	gttgggtgaa
		agaggagctg				
	1681	tctgctcatt	tcaggggggt	gggggttgag	aaagggcagt	ccctggcagg
30		agtaaagatg				
	1741	agtaaccac	aggaggccat	cataacgttc	accctagaac	caaaggggtc
		agccctggac				
	1801	aacgcacgtg	ggggtaacag	gatgtggccc	ctcctcactt	gtctttccag
		atctcagga				
35	1861	gttgatgacc	ttgttttcag	aaggtgactc	aggtcaacac	aggggccccca
		tctggtcgac				
	1921	agatgcagtg	gttctaggat	ctgccaagca	tccagggtga	gagcctgagg
		taggattgag				
40	1981	ggtacccctg	ggccagaatg	cagcaagggg	gccccataga	aatctgcctt
		gcccctgcgg				
	2041	ttacttcaga	gaccctgggc	agggctgtca	gctgaagtcc	ctccattatc
		ctgggatctt				
	2101	tgatgtcagg	gaaggggagg	ccttggtctg	aaggggctgg	agtcagggtca
		gtagagggag				
45	2161	ggtctcaggc	cctgccagga	gtggacgtga	ggaccaagcg	gactcgtcac
		ccaggacacc				
	2221	tggactccaa	tgaatttgga	catctctcgt	tgtccttcgc	gggaggacct
		ggtcacgtat				
	2281	ggccagatgt	gggtcccttc	atatccttct	gtaccatata	agggatgtga
50		gttcttgaca				
	2341	tgagagattc	tcaagccagc	aaaaggggtg	gattaggccc	tacaaggaga
		aaggtgaggg				
	2401	ccctgagtga	gcacagaggg	gaccctccac	ccaagtagag	tggggacctc
		acggagtctg				
55	2461	gccaaccctg	ctgagacttc	tgggaatccg	tggctgtgct	tgcagtctgc
		acactgaagg				
	2521	cccgtgcatt	cctctcccag	gaatcaggag	ctccaggaac	caggcagtga
		ggccttggtc				

	2581	tgagtcagtg	tcctcaggtc	acagagcaga	ggggacgcag	acagtgccaa
	cactgaaggt					
	2641	ttgcctggaa	tgacacccaa	gggccccacc	cgcccagAAC	aaatgggact
	ccagaaggcc					
5	2701	tggcctcacc	ctccctattc	tcagtcctgc	agcctgagca	tgtgctggcc
	ggctgtaccc					
	2761	tgaggtgccc	tcccacttcc	tccttcaggt	tctgaggggg	acaggctgac
	aagtaggacc					
10	2821	cgaggcactg	gaggagcatt	gaaggagaag	atctgtaagt	aagcctttgt
	cagagcctcc					
	2881	aaggttcagt	tcagttctca	cctaaggcct	cacacacgct	ccttctctcc
	ccaggcctgt					
	2941	gggtcttcat	tgcccagctc	ctgcccgcac	tcctgcctgc	tgccctgacc
	agagtcacat					
15	3001	tgccctcttga	gcagaggagt	cagcactgca	agcctgaaga	aggccttgag
	gcccaggag					
	3061	aggccctggg	cctgggtgggt	gcgcaggctc	ctgctactga	ggagcagcag
	accgcttctt					
	3121	cctctttctac	tctagtggaa	gttaccctgg	gggaggtgcc	tgctgccgac
20	tcaccgagtc					
	3181	ctccccacag	tcctcagga	gcctccagct	tctcgactac	catcaactac
	actctttgga					
	3241	gacaatccga	tgagggctcc	agcaaccaag	aagaggaggg	gccaagaatg
	tttcccgacc					
25	3301	tgagtcgga	gttccaagca	gcaatcagta	ggaagatggg	tgagttgggt
	cattttctgc					
	3361	tcctcaagta	tcgagccagg	gagccggctc	caaaggcaga	aatgctggag
	agtgtcctca					
	3421	gaaattgcca	ggacttcttt	cccgtgatct	tcagcaaagc	ctccgagtac
30	ttgcagctgg					
	3481	tctttggcat	cgaggtgggt	gaagtgggtc	ccatcagcca	cttgtacatc
	cttgtcacct					
	3541	gcctgggcct	ctcctacgat	ggcctgctgg	gcgacaatca	ggtcatgccc
	aagacaggcc					
35	3601	tcctgataat	cgctcctggc	ataatcgcaa	tagagggcga	ctgtgcccct
	gaggagaaaa					
	3661	tctgggagga	gctgagtatg	ttggaggtgt	ttgaggggag	ggaggacagt
	gtcttcgcac					
	3721	atcccaggaa	gctgctcatg	caagatctgg	tgacaggaaa	ctacctggag
40	taccggcagg					
	3781	tgcccggcag	tgatcctgca	tgctacgagt	tcctgtgggg	tccaagggcc
	ctcattgaaa					
	3841	ccagctatgt	gaaagtccct	caccatacac	taaagatcgg	tggagaacct
	cacatttcct					
45	3901	acccacccct	gcctgaacgg	gctttgagag	agggagaaga	gtgagtctca
	gcacatgttg					
	3961	cagccagggc	cagtgggagg	gggtctgggc	cagtgcacct	tccagggccc
	catccattag					
	4021	cttcactgct	ctcgtgtgat	atgaggccca	ttcctgcctc	tttgaagaga
50	gcagtcagca					
	4081	ttcttagcag	tgagtttctg	ttctgttgga	tgactttgag	atttatcttt
	ctttcctggt					
	4141	ggaattgttc	aaatgttctc	tttaacaaat	ggttgatga	acttcagcat
	ccaagtttat					
55	4201	gaatgacagt	agtcacacat	agtgtgtttt	atatagttta	gggtaagag
	tcctgttttt					
	4261	tattcagatt	gggaaatcca	ttccattttg	tgagttgtca	cataataaca
	gcagtggaa					

```

      4321 atgtatttgc ctatattgtg aacgaattag cagtaaaata catgatacaa
ggaactcaaa
      4381 agatagttaa ttcttgcctt atacctcagt ctattatgta aaattaaana
tatgtgtatg
5      4441 tttttgcttc tttgagaatg caaaagaaat taaatctgaa taaattcttc
ctgttctactg
      4501 gctcatttct ttaccattca ctcagcatct gctctgtgga aggccctggt
agtagtggg
//
10
U03735. Human MAGE-3 anti...[gi:468825]
LOCUS          HSU03735          4204 bp          DNA          PRI
07-APR-1994
DEFINITION Human MAGE-3 antigen (MAGE-3) gene, complete cds.
15 ACCESSION U03735
VERSION U03735.1 GI:468825
KEYWORDS
SOURCE human.
ORGANISM Homo sapiens
20      Eukaryota; Metazoa; Chordata; Craniata;
Vertebrata; Euteleostomi;
      Mammalia; Eutheria; Primates; Catarrhini;
Hominidae; Homo.
REFERENCE 1 (bases 1 to 4204)
25 AUTHORS Gaugler,B., Van den Eynde,B., van der Bruggen,P.,
Romero,P.,
      Gaforio,J.J., De Plaen,E., Lethe,B., Brasseur,F.
and Boon,T.
TITLE Human gene MAGE-3 codes for an antigen recognized on
30 a melanoma by
      autologous cytolytic T lymphocytes
JOURNAL J. Exp. Med. 179, 921-930 (1994)
MEDLINE 94157413
REFERENCE 2 (bases 1 to 4204)
35 AUTHORS Gaugler,B.
TITLE Direct Submission
JOURNAL Submitted (25-NOV-1993) Beatrice Gaugler, Ludwig
Institute for
      Cancer Research, 74 Avenue Hippocrate, Brussels
40 1200, Belgium
FEATURES Location/Qualifiers
      source 1..4204
      /organism="Homo sapiens"
      /isolate="patient MZ2"
45      /db_xref="taxon:9606"
      /chromosome="X"
      /sex="female"
      /cell_type="lymphocyte"
      /tissue_type="blood"
50      /dev_stage="adult"
      exon 434..468
      /number=1
      exon 2254..2319
      /number=2
55      exon 2400..3978
      /number=3
      gene 2465..3409
      /gene="MAGE-3"

```

```

CDS                2465..3409
                    /gene="MAGE-3"
                    /codon_start=1
                    /product="MAGE-3 antigen"
5                   /protein_id="AA17446.1"
                    /db_xref="GI:468826"

/translation="MPLEQRSQHCKPEEGLEARGEALGLVGAQAPATEEQEAASSSSTLVEVTLGE
10 VPAAESPDPPQSPQGASSLPTTMNYPLWSQSYEDSSNQEEGPSTFPDLESEFQAALSRKVAELVH
  FLLKLYRAREPVTKAEMLGSVVGNWQYFFPVIFSKASSSLQLVFGIELMEVDPIGHLYIFATCLGL
  SYDGLLGDNQIMPKAGLLIIVLAI IAREGDCAPEEKIWEELSVLEVFEGREDSILGDPKKLLTQHF
  VQENYLEYRQVPGSDPACYEFLWGPRALVETSYVKVLHMHVKISGGPHISYPPLHEWVLREGEE"
      polyA_signal    3958..3963
      polyA_site      3979
15  BASE COUNT      944 a   1144 c   1223 g    893 t
  ORIGIN
      1  acgcaggcag  tgatgtcacc  cagaccacac  cccttcccc  aatgccactt
  caggggggtac
      61  tcagagtcag  agacttggtc  tgaggggagc  agaagcaatc  tgcagaggat
20  ggcgggtccag
      121  gctcagccag  gcatcaactt  caggaccctg  agggatgacc  gaaggccccg
  cccaccacc
      181  cccaactccc  ccgacccac  caggatctac  agcctcagga  ccccgctccc
  aatccttacc
25  241  ccttgcccc  tcaccatctt  catgcttacc  tccaccccc  tccgatcccc
  atccaggcag
      301  aatccagttc  caccctgcc  cggaaccag  ggtagtaccg  ttgccaggat
  gtgacgccac
      361  tgacttgccg  attggaggtc  agaagaccgc  gagattctcg  ccctgagcaa
30  cgagcgacgg
      421  cctgacgtcg  gcggagggaa  gccggcccag  gctcggtgag  gaggcaaggt
  aagacgctga
      481  gggaggactg  aggcgggcct  cacctcagac  agagggcctc  aaataatcca
  gtgctgcctc
35  541  tgctgccggg  cctgggccac  cccgcagggg  aagacttcca  ggctgggtcg
  ccactacctc
      601  accccgccga  ccccgccgc  tttagccacg  gggaaactctg  gggacagagc
  ttaatgtggc
      661  cagggcaggg  ctggttagaa  gaggtcaggg  cccacgctgt  ggcaggaatc
40  aaggtcagga
      721  ccccgagagg  gaactgaggg  cagcctaacc  accaccctca  ccaccattcc
  cgtcccccaa
      781  cacccaaccc  cacccccac  cccattccc  atccccacc  ccaccctat
  cctggcagaa
45  841  tccgggcttt  gcccctggta  tcaagtcacg  gaagctccgg  gaatggcggc
  caggcacgtg
      901  agtcctgagg  ttcacatcta  cggctaaggg  agggaagggg  ttcggtatcg
  cgagtatggc
      961  cgttgggagg  cagcgaaagg  gcccaggcct  cctggaagac  agtggagtcc
50  tgaggggacc
      1021  cagcatgcca  ggacaggggg  cccactgtac  ccctgtctca  aaccgaggca
  ccttttcatt
      1081  cggctacggg  aatcctaggg  atgcagaccc  acttcagcag  ggggttgggg
  cccagccctg
55  1141  cgaggagtca  tggggaggaa  gaagagggag  gactgagggg  acctggaggt
  ccagatcagt
      1201  ggcaaccttg  ggctggggga  tgctgggcac  agtggccaaa  tgtgctctgt
  gtcattgcg

```

	1261	ccttcagggt	gaccagagag	ttgagggctg	tggctctgaag	agtgggactt
		caggtcagca				
	1321	gagggaggaa	tcccaggatc	tgcagggccc	aaggtgtacc	cccaaggggc
5		ccctatgtgg				
	1381	tggacagatg	cagtggctct	aggatctgcc	aagcatccag	gtgaagagac
		tgagggaggaa				
	1441	ttgaggggtac	ccctgggaca	gaatgcggac	tggggggccc	ataaaaatct
		gccctgctcc				
10		1501	tgctgttacc	tcagagagcc	tgggcagggc	tgctcagctga
		ttatcctagg				ggccctcca
		1561	atcactgatg	tcaggggaagg	ggaagccttg	gtctgagggg
		gggcagtaga				gctgcactca
		1621	gggaggctct	cagaccctac	taggagtgga	ggtagaggacc
		ctcaccagg				aagcagtctc
15		1681	gtacatggac	ttcaataaat	ttggacatct	ctcgttgtcc
		gacctgggaa				tttccgggag
		1741	tgtatggcca	gatgtgggtc	ccctcatgtt	tttctgtacc
		tgtgagttct				atatcaggta
20		1801	tgacatgaga	gattctcagg	ccagcagaag	ggagggatta
		ggagaaagg				ggccctataa
		1861	gagggccctg	agtgagcaca	gaggggatcc	tccaccccag
		acctcacaga				tagagtgggg
		1921	gtctggccaa	ccctcctgac	agttctggga	atccgtggct
		tctgcacatt				gcgtttgctg
25		1981	gggggcccgt	ggattcctct	cccaggaatc	aggagctcca
		agtgaggact				ggaacaaggc
		2041	tggctctgagg	cagtgtcctc	aggtcacaga	gtagaggggg
		gccaacggtg				ctcagatagt
30		2101	aaggtttgcc	ttggattcaa	accaagggcc	ccacctgccc
		ggactccaga				cagaacacat
		2161	gcgcctggcc	tcaccctcaa	tactttcagt	cctgcagcct
		tggccggatg				cagcatgcgc
		2221	taccctgagg	tgccctctca	cttcctcctt	caggttctga
		tgacctggag				ggggacaggc
35		2281	gaccagaggc	ccccggagga	gcactgaagg	agaagatctg
		tttgtttagag				taagtaagcc
		2341	cctccaagg	tccattcagt	actcagctga	ggctctcac
		tctccccagg				atgctccctc
		2401	ccagtgggtc	tccattgccc	agctcctgcc	cacactcccg
40		tgaccagagt				cctggtgccc
		2461	catcatgcct	cttgagcaga	ggagtcagca	ctgcaagcct
		ttgaggccc				gaagaaggcc
		2521	aggagaggcc	ctgggcctgg	tgggtgcgca	ggctcctgct
		aggaggctgc				actgaggagc
45		2581	ctcctcctct	tctactctag	ttgaagtcac	cctggggggag
		ccgagtcacc				gtgcctgctg
		2641	agatcctccc	cagagtcctc	agggagcctc	cagcctcccc
		actaccctct				actaccatga
50		2701	ctggagccaa	tcctatgagg	actccagcaa	ccaagaagag
		gcaccttccc				gaggggccaa
		2761	tgacctggag	tccgagttcc	aagcagcact	cagtaggaag
		tggttcattt				gtggccgagt
		2821	tctgctcctc	aagtatcgag	ccagggagcc	ggtcacaaag
		tggggagtg				gcagaaatgc
55		2881	cgtcggaaat	tggcagtatt	tctttcctgt	gatcttcagc
		gttccttgca				aaagcttcca
		2941	gctggtcttt	ggcatcgagc	tgatggaagt	ggaccccatc
		acatctttgc				ggccacttgt

```

      3001 cacctgcctg ggcctctcct acgatggcct gctgggtgac aatcagatca
tgcceaaggc
      3061 aggcctcctg ataatcgctc tggccataat cgcaagagag ggcgactgtg
cccttgagga
5      3121 gaaaatcttg gaggagctga gtgtgttaga ggtgtttgag gggaggggaa
acagtatctt
      3181 gggggatccc aagaagctgc tcacccaaca tttcgtgcag gaaaactacc
tggagtaccg
      3241 gcagggtccc ggcagtgate ctgcatgtta tgaattcctg tggggtccaa
10 gggccctcgt
      3301 tgaaaccagc tatgtgaaag tcctgcacca tatggtaaag atcagtggag
gacctcacat
      3361 ttcctaccca ccctgcatg agtgggtttt gagagagggg gaagagtgtg
tctgagcacg
15      3421 agttgcagcc agggccagtg ggaggggggtc tgggccagtg caccttccgg
ggccgcctcc
      3481 cttagtcttc actgcctcct gtgacgtgag gccattctt cactctttga
agcgagcagt
      3541 cagcattctt agtagtgggt ttctgttctg ttggatgact ttgagattat
20 tctttgtttc
      3601 ctgttgaggt tgttcaaagt ttcttttaa cggatgggtg aatgagcgtc
agcatccagg
      3661 tttatgaatg acagtagtca cacatagtgc tgtttatata gtttaggagt
aagagtcttg
25      3721 ttttttactc aaattgggaa atccattcca ttttgtgaat tgtgacataa
taatagcagt
      3781 ggtaaaagta tttgcttaaa attgtgagcg aattagcaat aacatacatg
agataactca
      3841 agaaatcaaa agatagttaga ttcttgccct gtacctcaat ctattctgta
30 aaattaaaca
      3901 aatatgcaaa ccaggatttc cttgacttct ttgagaatgc aagcgaaatt
aaatctgaat
      3961 aaataattct tcctcttcac tggctcgttt cttttccgtt cactcagcat
ctgctctgtg
35      4021 ggaggccctg ggttagtagt ggggatgcta aggtaagcca gactcacgcc
taccatagg
      4081 gctgtagagc ctaggacctg cagtcatata attaaggtgg tgagaagtcc
tgtaagatgt
      4141 agaggaaatg taagagaggg gtgagggtgt ggcgctccgg gtgagagtag
40 tggagtgtca
      4201 gtgc
//

```

```

45 AF043498. Homo sapiens pros...[gi:2909843]
LOCUS AF043498 990 bp mRNA PRI 24-
FEB-1998
DEFINITION Homo sapiens prostate stem cell antigen (PSCA)
mRNA, complete cds.
ACCESSION AF043498
50 VERSION AF043498.1 GI:2909843
KEYWORDS .
SOURCE human.
ORGANISM Homo sapiens
55 Euteleostomi;
Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
Homo.
REFERENCE 1 (bases 1 to 990)

```

AUTHORS Reiter,R.E., Gu,Z., Watabe,T., Thomas,G., Kinga,S.,
 Davis,E., Wahl,M., Nisitani,S., Yamashiro,J., Le Beau,M.M.,
 Losa,M. and Witte,O.N.
 5 TITLE Prostate stem cell antigen: a cell surface marker
 overexpressed in
 prostate cancer
 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 95 (4), 1735-1740 (1998)
 10 MEDLINE 98132661
 REFERENCE 2 (bases 1 to 990)
 AUTHORS Reiter,R.E.
 TITLE Direct Submission
 JOURNAL Submitted (19-JAN-1998) Urology, UCLA, 66-134 CHS
 15 10833 Le Conte
 Ave., Los Angeles, CA 90095, USA
 FEATURES Location/Qualifiers
 source 1..990
 20 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /chromosome="8"
 /map="8q24.2"
 /note="LAPC-4 prostate cancer xenograft"
 gene 1..990
 25 /gene="PSCA"
 CDS 18..389
 /gene="PSCA"
 /note="GPI-anchored cell surface protein"
 /codon_start=1
 30 /product="prostate stem cell antigen"
 /protein_id="AAC39607.1"
 /db_xref="GI:2909844"

 /translation="MKAVLLALLMAGLALQPGTALLCYSCAQVSNEDCLQVENCTQLGEQCW
 35 TARIRAVGLLTVISKGCSLNCVDDSDYYVGKKNITCCDTDLNAGAHALQPAAAILALLPA
 LGLLLWGPGQL"
 BASE COUNT 193 a 299 c 285 g 202 t 11 others
 ORIGIN
 1 agggagagggc agtgaccatg aaggctgtgc tgcttgccct gttgatggca
 40 ggcttgggccc
 61 tgcagccagg cactgccctg ctgtgctact cctgcaaagc ccagggtgagc
 aacgaggact
 121 gcctgcaggt ggagaactgc acccagctgg gggagcagtg ctggaccgcg
 45 cgcacccgcg
 181 cagttggcct cctgaccgtc atcagcaaag gctgcagctt gaactgcgtg
 gatgactcac
 241 aggactacta cgtgggcaag aagaacatca cgtgctgtga caccgacttg
 tgcaacgcca
 301 gcgggggcca tgccctgcag cgggctgccg ccatccttgc gctgctccct
 50 gcactcggcc
 361 tgetgctctg gggaccgggc cagctatagg ctctgggggg ccccgctgca
 gccacactg
 421 ggtgtggtgc ccaggcctt tgtgccactc ctcacagaac ctggcccagt
 gggagcctgt
 55 481 cctggttcct gaggcacatc ctaacgcaag tttgaccatg tatgtttgca
 ccccttttcc
 541 ccnaaccctg accttcccat gggccttttc caggattccn accnggcaga
 tcagttttag


```

        601 tganacanat cgcntgcag atggccctc caacctttt tggtgntggt
tccatggccc
        661 agcattttcc acccttaacc ctgtgttcag gcacttnttc ccccaggaag
ccttccctgc
5       721 ccacccatt tatgaattga gccaggtttg gtccgtggtg tccccgcac
ccagcagggg
        781 acaggcaatc aggagggccc agtaaaggct gagatgaagt ggactgagta
gaactggagg
        841 acaagagttg acgtgagttc ctgggagttt ccagagatgg ggctggagg
10      cctggaggaa
        901 ggggccaggc ctcacatttg tggggntccc gaatggcagc ctgagcacag
cgtaggcct
        961 taataaacac ctgttgata agccaaaaaa
//
15
P06870. GLANDULAR KALLIKR...[gi:125170]
LOCUS      KLK1_HUMAN      262 aa                      PRI      20-
AUG-2001
DEFINITION      GLANDULAR KALLIKREIN 1 PRECURSOR (TISSUE
20      KALLIKREIN)
          (KIDNEY/PANCREAS/SALIVARY GLAND KALLIKREIN).
ACCESSION      P06870
PID            g125170
VERSION        P06870 GI:125170
25      DBSOURCE      swissprot: locus KLK1_HUMAN, accession P06870;
          class: standard.
          extra accessions:Q9UMJ1,created: Jan 1, 1988.
          sequence updated: Jan 1, 1988.
          annotation updated: Aug 20, 2001.
30      xrefs: gi: gi: 186652, gi: gi: 186653, gi: gi: 186649,
          gi: gi:
          186651, gi: gi: 186645, gi: gi: 186646, gi: gi:
          186647, gi: gi:
          186648, gi: gi: 34026, gi: gi: 34027, gi: gi: 186643,
35      gi: gi:
          386843, gi: gi: 67558
          xrefs (non-sequence databases): HSSP P00757, MEROPS
          S01.160,
          GlycoSuiteDB P06870, MIM 147910, InterPro IPR001314,
40      InterPro
          IPR001254, Pfam PF00089, PRINTS PR00722, PROSITE
          PS50240, PROSITE
          PS00134, PROSITE PS00135
KEYWORDS      Hydrolase; Serine protease; Glycoprotein; Multigene
45      family;
          Zymogen; Signal.
SOURCE      human.
          ORGANISM      Homo sapiens
          Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
50      Euteleostomi;
          Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
          Homo.
REFERENCE      1 (residues 1 to 262)
          AUTHORS      Fukushima,D., Kitamura,N. and Nakanishi,S.
55      TITLE      Nucleotide sequence of cloned cDNA for human
          pancreatic kallikrein
          JOURNAL      Biochemistry 24, 8037-8043 (1985)
          REMARK      SEQUENCE FROM N.A.

```

- TISSUE=Pancreas
 REFERENCE 2 (residues 1 to 262)
 AUTHORS Evans,B.A., Yun,Z.X., Close,J.A., Tregear,G.W.,
 Kitamura,N.,
 5 Nakanishi,S., Callen,D.F., Baker,E., Hyland,V.J.,
 Sutherland,G.R.
 and Richards,R.I.
 TITLE Structure and chromosomal localization of the human
 renal
 10 kallikrein gene
 JOURNAL Biochemistry. 27 (9), 3124-3129 (1988)
 MEDLINE 88269498
 PUBMED 2898948
 REMARK SEQUENCE FROM N.A.
 15 TISSUE=Kidney
 REFERENCE 3 (residues 1 to 262)
 AUTHORS Angermann,A., Bergmann,C. and Appelhans,H.
 TITLE Cloning and expression of human salivary-gland
 kallikrein in
 20 Escherichia coli
 JOURNAL The Biochemical journal. 262 (3), 787-793 (1989)
 MEDLINE 90073574
 PUBMED 2686621
 REMARK SEQUENCE FROM N.A.
 25 TISSUE=Salivary gland
 REFERENCE 4 (residues 1 to 262)
 AUTHORS Baker,A.R. and Shine,J.
 TITLE Human kidney kallikrein: cDNA cloning and sequence
 analysis
 30 JOURNAL DNA (Mary Ann Liebert, Inc.) 4 (6), 445-450 (1985)
 MEDLINE 86135264
 PUBMED 3853975
 REMARK SEQUENCE OF 17-262 FROM N.A.
 TISSUE=Kidney
 35 REFERENCE 5 (residues 1 to 262)
 AUTHORS Lu,H.S., Lin,F.K., Chao,L. and Chao,J.
 TITLE Human urinary kallikrein. Complete amino acid sequence
 and sites of
 glycosylation
 40 JOURNAL International journal of peptide and protein research.
 33 (4),
 237-249 (1989)
 MEDLINE 89326688
 PUBMED 2666327
 45 REMARK SEQUENCE OF 25-262.
 TISSUE=Urine
 REFERENCE 6 (residues 1 to 262)
 AUTHORS Kellermann,J., Lottspeich,F., Geiger,R. and
 Deutzmann,R.
 50 TITLE Human urinary kallikrein--amino acid sequence and
 carbohydrate
 attachment sites
 JOURNAL Protein sequences & data analysis. 1 (3), 177-182
 (1988)
 55 MEDLINE 88203586
 PUBMED 3163150
 REMARK SEQUENCE OF 25-262, AND CARBOHYDRATE-LINKAGE SITES.
 TISSUE=Urine

REFERENCE 7 (residues 1 to 262)
 AUTHORS Lottspeich,F., Geiger,R., Henschen,A. and Kutzbach,C.
 TITLE N-Terminal amino acid sequence of human urinary
 kallikrein homology
 5 with other serine proteases
 JOURNAL Hoppe-Seyler's Zeitschrift fur physiologische Chemie.
 360 (12),
 1947-1950 (1979)
 MEDLINE 80114126
 10 PUBMED 393608
 REMARK SEQUENCE OF 25-55.
 TISSUE=Urine
 REFERENCE 8 (residues 1 to 262)
 AUTHORS Takahashi,S., Irie,A., Katayama,Y., Ito,K. and
 15 Miyake,Y.
 TITLE N-terminal amino acid sequence of human urinary
 prokallikrein
 JOURNAL Journal of biochemistry. 99 (3), 989-992 (1986)
 MEDLINE 86223893
 20 PUBMED 3635530
 REMARK SEQUENCE OF 28-47.
 TISSUE=Urine
 [FUNCTION] GLANDULAR KALLIKREINS CLEAVE MET-LYS AND
 ARG-SER BONDS
 25 IN KININOGEN TO RELEASE LYS-BRADYKININ.
 [CATALYTIC ACTIVITY] PREFERENTIAL CLEAVAGE OF ARG-|-
 XAA BONDS IN
 SMALL MOLECULE SUBSTRATES. HIGHLY SELECTIVE ACTION TO
 30 RELEASE
 KALLIDIN (LYSYL-BRADYKININ) FROM KININOGEN INVOLVES
 HYDROLYSIS OF
 MET-|-XAA OR LEU-|-XAA.
 [SIMILARITY] BELONGS TO PEPTIDASE FAMILY S1; ALSO
 35 KNOWN AS THE
 TRYPSIN FAMILY. KALLIKREIN SUBFAMILY.
 FEATURES Location/Qualifiers
 source 1..262
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 40 1..262
 Protein 1..262
 /product="GLANDULAR KALLIKREIN 1 PRECURSOR"
 /EC_number="3.4.21.35"
 Region 1..18
 45 /region_name="Signal"
 /note="PROBABLE."
 Region 19..24
 /region_name="Propeptide"
 /note="ACTIVATION PEPTIDE (PROBABLE)."
 50 Region 25..262
 /region_name="Mature chain"
 /note="GLANDULAR KALLIKREIN 1."
 Bond bond(31,174)
 /bond_type="disulfide"
 55 Bond bond(50,66)
 /bond_type="disulfide"
 /note="BY SIMILARITY."

```

Site      65
           /site_type="active"
           /note="CHARGE RELAY SYSTEM."
5      Site      93
           /site_type="glycosylation"
           /note="O-LINKED."
      Site      102
           /site_type="glycosylation"
           /note="N-LINKED (GLCNAC...)."
10     Site      104
           /site_type="glycosylation"
           /note="O-LINKED."
      Site      108
           /site_type="glycosylation"
           /note="N-LINKED (GLCNAC...)."
15     Site      120
           /site_type="active"
           /note="CHARGE RELAY SYSTEM."
      Region     145
           /region_name="Variant"
           /note="Q -> E. /FTId=VAR_006625."
      Bond      bond(153,220)
           /bond_type="disulfide"
           /note="BY SIMILARITY."
25     Site      165
           /site_type="glycosylation"
           /note="N-LINKED (GLCNAC...) (PARTIAL)."
      Site      167
           /site_type="glycosylation"
           /note="O-LINKED."
30     Bond      bond(185,199)
           /bond_type="disulfide"
           /note="BY SIMILARITY."
      Region     186
           /region_name="Variant"
           /note="E -> K. /FTId=VAR_006626."
35     Bond      bond(210,235)
           /bond_type="disulfide"
           /note="BY SIMILARITY."
40     Site      214
           /site_type="active"
           /note="CHARGE RELAY SYSTEM."
ORIGIN
45     1  mwflvlclal slggtgaapp iqsrivggwe ceqhsqpwwa alyhfstfqc
      ggilvhrqvw
      61 ltaahcisdn yqlwlgrhnl fddentaqfv hvsesfphpg fnmsllenht
      rqadedyshd
      121 lmlrltpepa dtitdavkvv elptqepevg stclasgwgs iepenfsfpd
      dlqcvdlkil
50     181 pndecekahv qkvtdfmlcv ghleggkdtc vgdsggplmc dgvlqgvtsw
      gypvcgtpnk
      241 psvavrvlsy vkwiedtia ns
      //
55
P08217. ELASTASE 2A PRECU...[gi:119255]
LOCUS      EL2A_HUMAN      269 aa      PRI      20-
AUG-2001

```

DEFINITION ELASTASE 2A PRECURSOR.
 ACCESSION P08217
 PID g119255
 VERSION P08217 GI:119255
 5 DBSOURCE swissprot: locus EL2A_HUMAN, accession P08217;
 class: standard.
 created: Aug 1, 1988.
 sequence updated: Aug 1, 1988.
 annotation updated: Aug 20, 2001.
 10 xrefs: gi: gi: 182022, gi: gi: 182023, gi: gi: 182057,
 gi: gi:
 182058, gi: gi: 88298, gi: gi: 88299
 xrefs (non-sequence databases): MEROPS S01.155,
 InterPro IPR001314,
 15 InterPro IPR001254, Pfam PF00089, PRINTS PR00722,
 PROSITE PS50240,
 PROSITE PS00134, PROSITE PS00135
 KEYWORDS Hydrolase; Serine protease; Pancreas; Zymogen; Signal.
 SOURCE human.
 20 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
 Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
 Homo.
 25 REFERENCE 1 (residues 1 to 269)
 AUTHORS Kawashima,I., Tani,T., Shimoda,K. and Takiguchi,Y.
 TITLE Characterization of pancreatic elastase II cDNAs: two
 elastase II
 mRNAs are expressed in human pancreas
 30 JOURNAL DNA 6 (2), 163-172 (1987)
 MEDLINE 87217962
 REMARK SEQUENCE FROM N.A.
 REFERENCE 2 (residues 1 to 269)
 35 AUTHORS Fletcher,T.S., Shen,W.F. and Largman,C.
 TITLE Primary structure of human pancreatic elastase 2
 determined by
 sequence analysis of the cloned mRNA
 JOURNAL Biochemistry 26 (23), 7256-7261 (1987)
 40 MEDLINE 88107669
 REMARK SEQUENCE FROM N.A.
 [FUNCTION] ACTS UPON ELASTIN.
 [CATALYTIC ACTIVITY] PREFERENTIAL CLEAVAGE: LEU-|-XAA,
 MET-|-XAA
 45 AND PHE-|-XAA. HYDROLYSES ELASTIN.
 [SUBCELLULAR LOCATION] SECRETED.
 [TISSUE SPECIFICITY] PANCREAS.
 [SIMILARITY] BELONGS TO PEPTIDASE FAMILY S1; ALSO
 KNOWN AS THE
 TRYPSIN FAMILY. ELASTASE SUBFAMILY.
 50 FEATURES Location/Qualifiers
 source 1..269
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 Region 1..16
 55 /region_name="Signal"
 Protein 1..269
 /product="ELASTASE 2A PRECURSOR"
 /EC_number="3.4.21.71"

```

      Region      17..28
                  /region_name="Propeptide"
                  /note="ACTIVATION PEPTIDE."
5      Region      29..269
                  /region_name="Mature chain"
                  /note="ELASTASE 2A."
      Bond        bond(58,74)
                  /bond_type="disulfide"
                  /note="BY SIMILARITY."
10     Site       73
                  /site_type="active"
                  /note="CHARGE RELAY SYSTEM (BY SIMILARITY)."
```

15

```

      Site       121
                  /site_type="active"
                  /note="CHARGE RELAY SYSTEM (BY SIMILARITY)."
```

20

```

      Bond        bond(155,222)
                  /bond_type="disulfide"
                  /note="BY SIMILARITY."
      Bond        bond(186,202)
                  /bond_type="disulfide"
                  /note="BY SIMILARITY."
      Bond        bond(212,243)
                  /bond_type="disulfide"
                  /note="BY SIMILARITY."
25     Site       216
                  /site_type="active"
                  /note="CHARGE RELAY SYSTEM (BY SIMILARITY)."
```

ORIGIN

```

30     1 mirtlllsl1 vagalscgdp typpyvtrvv ggeearpnsw pwqvsllqyss
      ngkwyhtcgg
      61 slianswvlt aahci:sssrt yrvglgrhnl yvaesgslav svskivvhkd
      wnsnqiskgn
      121 diallklanp vsltldkiqla clppagtilp nnypcyvtgw grlqtngavp
      dvlqggrllv
35     181 vdyatc:ssa wwgssvktsm icaggdgvis scngdsggpl ncqasdgwrw
      vhgivsfgsr
      241 lgcnyyhkps vftrvsnyid winsviann
      //
```

40

```

NP_056933. pancreatic elasta...[gi:7705648]
LOCUS      NP_056933      269 aa                      PRI      02-
NOV-2000
DEFINITION pancreatic elastase IIB [Homo sapiens].
ACCESSION  NP_056933
45     PID      g7705648
      VERSION   NP_056933.1  GI:7705648
      DBSOURCE  REFSEQ: accession NM_015849.1
      KEYWORDS  .
      SOURCE    human.
50     ORGANISM Homo sapiens
                  Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
                  Euteleostomi;
                  Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
                  Homo.
55     REFERENCE 1 (residues 1 to 269)
      AUTHORS    Kawashima,I., Tani,T., Shimoda,K. and Takiguchi,Y.
      TITLE      Characterization of pancreatic elastase II cDNAs: two
                  elastase II
```

mRNAs are expressed in human pancreas

JOURNAL DNA 6 (2), 163-172 (1987)

MEDLINE 87217962

COMMENT PROVISIONAL REFSEQ: This record has not yet been
 5 subject to final
 NCBI review. The reference sequence was derived from
 M16653.1.

FEATURES

	Location/Qualifiers
10	source 1..269
	/organism="Homo sapiens"
	/db_xref="taxon:9606"
	/chromosome="12"
	/map="12"
15	Protein 1..269
	/product="pancreatic elastase IIB"
	sig_peptide 1..16
	/note="pancreatic elastase IIB signal
	peptide"
20	Region 28..262
	/region_name="Trypsin-like serine protease"
	/db_xref="CDD:Tryp_SpC"
	/note="Tryp_SpC"
	mat_peptide 29..269
25	/product="pancreatic elastase IIB mature
	peptide"
	Region 31..262
	/region_name="Trypsin"
	/db_xref="CDD:pfam00089"
	/note="trypsin"
30	CDS 1..269
	/gene="LOC51032"
	/db_xref="LocusID:51032"
	/coded_by="NM_015849.1:26..835"

ORIGIN

35 1 mirtlllstl vagalscgvs tyapdmsrml ggeearpnsw pwqvslyyss
 ngqwyhtcgg
 61 slianswvlt aahci:ssri yrvmlgqhn1 yvaesgslav svskivvhkd
 wnsnqvskgn
 121 diallkianp vsltldkiqla clppagtilp nnypcyvtgw grlqtngalp
 40 ddkkqgrllv
 181 vdyatcsssg wwgstvktnm icaggdgvic tcngdsggpl ncqasdgrwe
 vhgigsltsv
 241 lgcnyyykps iftrvsnynd winsviann
 //

45

PRAME

LOCUS NM_006115 2148 bp mRNA PRI 19-JUN-

50 2001

DEFINITION Homo sapiens preferentially expressed antigen in
 melanoma (PRAME),
 mRNA.

ACCESSION NM_006115

55 VERSION NM_006115.1 GI:5174640

KEYWORDS .

SOURCE human.

ORGANISM Homo sapiens

- Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
 Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
 Homo.
- 5 REFERENCE 1 (bases 1 to 2148)
 AUTHORS Ikeda,H., Lethe,B., Lehmann,F., Van Baren,N.,
 Baurain,J.-F., De
 Smet,C., Chambost,H., Vitale,M., Moretta,A., Boon,T. and
 Coulie,P.G.
- 10 TITLE Characterization of an antigen that is recognized on a
 melanoma
 showing partial HLA loss by CTL expressing an NK inhibitory
 receptor
- 15 JOURNAL Immunity 6 (2), 199-208 (1997)
 MEDLINE 97199265
 REFERENCE 2 (bases 1 to 2148)
 AUTHORS Williams JM, Chen GC, Zhu L and Rest RF.
 TITLE Using the yeast two-hybrid system to identify human
 epithelial cell
 20 proteins that bind gonococcal Opa proteins: intracellular
 gonococci
 bind pyruvate kinase via their Opa proteins and
 require host
 pyruvate for growth
- 25 JOURNAL Mol. Microbiol. 27 (1), 171-186 (1998)
 MEDLINE 98125741
 PUBMED 9466265
 REFERENCE 3 (bases 1 to 2148)
 AUTHORS van Baren,N., Chambost,H., Ferrant,A., Michaux,L., Ikeda
 30 Millard,I., Olive,D., Boon,T. and Coulie,P.G.
 TITLE PRAME, a gene encoding an antigen recognized on a
 human melanoma by
 cytolytic T cells, is expressed in acute leukaemia cells
- 35 JOURNAL Br. J. Haematol. 102 (5), 1376-1379 (1998)
 MEDLINE 98423996
 PUBMED 9753074
 REFERENCE 4 (bases 1 to 2148)
 AUTHORS Dunham I, Shimizu N, Roe BA, Chissole S, Hunt AR, Collins
 40 Bruskiewich R, Beare DM, Clamp M, Smink LJ, Ainscough
 R, Almeida
 JP, Babbage A, Bagguley C, Bailey J, Barlow K, Bates
 KN, Beasley O,
 Bird CP, Blakey S, Bridgeman AM, Buck D, Burgess J,
 Burrill WD,
- 45 O'Brien KP and et al.
 TITLE The DNA sequence of human chromosome 22
 JOURNAL Nature 402 (6761), 489-495 (1999)
 MEDLINE 20057165
 PUBMED 10591208
- 50 REMARK Erratum:[published erratum appears in Nature 2000 Apr
 20;404(6780):904]]
 COMMENT REVIEWED REFSEQ: This record has been curated by NCBI
 staff. The
 reference sequence was derived from U65011.1.
 55 Summary: The protein encoded by this gene has a 509
 amino acid
 antigen, lacking a signal sequence, and recognized on
 a human

melanoma cell line by a T-lymphocyte clone. A
significant level of
this mRNA is detected in normal testis as well as in
many
5 melanomas, non-small cell lung carcinomas, sarcomas,
head and neck
tumors and renal carcinomas. The encoded protein is
expressed
predominantly in acute leukemias carrying chromosomal
10 abnormalities
such as translocation t(8:21), which fuses the AML1
and ETO genes.
Its expression shares several characteristics with the
expression
15 patterns of MAGE, BAGE, and GAGE gene families, all of
which are
expressed in tumors. This protein is expressed in a
higher
proportion of samples than genes of the MAGE, BAGE,
20 and GAGE
families, and of these four groups, only this protein
is expressed
by acute myeloid leukemias.
COMPLETENESS: complete on the 3' end.
25 FEATURES Location/Qualifiers
source 1..2148
/organism="Homo sapiens"
/db_xref="taxon:9606"
/chromosome="22"
30 /map="22q11.22"
/haplotype="A24 A28 B13 B44 Cw6 Cw7"
gene 1..2148
/gene="PRAME"
/note="MAPE; OIP4"
35 /db_xref="LocusID:23532"
/db_xref="MIM:606021"
CDS 236..1765
/gene="PRAME"
/note="melanoma antigen preferentially
40 expressed in
tumors; Opa-interacting protein OIP4"
/codon_start=1
/db_xref="LocusID:23532"
/db_xref="MIM:606021"
45 /product="preferentially expressed antigen of
melanoma"
/protein_id="NP_006106.1"
/db_xref="GI:5174641"
50 /translation="MERRRLWGSIQSRYISMSVWTSPPRLVELAGQSLLKDEALATAALELLPREL
FPPLFMAAFDGRHSQTLKAMVQAWPFTCLPLGLVMKGQHLHLETFAVL DGLDVL LAQEVPRRWK
LQVL DLRKNSHQDFWTVWSGNRASLYSFPEPEAAQPM TKKRKVDGLSTEA EQPFIPVEVLVDLFLK
EGACDELFSY LIEKVRRKKNVLR LCCCKLKFAMP MQDIKMILKMVQLDSIEDLEV TCTWKLP TLA
KFSPYLGOMINLRRL LLSHIHASSYISPEKEEQYIAQFTSQFLSLQCLQALYVDSLFFLRGR L DQL
55 LRHVMNPLET L SITNCRLSEGDVMHLSQSPSVLSVLSL SGVMLTDVSPEPLQALLERASATLQD
LVFDECGITDDQL LALLPSLSHCSQLT T LSFYGN SISISALQSL LQHLIGLSNLTHVLYPVPLESY
BDIHGTLHLERLAYLHARLRELLCELGRPSMVWLSANPCPHCGDRTFYDPEILCPCFMPN"
variation 254

```

/allele="C"
/allele="T"
/db_xref="dbSNP:1129172"
5      variation      698
/allele="C"
/allele="A"
/db_xref="dbSNP:1063107"
      variation      1001
/allele="C"
10     /allele="T"
/db_xref="dbSNP:2229695"
      variation      1498
/allele="T"
/allele="C"
15     /db_xref="dbSNP:2229696"
      variation      1762
/allele="C"
/allele="T"
/db_xref="dbSNP:13604"
20     variation      1952
/allele="A"
/allele="G"
/db_xref="dbSNP:7104"
      variation      1984
25     /allele="A"
/allele="C"
/db_xref="dbSNP:8405"
      variation      2043
/allele="T"
/allele="A"
30     /db_xref="dbSNP:1804685"
polyA_signal      2108..2113
polyA_site        2130
BASE COUNT      534 a      548 c      558 g      508 t
ORIGIN
35      1 gcttcagggt acagctcccc cgcagccaga agccgggcct gcagccctc
agcaccgctc
      61 cgggacaccc caccgcttc ccaggcgtga cctgtcaaca gcaacttcgc
ggtgtggtga
40     121 actctctgag gaaaaaccat ttgattatt actctcagac gtgcgtggca
acaagtgact
      181 gagacctaga aatccaagcg ttggaggtcc tgaggccagc ctaagtcgct
tcaaaatgga
      241 acgaaggcgt ttgtgggggt ccattcagag ccgatacatc agcatgagtg
45     tgtggacaag
      301 cccacggaga cttgtggagc tggcagggca gagcctgctg aaggatgagg
ccctggccat
      361 tgccgccctg gagttgctgc ccaggagct ctccccgcca ctcttcattg
cagcctttga
50     421 cgggagacac agccagaccc tgaaggcaat ggtgcaggcc tggcccttca
cctgcctccc
      481 tctgggagtg ctgatgaagg gacaacatct tcacctggag accttcaaag
ctgtgcttga
      541 tggacttgat gtgctccttg ccaggaggt tcgccccagg aggtggaaac
55     ttcaagtgtc
      601 ggatttacgg aagaactctc atcaggactt ctggactgta tggctctggaa
acagggccag

```

661 tctgtactca tttccagagc cagaagcagc tcagcccatg acaaagaagc
gaaaagtaga
721 tggtttgagc acagaggcag agcagccctt cattccagta gaggtgctcg
tagacctggt
5 781 cctcaaggaa ggtgcctgtg atgaattggt ctccctacctc attgagaaaag
tgaagcgaaa
841 gaaaaatgta ctacgcctgt gctgtaagaa gctgaagatt tttgcaatgc
ccatgcagga
901 tatcaagatg atcctgaaaa tgggtgcagct ggactctatt gaagatttgg
10 aagtgcattg
961 tacctggaag ctaccacact tggcgaaatt ttctccttac ctgggccaga
tgattaatct
1021 gcgtagactc ctccctctccc acatccatgc atcttcctac atttccccgg
agaaggaaga
15 1081 gcagtatatc gccagttca cctctcagtt cctcagtcctg cagtgcctgc
aggctctcta
1141 tgtggactct ttatttttcc ttagaggccg cctggatcag ttgctcaggg
acgtgatgaa
1201 ccccttgga accctctcaa taactaactg ccggctttcg gaaggggatg
20 tgatgcacat
1261 gtcccagagt cccagcgtca gtcagctaag tgtcctgagt ctaagtgggg
tcatgctgac
1321 cgatgtaagt cccgagcccc tccaagctct gctggagaga gcctctgcca
ccctccagga
25 1381 cctggctctt gatgagtgtg ggatcacgga tgatcagctc cttgccctcc
tgccctccct
1441 gagccactgc tcccagctta caaccttaag cttctacggg aattccatct
ccatatctgc
1501 cttgcagagt ctccctgcagc acctcatcgg gctgagcaat ctgacccacg
30 tgctgatatc
1561 tgtccccctg gagagttatg aggacatcca tggtagcctc cacctggaga
ggcttgcccta
1621 tctgcatgcc aggtcaggg agttgctgtg tgagttgggg cggcccagca
tggctctggct
35 1681 tagtgccaac ccctgtcctc actgtgggga cagaaccttc tatgacccgg
agcccatcct
1741 gtgcccctgt ttcatgccta actagctggg tgcacatatc aaatgcttca
ttctgcatac
1801 ttggacacta aagccaggat gtgcatgcat cttgaagcaa caaagcagcc
40 acagtttcag
1861 acaaagtgtc agtgtgagtg agggaaacat gttcagtgag gaaaaaacat
tcagacaaat
1921 gttcagtgag gaaaaaaagg ggaagttggg gataggcaga tgttgacttg
aggagttaat
45 1981 gtgatctttg gggagataca tcttatagag ttagaaatag aatctgaatt
tctaaagggg
2041 gattctggct tgggaagtac atgtaggagt taatccctgt gtagactggt
gtaaagaaac
2101 tgttgaaaat aaagagaagc aatgtgaagc aaaaaaaaaa aaaaaaaaaa

ED-B domain of Fibronectin

LOCUS HSFIBEDB 2823 bp DNA linear
PRI 09-AUG-1999

5 DEFINITION Human fibronectin gene ED-B region.
ACCESSION X07717
VERSION X07717.1 GI:31406
KEYWORDS alternate splicing; fibronectin.
SOURCE human.

10 ORGANISM Homo sapiens
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
Euteleostomi;
Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
Homo.

15 REFERENCE 1 (bases 1 to 2823)
AUTHORS Paoletta,G., Henchcliffe,C., Sebastio,G. and
Baralle,F.E.
TITLE Sequence analysis and *in vivo* expression show that
alternative
20 splicing of ED-B and ED-A regions of the human
fibronectin gene are
independent events
JOURNAL Nucleic Acids Res. 16 (8), 3545-3557 (1988)
MEDLINE 88233940

25 FEATURES Location/Qualifiers
source 1..2823
/organism="Homo sapiens"
/db_xref="taxon:9606"
/clone="MA10"

30 exon 1..104
/number=1
/product="fibronectin"

CDS join(<2..104,1375..1647,2758..>2823)
/codon_start=1
35 /product="fibronectin"
/protein_id="CAB52437.1"
/db_xref="GI:5725425"

/translation="CTFDNLSPGLEYNVSVYTVKDDKESVPISDTIIEVPQLTDLSE"

VDITDSSIGLRWTPPLNSSTIIGYRITVVAAGEGIPFEDFVDSSVGYYTVTGLEPGID

YDISVITLINGGESAPTTLTQQTAVPPPTDLRFTNIGPDTMRVTW"

5 intron 105..1374
 /number=1
 exon 1375..1647
 /note="ED-B exon"
 /number=2
10 /product="fibronectin"
 intron 1648..2757
 /number=2
 exon 2758..2823
 /number=3
15 /product="fibronectin"
BASE COUNT 824 a 556 c 528 g 915 t
ORIGIN
 1 ctgcactttt gataacctga gtcccggcct ggagtacaat gtcagtgttt
acactgtcaa
20 61 ggatgacaag gaaagtgtcc ctatctctga taccatcatc ccaggtaata
gaaaataagc
 121 tgctatcctg agagtgacat tccaataaga gtggggatta gcatcttaat
ccccagatgc
 181 ttaaggggtg caactatatt tgggatttaa ttccgatctc ccagctgcac
25 tttccaaaac
 241 caagaagtca aagcagcgat ttggacaaaa tgcttgctgt taacactgct
ttactgtctg
 301 tgcttcactg ggatgctgtg tgttgcagcg agtatgtaat ggagtggcag
ccatggcttt
30 361 aactctgtat tgtctgctca catggaagta tgactaaaac actgtcacgt
gtctgtactc
 421 agtactgata ggctcaaagt aatatggtaa atgcatccca tcagtacatt
tctgcccgat
 481 ttacaatcc atatcaattt ccaacagctg cctatttcat cttgcagttt
35 caaatccttc
 541 tttttgaaaa ttggatttta aaaaaaagtt aagtaaaagt cacaccttca
gggttgttct
 601 ttcttggtggc cttgaaagac aacattgcaa aggctgtcc taaggatagg
cttgtttgtc

661 cattgggtta taacataatg aaagcattgg acagatcgtg tccccctttg
gactcttcag
721 tagaatgctt ttactaacgc taattacatg ttttgattat gaatgaacct
aaaatagtgg
5 781 caatggcctt aacctaggcc tgtctttcct cagcctgaat gtgcttttga
atggcacatt
841 tcacaccata cattcataat gcattagcgt tatggccatg atgttgtcat
gagttttgta
901 tgggagaaaa aaaatcaatt tatcacccat ttattatttt ttccggttgt
10 tcatgcaagc
961 ttattttcta ctaaaacagt tttggaatta ttaaaagcat tgctgatact
tacttcagat
1021 attatgtcta ggctctaaga atgggttcga catcctaaac agccatatga
tttttaggaa
15 1081 tctgaacagt tcaaattgta ccttttaagg atgttttcaa aatgtaaaaa
atatatatat
1141 atatatatat tccctaaaag aatattcctg tttattcttc tagggaagca
aactgttcat
1201 gatgcttagg aagtcttttc agagaattta aaacagattg catattacca
20 tcattgcttt
1261 aacattccac caattttact actagtaacc tgatatacac tgcctttattt
tttctctttt
1321 ttttccctct attttccttt tgccctcccc tccctttgct ttgtaactca
atagaggtgc
25 1381 cccaactcac tgacctaagc tttgttgata taaccgattc aagcatcggc
ctgaggtgga
1441 ccccgctaaa ctcttcacc attattgggt accgcatcac agtagttgcg
gcaggagaag
1501 gtatccctat ttttgaagat tttgtggact cctcagtagg atactacaca
30 gtcacagggc
1561 tggagccggg cattgactat gatatcagcg ttatcactct cattaatggc
ggcgagagtg
1621 cccctactac actgacacaa caaacgggtg aattttgaaa acttctgcgt
ttgagacata
35 1681 gatggtgttg catgctgcc aagttactc cggttaaata tggatgttcc
atgggggaag
1741 tcagcaattg gccaaagatt cagataggtg gaattggggg gataaggaat
caaatgcac

1801 tgctaaactg attggagaaa aacacatgca atatcttcag tacactctca
tttaaaccac
1861 aagtagatat aaagcctaga gaaatacaga tgtctgctct gttaaataa
aaatagcaaa
5 1921 tggtcattca atttgaagac ctagaatttt tcttcttaaa taccaaacac
gaataccaaa
1981 ttgcgtaagt accaattgat aagaatatat caccaaaatg taccatcatg
ctcttccttc
2041 taccctttga taaactctac catgctcctt ctttgtagct aaaaacccat
10 caaaatttag
2101 ggtagagtgg atgggcattg ttttgaggta ggagaaaagt aaacttgga
ccattctagg
2161 ttttggtgct gtcactagggt aaagaaacac ctctttaacc acagtctggg
gacaagcatg
15 2221 caacatttta aaggttctct gctgtgcatg ggaaaagaaa catgctgaga
accaatttgc
2281 atgaacatgt tcacttgtaa gtagaattca ctgaatggaa ctgtagctct
agatatctca
2341 catgggggga agtttaggac cctcttgtct tttgtctgt gtgcatgtat
20 ttctttgtaa
2401 agtactgcta tgttctcttt tgetgtgtgg caacttaagc ctcttcggcc
tgggataaaa
2461 taatctgcag tggattaat aatgtacata aagtcaacat atttgaaagt
agattaaaat
25 2521 ctttttttaa tatatcaatg atggcaaaaa ggtaaagggt ggctaacag
tactgtgtgt
2581 agtggtttat ttttaacagt agtacactat aacttaaaat agacttagat
tagactgttt
2641 gcatgattat gattctgttt cctttatgca tgaaatattg attttacctt
30 tccagctact
2701 tcgtagctt taatttttaa atacattaac tgagtcttcc ttctgttcg
aaaccagctg
2761 ttctctctcc cactgacctg cgattcacca acattggtcc agacaccatg
cgtgtcacct
35 2821 ggg
//

CEA
LOCUS NM_004363 2974 bp mRNA linear
PRI 28-NOV-2000
DEFINITION Homo sapiens carcinoembryonic antigen-related cell
5 adhesion
molecule 5 (CEACAM5), mRNA.
ACCESSION NM_004363
VERSION NM_004363.1 GI:11386170
KEYWORDS .
10 SOURCE human.
ORGANISM Homo sapiens
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
Euteleostomi;
Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
15 Homo.
REFERENCE 1 (bases 1 to 2974)
AUTHORS Oikawa S, Nakazato H and Kosaki G.
TITLE Primary structure of human carcinoembryonic antigen
(CEA) deduced
20 from cDNA sequence
JOURNAL Biochem. Biophys. Res. Commun. 142 (2), 511-518 (1987)
MEDLINE 87128144
PUBMED 3814146
REFERENCE 2 (bases 1 to 2974)
25 AUTHORS Zimmermann W, Weber B, Ortlieb B, Rudert F, Schempp W,
Fiebig HH,
Shively JE, von Kleist S and Thompson JA.
TITLE Chromosomal localization of the carcinoembryonic
antigen gene
30 family and differential expression in various tumors
JOURNAL Cancer Res. 48 (9), 2550-2554 (1988)
MEDLINE 88184584
PUBMED 3356015
REFERENCE 3 (bases 1 to 2974)
35 AUTHORS Barnett, T., Goebel, S.J., Nothdurft, M.A. and
Elting, J.J.
TITLE Carcinoembryonic antigen family: characterization of
cDNAs coding

for NCA and CEA and suggestion of nonrandom sequence
variation in
their conserved loop-domains
JOURNAL Genomics 3 (1), 59-66 (1988)
5 MEDLINE 89122014
REFERENCE 4 (bases 1 to 2974)
AUTHORS Barnett T and Zimmermann W.
TITLE Workshop report: proposed nomenclature for the
carcinoembryonic
10 antigen (CEA) gene family
JOURNAL Tumour Biol. 11 (1-2), 59-63 (1990)
MEDLINE 90176333
PUBMED 2309067
REFERENCE 5 (bases 1 to 2974)
15 AUTHORS Schrewe H, Thompson J, Bona M, Hefta LJ, Maruya A,
Hassauer M,
Shively JE, von Kleist S and Zimmermann W.
TITLE Cloning of the complete gene for carcinoembryonic
antigen: analysis
20 of its promoter indicates a region conveying cell
type-specific
expression
JOURNAL Mol. Cell. Biol. 10 (6), 2738-2748 (1990)
MEDLINE 90258861
25 PUBMED 2342461
COMMENT PROVISIONAL REFSEQ: This record has not yet been
subject to final
NCBI review. The reference sequence was derived from
M29540.1.
30 FEATURES Location/Qualifiers
source 1..2974
/organism="Homo sapiens"
/db_xref="taxon:9606"
/chromosome="19"
35 /map="19q13.1-q13.2"
gene 1..2974
/gene="CEACAM5"
/note="CEA; CD66e"
/db_xref="LocusID:1048"

1 ctcagggcag agggaggaag gacagcagac cagacagtca cagcagcctt
gacaaaacgt
61 tcctggaact caagctcttc tccacagagg aggacagagc agacagcaga
gaccatggag
5 121 tctccctcgg cccctcccca cagatgggtgc atcccctggc agaggctcct
gtccacagcc
181 tcacttctaa ccttctggaa cccgcccacc actgccaagc tcactattga
atccacgcgc
241 ttcaatgtcg cagaggggaa ggaggtgctt ctacttgtcc acaatctgcc
10 ccagcatctt
301 tttggctaca gctggtacaa aggtgaaaga gtggatggca accgtcaa
tataggatat
361 gtaataggaa ctcaacaagc taccacaggg cccgcataca gtggctcgaga
gataatatac
15 421 cccaatgcat ccctgctgat ccagaacatc atccagaatg acacaggatt
ctacacccta
481 cacgtcataa agtcagatct tgtgaatgaa gaagcaactg gccagttccg
ggtatacccg
541 gagctgcccc agccctccat ctceagcaac aactccaaac ccgtggagga
20 caaggatgct
601 gtggccttca cctgtgaacc tgagactcag gacgcaacct acctgtggtg
ggtaaacaat
661 cagagcctcc cggtcagtcc caggctgcag ctgtccaatg gcaacaggac
cctcactcta
25 721 ttcaatgtca caagaaatga cacagcaagc taaaaatgtg aaaccagaa
cccagtgagt
781 gccaggcgca gtgattcagt catcctgaat gtccctctatg gcccggatgc
ccccaccatt
841 tccccctctaa acacatctta cagatcaggg gaaaatctga acctctcctg
30 ccacgcagcc
901 tctaaccacac ctgcacagta ctcttggttt gtcaatggga ctttccagca
atccacccaa
961 gagctcttta tccccaacat cactgtgaat aatagtggat cctatacgtg
ccaagcccat
35 1021 aactcagaca ctggcctcaa taggaccaca gtcacgacga tcacagtcta
tgcagagcca
1081 cccaaaccct tcatcaccag caacaactcc aaccccgtagg aggatgagga
tgctgtagcc

1141 ttaacctgtg aacctgagat tcagaacaca acctacctgt ggtgggtaaa
 taatcagagc
 1201 ctcccggtea gtcccaggct gcagctgtcc aatgacaaca ggaccctcac
 tctactcagt
 5 1261 gtcacaagga atgatgtagg acctatgag tgtggaatcc agaacgaatt
 aagtgttgac
 1321 cacagcgacc cagtcaccc cagtcaccc tatggcccag acgacccccc
 catttcccc
 1381 tcatacacct attaccgtcc aggggtgaac ctcagccctc cctgccatgc
 10 agcctctaac
 1441 ccacctgcac agtattcttg gctgattgat gggaacatcc agcaacacac
 acaagagctc
 1501 tttatctcca acatcaatga gaagaacagc ggactctata cctgccaggc
 caataactca
 1561 gccagtggcc acagcaggac tacagtcaag acaatcacag tctctgcgga
 gctgcccagg
 1621 ccctccatct ccagcaacaa ctccaaaccc gtggaggaca aggatgctgt
 ggccctcacc
 1681 tgtgaacctg aggctcagaa cacaacctac ctgtggtggg taaatgggtca
 20 gagcctccca
 1741 gtcagtccca ggctgcagct gtccaatggc aacaggaccc tcactctatt
 caatgtcaca
 1801 agaaatgacg caagagccta tgtatgtgga atccagaact cagtgagtgc
 aaaccgcagt
 25 1861 gaccagtcac ccctggatgt cctctatggg ccggacaccc ccatcatttc
 cccccagac
 1921 tcgtcttacc ttccgggagc gaacctcaac ctctcctgcc actcggcctc
 taaccatcc
 1981 ccgcagtatt cttggcgat caatgggata ccgcagcaac acacacaagt
 30 tctctttatc
 2041 gccaaaatca cgccaaataa taacgggacc tatgcctgtt ttgtctctaa
 cttggctact
 2101 ggccgcaata attccatagt caagagcatc acagtctctg catctggaac
 ttctcctggt
 35 2161 ctctcagctg gggccactgt cggcatcatg attggagtgc tgggtggggg
 tgctctgata
 2221 tagcagccct ggtgtagttt cttcatttca ggaagactga cagttgtttt
 gcttcttctc

2281 taaagcattt gcaacagcta cagtctaaaa ttgcttcttt accaaggata
 ttacagaaa
 2341 agactctgac cagagatcga gaccatccta gccaacatcg tgaaaccca
 tctctactaa
 5 2401 aaatacaaaa atgagctggg cttgggtggcg cgcacctgta gtcccagtta
 ctcgaggaggc
 2461 tgaggcagga gaatcgcttg aaccggggag gtggagattg cagtgagccc
 agatcgcacc
 2521 actgcactcc agtctggcaa cagagcaaga ctccatctca aaaagaaaag
 10 aaaagaagac
 2581 tctgacctgt actcttgaat acaagtttct gataccactg cactgtctga
 gaatttccaa
 2641 aactttaatg aactaactga cagcttcatg aaactgtcca ccaagatcaa
 gcagagaaaa
 15 2701 taattaattt catgggacta aatgaactaa tgaggattgc tgattcttta
 aatgtcttgt
 2761 ttcccagatt tcaggaaact ttttttcttt taagctatcc actcttacag
 caatttgata
 2821 aaatatactt ttgtgaacaa aaattgagac atttacattt tctccctatg
 20 tggtcgctcc
 2881 agacttggga aactattcat gaatatttat attgtatggt aatatagtta
 ttgcacaagt
 2941 tcaataaaaa tctgctcttt gtataacaga aaaa
 //

25

Her2/Neu

LOCUS HUMHER2A 4530 bp mRNA linear
 PRI 18-SEP-1995
 30 DEFINITION Human tyrosine kinase-type receptor (HER2) mRNA,
 complete cds.
 ACCESSION M11730
 VERSION M11730.1 GI:183986
 KEYWORDS tyrosine kinase.
 35 SOURCE Homo sapiens (clone: lambda-HER2-436) fetal cDNA to
 mRNA.
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
 Euteleostomi;

Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
Homo.
REFERENCE 1 (bases 1 to 4530)
AUTHORS Coussens, L., Yang-Feng, T.L., Liao, Y.-C., Chen, E.,
5 Gray, A.,
McGrath, J., Seeburg, P.H., Libermann, T.A.,
Schlessinger, J.,
Francke, U., Levinson, A. and Ullrich, A.
TITLE Tyrosine kinase receptor with extensive homology to
10 EGF receptor
shares chromosomal location with neu oncogene
JOURNAL Science 230 (4730), 1132-1139 (1985)
MEDLINE 86070181
REFERENCE 2 (bases 1701 to 1719)
15 AUTHORS Ullrich, A.
JOURNAL Unpublished (1988)
FEATURES Location/Qualifiers
source 1..4530
/organism="Homo sapiens"
20 /db_xref="taxon:9606"
/clone="lambda-HER2-436"
/dev_stage="fetal"
mRNA <1..4530
/product="HER2 mRNA"
25 CDS 151..3918
/note="HER2 receptor"
/codon_start=1
/protein_id="AAA75493.1"
/db_xref="GI:306840"
30
/translation="MELAALCRWGLLLALLPPGAASTQVCTGTDMKLRLPASPETHLD
MLRHLYQGCQVVQGNLELTYPNTNASLSFLQDIQEVQGYVLIHNSQVRQVPLQRLRIV
RGTQLFEDNYALAVLDNGDPLNNTTPVTGASPGGLRELQLRSLTEILKGGVLIQRNPQ
LCYQDTILWKDIFHKNNQLALTIDTNRSRACHPCSPMCKGSRWGESSEDCQSLTRT
35 VCAGGCARCKGPLPTDCCHEQCAAGCTGPKHSDCLACLFNHSGICELHCPALVTYNT
DTFESMPNPEGRYTFGASCVTACPYNYLSTDVGSCTLVCPLHNQEVTAE DGTQRCEKC
SKPCARVCYGLGMEHLREVRAVTSANIQEFAGCKKIFGSLAFLPESFDGDPASNTAPL
QPEQLQVFETLEEITGYLYISAWPDSLPLSVFQNLQVIRGRILHNGAYSLTLQGLGI
SWLGLRSLRELGSGLALIHNTLHCFVHTVPWDQLFRNPHQALLHTANRPEDECVGEG

LACHQLCARGHCWGPPTQCVNCSQFLRGQECVEECRVLQGLPREYVNARHCLPCHPE
 CQFQNGSVTCFGPEADQCVACAHYKDPFPCVARCPGSKPDLSPYMPIWKFPDEEGACQ
 PCPINCTHSCVDLDDKGCPAEQRASPLTSIVSAVVGILLVVVLGVVFGILIKRRQQKI
 RKYTMRRLLQETELVEPLTPSGAMPNQAQMRIKETELRKVKVLGSGAFGTVYKGIWI
 5 PDGENVKIPVAIKVLRENTSPKANKEILDEAYVMAGVGSPYVSRLLGICLTSTVQLVT
 QIMPYGCLLDHVRENRLGSLQDLLNWCMIAGMSYLEDVRLVHRDLAARNVLVKSP
 NHVKITDFGLARLLDIDETEHADGGKVPIKWMALLESILRRRFTHQSDVWSYGVTVWE
 LMTFGAKPYDGIPIAREIPDLLEKGERLPQPPICTIDVYMIMVKCWMIDSECRPRFREL
 VSEFSRMARDPQRFVVIQNEGLGPASPLDSTFYRSLLLEDDMDGLVDAEYLVLPQQGF
 10 FCPDPAPGAGGMVHHRSSSTRSGGDLTLGLEPSEEEAPRSPLAPSEGAGSDVFDG
 DLGMAAKGLQSLPTHDPSPQLQRYSEDPTVPLPSETDGYVAPLTCSPQPEYVNQPDVR
 PQPPSPREGPLPAARPAGATLERAKTLPSPGKNGVVKDVFAFGGAVENPEYLTPOGGAA
 PQPHPPPAFSPAFDNLYYWDQDPPERGAPPSTFKGTPTAENPEYLGLDVPV"

old_sequence 1701..1719

15

/citation=[1]

BASE COUNT 922 a 1382 c 1346 g 880 t

ORIGIN . Chromosome 17q21-q22.

1 aattctcgag ctctctgacc ggtcgacgag ctcgagggtc gacgagctcg
 agggcgcgcg
 20 61 cccggccccc acccctcgca gcaccccgcg ccccgcgccc tcccagccgg
 gtccagccgg
 121 agccatgggg ccggagccgc agtgagcacc atggagctgg cggccttggtg
 ccgctggggg
 181 ctctctctcg ccctcttgcc ccccgagacc gcgagcacc aagtgtgcac
 25 cggcacagac
 241 atgaagctgc ggctccctgc cagtcccgag acccacctgg acatgctccg
 ccacctctac
 301 cagggtgcc aggtggtgca gggaaacctg gaactcacct acctgcccac
 caatgccagc
 361 ctgtccttcc tgcaggatat ccaggaggtg cagggtctacg tgctcatcgc
 tcacaaccaa
 421 gtgaggcagg tcccactgca gaggtctcgg attgtgcgag gcacccagct
 ctttgaggac
 481 aactatgcc tggccgtgct agacaatgga gaccgctga acaataccac
 35 ccctgtcaca
 541 ggggcctccc caggaggcct gcgggagctg cagcttcgaa gcctcacaga
 gatcttgaaa
 601 ggaggggtct tgatccagcg gaacccccag ctctgctacc aggacacgat
 tttgtggaag

661 gacatcttcc acaagaacaa ccagctggct ctcacactga tagacaccaa
ccgctctcgg
721 gcctgccacc cctgttctcc gatgtgtaag ggctcccgt gctggggaga
gagttctgag
5 781 gattgtcaga gcctgacgcg cactgtctgt gccggtggct gtgcccgctg
caagggggcca
841 ctgcccactg actgctgcca tgagcagtgt gctgccggct gcacggggccc
caagcactct
901 gactgcctgg cctgcctcca cttcaaccac agtggcatct gtgagctgca
10 ctgcccagcc
961 ctggtcacct acaacacaga cacgtttgag tccatgcccc atcccagagg
ccggtatata
1021 ttccggcgcca gctgtgtgac tgctgtccc tacaactacc tttctacgga
cgtgggatcc
15 1081 tgcaccctcg tctgccccct gcacaaccaa gaggtgacag cagaggatgg
aacacagcgg
1141 tgtgagaagt gcagcaagcc ctgtgcccga gtgtgctatg gtctgggcat
ggagcacttg
1201 cgagaggtga gggcagttac cagtgccaat atccaggagt ttgctggctg
20 caagaagatc
1261 tttgggagcc tggcatttct gccggagagc tttgatgggg acccagcctc
caacactgcc
1321 ccgctccagc cagagcagct ccaagtgttt gagactctgg aagagatcac
aggttaccta
25 1381 tacatctcag catggccgga cagcctgcct gacctcagcg tcttcagaa
cctgcaagta
1441 atccggggac gaattctgca caatggcgcc tactcgctga ccctgcaagg
gctgggcatc
1501 agctggctgg ggctgcgctc actgaggga ctgggcagt gactggccct
30 catccaccat
1561 aacaccacc tctgcttcgt gcacacggtg ccctgggacc agctctttcg
gaaccgcac
1621 caagctctgc tccacactgc caaccggcca gaggacgagt gtgtgggcga
gggcctggcc
35 1681 tgccaccagc tgtgcgccc agggcactgc tggggtccag ggccaccca
gtgtgtcaac
1741 tgcagccagt tccttcgggg ccaggagtgc gtggaggaat gccgagtact
gcaggggctc

1801 cccagggagt atgtgaatgc caggcactgt ttgccgtgcc accctgagtg
tcagccccag
1861 aatggctcag tgacctgttt tggaccggag gctgaccagt gtgtggcctg
tgcccactat
5 1921 aaggaccctc ccttctgcgt ggcccgtgc cccagcggtg tgaaacctga
cctctcctac
1981 atgcccattct ggaagtttcc agatgaggag ggcgcattgcc agccttgccc
catcaactgc
2041 acccactcct gtgtggacct ggatgacaag ggctgccccg ccgagcagag
10 agccagccct
2101 ctgacgtcca tcgtctctgc ggtggttggc attctgctgg tcgtggtctt
gggggtggtc
2161 tttgggatcc tcatcaagcg acggcagcag aagatccgga agtacacgat
gcggagactg
15 2221 ctgcaggaaa cggagctggg ggagccgctg acacctagcg gagcgatgcc
caaccaggcg
2281 cagatgcgga tcctgaaaga gacggagctg aggaaggtga aggtgcttgg
atctggcgt
2341 tttggcacag tctacaaggg catctggatc cctgatgggg agaattgtgaa
20 aattccagtg
2401 gccatcaaag tgttgaggga aaacacatcc cccaaagcca acaaagaaat
cttagacgaa
2461 gcatacgtga tggctgggtg gggctcccca tatgtctccc gccttctggg
catctgcctg
25 2521 acatccacgg tgcagctggg gacacagctt atgcctatg gctgcctctt
agaccatgtc
2581 cgggaaaacc gcggacgcct gggctcccag gacctgctga actggtgtat
gcagattgcc
2641 aaggggatga gctacctgga ggatgtgcgg ctcgtaaca gggacttggc
30 cgctcggaac
2701 gtgctggtca agagtcccaa ccatgtcaaa attacagact tcgggctggc
tcggctgctg
2761 gacattgacg agacagagta ccatgcagat gggggcaagg tgcccatcaa
gtggatggcg
35 2821 ctggagtcca ttctccgccg gcggttcacc caccagagtg atgtgtggag
ttatgggtgtg
2881 actgtgtggg agctgatgac ttttggggcc aaaccttacg atgggatccc
agcccgggag

2941 atccctgacc tgctggaaaa gggggagcgg ctgccccagc ccccatctg
caccattgat
3001 gtctacatga tcatgggtcaa atgttgatg attgactctg aatgtcggcc
aagattccgg
5 3061 gagttggtgt ctgaattctc ccgcatggcc agggaccccc agcgctttgt
ggtcateccag
3121 aatgaggact tggggccagc cagtcccttg gacagcacct tctaccgctc
actgctggag
3181 gacgatgaca tgggggacct ggtggatgct gaggagtatc tggtagccca
10 gcagggcttc
3241 ttctgtccag accctgcccc gggcgctggg ggcattggtcc accacaggca
ccgcagctca
3301 tctaccagga gtggcggtgg ggacctgaca ctagggtctg agccctctga
agaggaggcc
15 3361 cccaggctctc cactggcacc ctccgaaggg gctgggtccg atgtatttga
tggtgacctg
3421 ggaatggggg cagccaaggg gctgcaaagc ctccccacac atgacccag
ccctctacag
3481 cggtagagtg aggacccac agtaccctg ccctctgaga ctgatggcta
20 cgttgccccc
3541 ctgacctgca gccccagcc tgaatatgtg aaccagccag atgttcggcc
ccagccccct
3601 tcgccccgag agggccctct gcctgctgcc cgacctgctg gtgccactct
ggaaagggcc
25 3661 aagactctct cccagggaa gaatggggtc gtcaaagacg tttttgcctt
tgggggtgcc
3721 gtggagaacc ccgagtactt gacacccag ggaggagctg cccctcagcc
ccaccctcct
3781 cctgccttca gccagcctt cgacaacctc tattactggg accaggacct
30 accagagcgg
3841 ggggctccac ccagcacctt caaagggaca cctacggcag agaaccaga
gtacctgggt
3901 ctggacgtgc cagtgtgaac cagaaggcca agtccgcaga agccctgatg
tgtcctcagg
35 3961 gagcagggaa ggcctgactt ctgctggcat caagaggtgg gagggccctc
cgaccacttc
4021 caggggaacc tgccatgcc aagacctgtc ctaaggaacc ttccctcctg
cttgagttcc

```

      4081 cagatggctg gaaggggtcc agcctcgttg gaagaggaac agcactggg
      agtctttgtg
      4141 gattctgagg ccctgccccaa tgagactcta ggggccagtg gatgccacag
      cccagcttgg
5      4201 ccctttcctt ccagatcctg ggtactgaaa gccttaggga agctggcctg
      agaggggaag
      4261 cggccctaag ggagtgtcta agaacaaaag cgacccattc agagactgtc
      cctgaaacct
      4321 agtactgccc cccatgagga aggaacagca atggtgtcag tatccaggct
10     ttgtacagag
      4381 tgcttttctg tttagttttt actttttttg ttttgttttt ttaaagacga
      aataaagacc
      4441 caggggagaa tgggtgttgt atggggaggc aagtgtgggg ggtccttctc
      cacaccact
15     4501 ttgtccattt gcaaataatat tttgaaaaac
      //

```

```

SCP-1
20 LOCUS      HSSCP1PRT      3393 bp      mRNA      linear
      PRI 26-FEB-1997
      DEFINITION H.sapiens mRNA for SCP1 protein.
      ACCESSION X95654
      VERSION X95654.1 GI:1212982
25 KEYWORDS SCP1 gene.
      SOURCE human.
      ORGANISM Homo sapiens
      Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
      Euteleostomi;
30 Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
      Homo.
      REFERENCE 1 (bases 1 to 3393)
      AUTHORS Meuwissen,R.L., Meerts,I., Hoovers,J.M., Leschot,N.J.
      and
35 Heyting,C.
      TITLE Human synaptonemal complex protein 1 (SCP1): isolation
      and
      characterization of the cDNA and chromosomal
      localization of the

```

gene
JOURNAL Genomics 39 (3), 377-384 (1997)
MEDLINE 97224467
REFERENCE 2 (bases 1 to 3393)
5 AUTHORS Meuwissen,R.J.L.
TITLE Direct Submission
JOURNAL Submitted (13-FEB-1996) Dr. R.L.J. Meuwissen,
Agricultural
University, Genetics, Dreijenlaan 2, 6703 HA
10 WAGENINGEN,
NETHERLANDS
FEATURES Location/Qualifiers
source 1..3393
/organism="Homo sapiens"
15 /db_xref="taxon:9606"
/map="1p12-p13"
/tissue_type="testis"
gene 95..3025
/gene="SCP1"
20 CDS 95..3025
/gene="SCP1"
/codon_start=1
/evidence=experimental
/product="polypeptide of 976 aa"
25 /protein_id="CAA64956.1"
/db_xref="GI:1212983"
/db_xref="SWISS-PROT:Q15431"

/translation="MEKQKPFALFVPPRSSSSQVSAVKPQTLGGDSTFFKSFNKCTED
30 DLEFPFAKTNLSKNGENIDSDPALQKVNFLPVLEQVGNSDCHYQEGDKDSDLENSEGL
SRVFSKLYKEAEKIKKWKVSTAEALRQKESKLQENRKII EAQRKAIQELQFGNEKVS
KLEEGIQENKDLIKENNATRHLCNLLKETCARSAEKTKKYBYEREETRQVYMDLNNNI
EKMITAHGELRVQAENSRLMHFKLKEDYEKIQHLEQYKKEINDKEKQVSLLLIQIT
EKENKMKDLTFLLEESRDKNVQLLEKTKLQSENKQSI EKQHHLTKLEDIKVSLQRS
35 VSTQKALEEDLQIATKTI COLTEEKETQMEESNKARAAHSFVVTEFETTVCSLEELLR
TEQQRLEKNEDQLKIILTMELQKKSSELEEMTKLTNNKEVELEELKKVLGEKETLLYEN
KQFEKIAEELKGTEQELIGLLQAREKEVHDLEIQLTAITTSQYYSKEVKDLKTELEN
EKLKNTELTSHCNKLSLENKELTQETSDMTLELKNQQEDINNNKKQBERMLKQIENLQ
ETETQLRNELEYVREELKQKRDEVKCKLKDSEENCNNLRKQVENKNKYIEELQQENKA

LKKKGTAESKQLNVYEIKVKNLELELES AKQKFGEITDTYQKEIEDKKISEENLLEEV
 EKAKVIAD EAVKLQKEIDKRCQH K I AEMVALMEKHKHQYDKI IEERDSELGLYKSKEQ
 EQSSLRASLEIELSNLKAELLSVKKQLEIEREEKEKLKREAKENTATLKEKKDKKTQT
 FLLETPEIYWKLD SKAVPSQTVSRNFTSVDHGISKDKRDYLWTS AKNTLSTPLPKAYT
 5 VKTPTKPKLQORENLNIP I EESKKKRKMAFEFDINS DSSETTDLLSMVSEETLKTLY
 RNNNPPASHLCVKTPKKAPSSLTTPGPTLKFGAIRKMREDRWAVIAKMDRKKKLKEAE
 KLFV"
 BASE COUNT 1440 a 483 c 648 g 822 t
 ORIGIN
 10 1 gccctcatag accgtttggt gtagttcgcg tgggaacagc aaccacggt
 ttcccgatag
 61 ttcttcaaag atatttaca ccgtaacaga gaaaatggaa aagcaaaagc
 cctttgcatt
 121 gtctgtacca ccgagatcaa gcagcagtca ggtgtctgcg gtgaaacctc
 15 agacctggg
 181 aggcgattcc actttcttca agagtttcaa caaatgtact gaagatgatt
 tggagtttcc
 241 atttgcaaag actaatctct ccaaaaatgg ggaaaacatt gattcagatc
 ctgctttaca
 20 301 aaaagttaat ttcttgcccg tgcttgagca ggttggaat tctgactgtc
 actatcagga
 361 aggactaaaa gactctgatt tggagaattc agagggattg agcagagtgt
 tttcaaaact
 421 gtataaggag gctgaaaaga taaaaaatg gaaagtaagt acagaagctg
 25 aactgagaca
 481 gaaagaaagt aagttgcaag aaaacagaaa gataattgaa gcacagcgaa
 aagccattca
 541 ggaactgcaa tttggaaatg aaaaagtaag tttgaaatta gaagaaggaa
 tacaagaaaa
 30 601 taaagattta ataaaagaga ataatgccac aaggcattta tgtaatctac
 tcaaagaaac
 661 ctgtgctaga tctgcagaaa agacaaagaa atatgaatat gaacgggaag
 aaaccaggca
 721 agtttatatg gatctaaata ataacattga gaaaatgata acagctcatg
 35 gggaacttcg
 781 tgtgcaagct gagaattcca gactggaaat gcattttaag ttaaaggaag
 attatgaaaa
 841 aatccaacac cttgaacaag aatacaagaa ggaaataaat gacaaggaaa
 agcaggtatc

901 actactattg atccaaatca ctgagaaaga aaataaaatg aaagatttaa
catttctgct
961 agaggaatcc agagataaag ttaatcaatt agaggaaaag acaaaattac
agagtgaaaa
5 1021 cttaaaacaa tcaattgaga aacagcatca tttgactaaa gaactagaag
atattaaagt
1081 gtcattacaa agaagtgtga gtactcaaaa ggcttttagag gaagatttac
agatagcaac
1141 aaaaacaatt tgtcagctaa ctgaagaaaa agaaactcaa atggaagaat
10 ctaataaagc
1201 tagagctgct cattcgtttg tggttactga atttgaaact actgtctgca
gcttggaaga
1261 attattgaga acagaacagc aaagattgga aaaaaatgaa gatcaattga
aaatacttac
15 1321 catggagctt caaaagaaat caagtgaact ggaagagatg actaagctta
caaataacaa
1381 agaagtagaa cttgaagaat tgaaaaaagt cttgggagaa aaggaaacac
ttttatatga
1441 aaataaaciaa tttgagaaga ttgctgaaga attaaaagga acagaacaag
20 aactaattgg
1501 tcttctccaa gccagagaga aagaagtaca tgatttgga atacagttaa
ctgccattac
1561 cacaagtga cagtattatt caaaagaggt taaagatcta aaaactgagc
ttgaaaacga
25 1621 gaagcttaag aatactgaat taacttcaca ctgcaacaag ctttcactag
aaaacaaaga
1681 gctcacacag gaaacaagt atatgaccct agaactcaag aatcagcaag
aagatattaa
1741 taataacaaa aagcaagaag aaaggatgtt gaaacaaata gaaaatcttc
30 aagaaacaga
1801 aaccaatta agaatgaac tagaatatgt gagagaagag ctaaaacaga
aaagagatga
1861 agttaaatgt aaattggaca agagtgaaga aaattgtaac aatttaagga
aacaagttga
35 1921 aaataaaaaac aagtatattg aagaacttca gcaggagaat aaggccttga
aaaaaaaagg
1981 tacagcagaa agcaagcaac tgaatgttta tgagataaag gtcaataaat
tagagttaga

2041 actagaaagt gccaaacaga aatttggaga aatcacagac acctatcaga
aagaaattga
2101 ggacaaaaag atatcagaag aaaatctttt ggaagagggt gagaaagcaa
aagtaatagc
5 2161 tgatgaagca gtaaaattac agaaagaaat tgataagcga tgtcaacata
aaatagctga
2221 aatggtagca cttatggaaa aacataagca ccaatatgat aagatcattg
aagaaagaga
2281 ctcagaatta ggactttata agagcaaaga acaagaacag tcatcactga
10 gagcatcttt
2341 ggagattgaa ctatccaatc tcaaagctga acttttgtct gttaagaagc
aacttgaaat
2401 agaaagagaa gagaaggaaa aactcaaaag agaggcaaaa gaaaacacag
ctactcttaa
15 2461 agaaaaaaaa gacaagaaaa cacaaacatt tttattggaa acacctgaaa
tttattggaa
2521 attggattct aaagcagttc cttcaciaaac tgtatctcga aatttcacat
cagttgatca
2581 tggcatatcc aaagataaaa gagactatct gtggacatct gccaaaaata
20 ctttatctac
2641 accattgcc aagggcatata cagtgaagac accaacaaaa ccaaaactac
agcaaagaga
2701 aaacttgaat ataccattg aagaaagtaa aaaaaagaga aaaatggcct
ttgaatttga
25 2761 tattaattca gatagttcag aaactactga tcttttgagc atggtttcag
aagaagagac
2821 attgaaaaca ctgtatagga acaataatcc accagcttct catctttgtg
tcaaaacacc
2881 aaaaaaggcc cttcatctc taacaacccc tggacctaca ctgaagtttg
30 gagctataag
2941 aaaaatgagg gaggaccgtt gggctgtaat tgctaaaatg gatagaaaaa
aaaaactaaa
3001 agaagctgaa aagttatttg ttttaatttca gagaatcagt gtagttaagg
agcctaataa
35 3061 cgtgaaactt atagttaata ttttgttctt atttgccaga gccacatttt
atctggaagt
3121 tgagacttaa aaaatacttg catgaatgat ttgtgtttct ttatatTTTT
agcctaaatg

3181 ttaactacat attgtctgga aacctgtcat tgtattcaga taattagatg
attatatatt
3241 gttgttactt tttcttgtat tcatgaaaac tgtttttact aagttttcaa
atttgtaaag
5 3301 ttagcctttg aatgctagga atgcattatt gagggtcatt ctttattott
tactattaaa
3361 atatttttga tgcaaaaaaa aaaaaaaaaa aaa
//

10

SSX-4

LOCUS NM_005636 576 bp mRNA linear

PRI 10-DEC-2001

DEFINITION Homo sapiens synovial sarcoma, X breakpoint 4 (SSX4),
15 mRNA.

ACCESSION NM_005636

VERSION NM_005636.1 GI:5032122

KEYWORDS .

SOURCE human.

20

ORGANISM Homo sapiens

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;

Euteleostomi;

Mammalia; Eutheria; Primates; Catarrhini; Hominidae;

Homo.

25

REFERENCE 1 (bases 1 to 576)

AUTHORS Gure,A.O., Tureci,O., Sahin,U., Tsang,S.,

Scanlan,M.J., Knuth,A.,

Pfreundschuh,M., Old,L.J. and Chen,Y.T.

TITLE SSX: a multigene family with several members

30

transcribed in normal

testis and human cancer

JOURNAL Int. J. Cancer 72 (6), 965-971 (1997)

MEDLINE 98021352

COMMENT PROVISIONAL REFSEQ: This record has not yet been

35

subject to final

NCBI review. The reference sequence was derived from

U90841.1.

FEATURES Location/Qualifiers

source 1..576


```

/organism="Homo sapiens"
/db_xref="taxon:9606"
/chromosome="X"
/map="Xp11.3"
5      gene      1..576
          /gene="SSX4"
          /db_xref="LocusID:6759"
          /db_xref="MIM:300326"
          CDS      1..567
10      /gene="SSX4"
          /note="Kruppel-associated box containing SSX
gene"
          /codon_start=1
          /db_xref="LocusID:6759"
15      /db_xref="MIM:300326"
          /product="synovial sarcoma, X breakpoint 4"
          /protein_id="NP_005627.1"
          /db_xref="GI:5032123"

20      /translation="MNGDDAFARRPRDDAQISEKLRKAFDDIAKYFSKKEWEKMKSSSEKIVY
VYMKLNYEVMTKLGFKVTLPPFMRSKRAADFHGNDFGNDRNHRNQVERPQMTFG
SLQRIFPKIMPKKPAEBENGLKEVPEASGPQNDGKQLCPPGNPSTLEKINKTSGPKRG
KHAWTHRLRERKQLVVYEEISDPEEDDE"
          misc_feature      70..246
25      /note="KRAB; Region: krueppel associated box"
BASE COUNT      187 a      127 c      150 g      112 t
ORIGIN
          1 atgaacggag acgacgcctt tgcaaggaga cccagggatg atgctcaa
atcagagaag
30      61 ttacgaaagg ccttcgatga tattgccaaa tacttctcta agaaagagt
ggaaaagatg
          121 aaatcctcgg agaaaatcgt ctatgtgtat atgaagctaa actatgaggt
catgactaaa
          181 ctaggtttca aggtcacctt cccacctttc atgcgtagta aacgggctgc
35      agacttccac
          241 gggaatgatt ttggtaacga tcgaaaccac aggaatcagg ttgaacgtcc
tcagatgact
          301 ttcggcagcc tccagagaat cttcccgaag atcatgccca agaagccagc
agaggaagaa

```

361 aatggtttga aggaagtgcc agaggcatct ggcccacaaa atgatgggaa
acagctgtgc

421 cccccgggaa atccaagtac cttggagaag attaacaaga catctggacc
caaaagggggg

5 481 aaacatgcct ggacccacag actgcgtgag agaaagcagc tggtggttta
tgaagagatc

541 agcgaccctg aggaagatga cgagtaactc ccctcg

10 All patents and publications mentioned in the specification
are indicative of the levels of those skilled in the art to which
the invention pertains. All patents and publications are herein
incorporated by reference to the same extent as if each individual
publication was specifically and individually indicated to be
incorporated by reference.

15 The invention illustratively described herein suitably may be
practiced in the absence of any element or elements, limitation or
limitations which is not specifically disclosed herein. The terms
and expressions which have been employed are used as terms of
description and not of limitation, and there is no intention that
20 in the use of such terms and expressions indicates the exclusion
of equivalents of the features shown and described or portions
thereof. It is recognized that various modifications are possible
within the scope of the invention claimed. Thus, it should be
understood that although the present invention has been
25 specifically disclosed by preferred embodiments and optional
features, modification and variation of the concepts herein
disclosed may be resorted to by those skilled in the art, and that
such modifications and variations are considered to be within the
scope of this invention as defined by the appended claims.

30

WHAT IS CLAIMED IS:

1. An isolated epitope, comprising a component selected from the group consisting of:
 - (i) a polypeptide having the sequence as disclosed in TABLE 1;
 - (ii) an epitope cluster comprising the polypeptide of (i);
 - (iii) a polypeptide having substantial similarity to (i) or (ii);
 - (iv) a polypeptide having functional similarity to any of (i) through (iii); and
 - (v) a nucleic acid encoding the polypeptide of any of (i) through (iv).
2. The epitope of claim 1, wherein the epitope is immunologically active.
3. The epitope of claim 1, wherein the polypeptide is less than about 30 amino acids in length.
4. The epitope of claim 1, wherein the polypeptide is 8 to 10 amino acids in length.
5. The epitope of claim 1, wherein the substantial or functional similarity comprises addition of at least one amino acid.
6. The epitope of claim 5, wherein the at least one additional amino acid is at an N-terminus of the polypeptide.
7. The epitope of claim 1, wherein the substantial or functional similarity comprises a substitution of at least one amino acid.
8. The epitope of claim 1, the polypeptide having affinity to an HLA-A2 molecule.
9. The epitope of claim 8, wherein the affinity is determined by an assay of binding.
10. The epitope of claim 8, wherein the affinity is determined by an assay of restriction of epitope recognition.
11. The epitope of claim 8, wherein the affinity is determined by a prediction algorithm.
12. The epitope of claim 1, the polypeptide having affinity to an HLA-B7 or HLA-B51 molecule.
13. The epitope of claim 1, wherein the polypeptide is a housekeeping epitope.
14. The epitope of claim 1, wherein the polypeptide corresponds to an epitope displayed on a tumor cell.
15. The epitope of claim 1, wherein the polypeptide corresponds to an epitope displayed on a neovasculature cell.
16. The epitope of claim 1, wherein the peptide is an immune epitope.
17. The epitope of claim 1 wherein the epitope is a nucleic acid.
18. A pharmaceutical composition comprising the peptide of claim 1 and a pharmaceutically acceptable adjuvant, carrier, diluent, or excipient.
19. The composition of claim 18, where the adjuvant is a polynucleotide.

20. The composition of claim 19 wherein the polynucleotide comprises a dinucleotide.
21. The composition of claim 20 wherein the dinucleotide is CpG.
22. The composition of claim 18, wherein the adjuvant is encoded by a polynucleotide.
23. The composition of claim 18 wherein the adjuvant is a cytokine.
- 5 24. The composition of claim 23 wherein the cytokine is GM-CSF.
25. The composition of claim 18 further comprising a professional antigen-presenting cell (pAPC).
26. The composition of claim 25, wherein the pAPC is a dendritic cell.
27. The composition of claim 18, further comprising a second epitope.
- 10 28. The composition of claim 27, wherein the second epitope is a polypeptide.
29. The composition of claim 27, wherein the second epitope is a nucleic acid.
30. The composition of claim 27, wherein the second epitope is a housekeeping epitope.
31. The composition of claim 27, wherein the second epitope is an immune epitope.
- 15 32. A pharmaceutical composition comprising the nucleic acid of claim 1 and a pharmaceutically acceptable adjuvant, carrier, diluent, or excipient.
33. A recombinant construct comprising the nucleic acid of Claim 1.
34. The construct of claim 33, further comprising a plasmid, a viral vector, or an artificial chromosome.
- 20 35. The construct of claim 33, further comprising a sequence encoding at least one feature selected from the group consisting of a second epitope, an IRES, an ISS, an NIS, and ubiquitin.
36. A purified antibody that specifically binds to the epitope of claim 1.
37. A purified antibody that specifically binds to a peptide-MHC protein complex
- 25 comprising the epitope of claim 1.
38. The antibody of claim 36 or claim 37, wherein the antibody is a monoclonal antibody.
39. A multimeric MHC-peptide complex comprising the epitope of claim 1.
40. An isolated T cell expressing a T cell receptor specific for an MHC-peptide
- 30 complex, the complex comprising the epitope of claim 1.
41. The T cell of claim 40, produced by an *in vitro* immunization.
42. The T cell of claim 40, isolated from an immunized animal.
43. A T cell clone comprising the T cell of claim 40.
44. A polyclonal population of T cells comprising the T cell of claim 40.
- 35 45. A pharmaceutical composition comprising the T cell of claim 40 and a pharmaceutically acceptable adjuvant, carrier, diluent, or excipient.

46. An isolated protein molecule comprising the binding domain of a T cell receptor specific for an MHC-peptide complex, the complex comprising the epitope of claim 1.
47. The protein of claim 46, wherein the protein is multivalent.
48. An isolated nucleic acid encoding the protein of claim 46.
- 5 49. A recombinant construct comprising the nucleic acid of claim 48.
50. A host cell expressing the recombinant construct of claim 33 or 49.
51. The host cell of claim 50, wherein the host cell is a dendritic cell, macrophage, tumor cell, or tumor-derived cell.
52. The host cell of claim 50, wherein the host cell is a bacterium, fungus, or
10 protozoan.
53. A pharmaceutical composition comprising the host cell of claim 50 and a pharmaceutically acceptable adjuvant, carrier, diluent, or excipient.
54. A vaccine or immunotherapeutic composition comprising at least one component selected from the group consisting of the epitope of claim 1; the composition of claim 18, 32, 45, or
15 53; the construct of claim 33; the T cell of claim 40, and the host cell of claim 50.
55. A method of treating an animal, comprising:
administering to an animal the vaccine or immunotherapeutic composition of claim
54.
56. The method of claim 55, wherein the administering step comprises a mode of
20 delivery selected from the group consisting of transdermal, intranodal, perinodal, oral, intravenous, intradermal, intramuscular, intraperitoneal, mucosal, aerosol inhalation, and instillation.
57. The method of claim 55, further comprising a step of assaying to determine a characteristic indicative of a state of a target cell or target cells.
58. The method of claim 57, comprising a first assaying step and a second assaying
25 step, wherein the first assaying step precedes the administering step, and wherein the second assaying step follows the administering step.
59. The method of claim 58, further comprising a step of comparing the characteristic determined in the first assaying step with the characteristic determined in the second assaying step to obtain a result.
60. The method of claim 59, wherein the result is selected from the group consisting
30 of: evidence of an immune response, a diminution in number of target cells, a loss of mass or size of a tumor comprising target cells, a decrease in number or concentration of an intracellular parasite infecting target cells.
61. A method of evaluating immunogenicity of a vaccine or immunotherapeutic
35 composition, comprising:

- administering to an animal the vaccine or immunotherapeutic composition of claim 54; and
evaluating immunogenicity based on a characteristic of the animal.
62. The method of claim 61, wherein the animal is HLA-transgenic.
- 5 63. A method of evaluating immunogenicity, comprising:
in vitro stimulation of a T cell with the vaccine or immunotherapeutic composition of claim 54; and
evaluating immunogenicity based on a characteristic of the T cell.
64. The method of claim 63, wherein the stimulation is a primary stimulation.
- 10 65. A method of making a passive/adoptive immunotherapeutic, comprising:
combining the T cell of claim 40 or the host cell of claim 50 with a pharmaceutically acceptable adjuvant, carrier, diluent, or excipient.
66. A method of determining specific T cell frequency comprising the step of contacting T cells with a MHC-peptide complex comprising the epitope of claim 1.
- 15 67. The method of claim 66, wherein the contacting step comprises at least one feature selected from the group consisting of immunization, restimulation, detection, and enumeration.
68. The method of Claim 66, further comprising ELISPOT analysis, limiting dilution analysis, flow cytometry, *in situ* hybridization, the polymerase chain reaction or any combination thereof.
- 20 69. A method of evaluating immunologic response, comprising the method of claim 66 carried out prior to and subsequent to an immunization step.
70. A method of evaluating immunologic response, comprising:
determining frequency, cytokine production, or cytolytic activity of T cells, prior to and subsequent to a step of stimulation with MHC-peptide complexes comprising the epitope of claim 1.
- 25 71. A method of diagnosing a disease comprising:
contacting a subject tissue with at least one component selected from the group consisting of the T cell of claim 40, the host cell of claim 50, the antibody of claim 36, the protein of claim 46; and
diagnosing the disease based on a characteristic of the tissue or of the component.
- 30 72. The method of claim 71, wherein the contacting step takes place *in vivo*.
73. The method of claim 71, wherein the contacting step takes place *in vitro*.
74. A method of making a vaccine, comprising:
combining at least one component selected from the group consisting of the epitope of claim 1; the composition of claim 18, 32, 45, or 53; the construct of claim 33;
- 35

the T cell of claim 40, and the host cell of claim 50, with a pharmaceutically acceptable adjuvant, carrier, diluent, or excipient.

5 75. A computer readable medium having recorded thereon the sequence of any one of SEQ ID NOS: X -Y, in a machine having a hardware or software that calculates the physical, biochemical, immunologic, or molecular genetic properties of a molecule embodying said sequence.

 76. A method of treating an animal comprising combining the method of claim 55 combined with at least one mode of treatment selected from the group of radiation therapy, chemotherapy, biochemotherapy, and surgery.

10 77. An isolated polypeptide comprising an epitope cluster from a target-associated antigen having the sequence as disclosed in Tables 25-44, wherein the amino acid sequence consists of not more than about 80% of the amino acid sequence of the antigen.

 78. A vaccine or immunotherapeutic product comprising the polypeptide of claim 78.

 79. An isolated polynucleotide encoding the polypeptide of claim 78.

15 80. A vaccine or immunotherapeutic product comprising the polynucleotide of claim 80.

 81. The polynucleotide of claim 79 or 80, wherein the polynucleotide is DNA.

 82. The polynucleotide of claim 79 or 80, wherein the polynucleotide is RNA.

1			50
CTAG HUMAN NY-ESO	(1)	MOAEGRTGGSTGDDADGPGGPGIPDGPGGNAGGPGGEAGATGGRGPRGAGA	
AAD05202 - CAG-3	(1)	MOAEGRTGGSTGDDADGPGGPGIPDGPGGNAGGPGGEAGATGGRGPRGAGA	
CAA11044 -LAGE-1a	(1)	MOAEGRTGGSTGDDADGPGGPGIPDGPGGNAGGPGGEAGATGGRGPRGAGA	
CAA10194 - LAGE-1s	(1)	MOAEGRTGGSTGDDADGPGGPGIPDGPGGNAGGPGGEAGATGGRGPRGAGA	
CAA11043 - LAGE-1b	(1)	MOAEGRTGGSTGDDADGPGGPGIPDGPGGNAGGPGGEAGATGGRGPRGAGA	
CAA10196 - LAGE-1L	(1)	MOAEGRTGGSTGDDADGPGGPGIPDGPGGNAGGPGGEAGATGGRGPRGAGA	
AAH02833 CT-2	(1)	MOAEGRTGGSTGDDADGPGGPGIPDGPGGNAGGPGGEAGATGGRGPRGAGA	
Consensus	(1)	MOAEGRTGGSTGDDADGPGGPGIPDGPGGNAGGPGGEAGATGGRGPRGAGA	100
CTAG HUMAN NY-ESO	(51)	ARASGPGGAPRGPHGGAASLNGCCRCGANGPESRLLEFMMAMPFESPM	51
AAD05202 - CAG-3	(51)	ARASGPGGAPRGPHGGAASLNGCCRCGANGPESRLLEFMMAMPFESPM	
CAA11044 -LAGE-1a	(51)	ARASGPGGAPRGPHGGAASLNGCCRCGANGPESRLLEFMMAMPFESPM	
CAA10194 - LAGE-1s	(51)	ARASGPGGAPRGPHGGAASLNGCCRCGANGPESRLLEFMMAMPFESPM	
CAA11043 - LAGE-1b	(51)	ARASGPGGAPRGPHGGAASLNGCCRCGANGPESRLLEFMMAMPFESPM	
CAA10196 - LAGE-1L	(51)	ARASGPGGAPRGPHGGAASLNGCCRCGANGPESRLLEFMMAMPFESPM	
AAH02833 CT-2	(51)	ARASGPGGAPRGPHGGAASLNGCCRCGANGPESRLLEFMMAMPFESPM	
Consensus	(51)	ARASGPGGAPRGPHGGAASLNGCCRCGANGPESRLLEFMMAMPFESPM	100
CTAG HUMAN NY-ESO	(101)	EAELARRSLQDAFPLEVRCVPLKEFTVSGNLLTIRRAADHRLQLSIS	101
AAD05202 - CAG-3	(101)	EAELARRSLQDAFPLEVRCVPLKEFTVSGNLLTIRRAADHRLQLSIS	
CAA11044 -LAGE-1a	(101)	EAELARRSLQDAFPLEVRCVPLKEFTVSGNLLTIRRAADHRLQLSIS	
CAA10194 - LAGE-1s	(101)	EAELARRSLQDAFPLEVRCVPLKEFTVSGNLLTIRRAADHRLQLSIS	
CAA11043 - LAGE-1b	(101)	EAELARRSLQDAFPLEVRCVPLKEFTVSGNLLTIRRAADHRLQLSIS	
CAA10196 - LAGE-1L	(101)	EAELARRSLQDAFPLEVRCVPLKEFTVSGNLLTIRRAADHRLQLSIS	
AAH02833 CT-2	(101)	EAELARRSLQDAFPLEVRCVPLKEFTVSGNLLTIRRAADHRLQLSIS	
Consensus	(101)	EAELARRSLQDAFPLEVRCVPLKEFTVSGNLLTIRRAADHRLQLSIS	200
CTAG HUMAN NY-ESO	(151)	SGEGQSSMMWITQCFLPVFLAQPSGQRR	151
AAD05202 - CAG-3	(151)	SGEGQSSMMWITQCFLPVFLAQPSGQRR	
CAA11044 -LAGE-1a	(151)	SGEGQSSMMWITQCFLPVFLAQPSGQRR	
CAA10194 - LAGE-1s	(151)	SGEGQSSMMWITQCFLPVFLAQPSGQRR	
CAA11043 - LAGE-1b	(151)	SGEGQSSMMWITQCFLPVFLAQPSGQRR	
CAA10196 - LAGE-1L	(151)	SGEGQSSMMWITQCFLPVFLAQPSGQRR	
AAH02833 CT-2	(151)	SGEGQSSMMWITQCFLPVFLAQPSGQRR	
Consensus	(151)	SGEGQSSMMWITQCFLPVFLAQPSGQRR	200
CTAG HUMAN NY-ESO	(181)	-----	181
AAD05202 - CAG-3	(181)	-----	
CAA11044 -LAGE-1a	(181)	-----	
CAA10194 - LAGE-1s	(181)	-----	
CAA11043 - LAGE-1b	(201)	FNVMFSAPI	
CAA10196 - LAGE-1L	(201)	FNVMFSAPI	
AAH02833 CT-2	(201)	FNVMFSAPI	
Consensus	(201)	FNVMFSAPI	

FIG 1




FIG 2

FACSscan Analysis of Binding Assay to Determine the Binding
Ability of Tyrosinase 208-216 Peptide to MHC Class I




FIG 3A

**FACscan Analysis of Binding Assay to Determine the Binding
Ability of Tyrosinase 207-215 Peptide to MHC Class I**




FIG 3B




FIG 4

Comparison of Peptides Binding Affinity to HLA A2
by Binding Assay




FIG 5

SSX2₄₁₋₄₉ specific lysis by CTL from Peptide injected HHD1 mice




FIG 6




Fig. 7A. Pool sequencing of PSMA_163-192 Digested for 60 min by Proteasome




Fig. Pool sequencing of PSMA_163-192 Digested for 60 min by Proteasome

FIG 7B




Fig. Pool sequencing of PSMA_163-192 Digested for 60 min by Proteasome

FIG 7C

HLA A2 Binding Assay




FIG 8




Fig. Pool sequencing of PSMA_281-310 digested for 60 min by Proteasome

Comparison of Peptides Binding Affinity to HLA A2 by Binding Assay




FIG 10

Autologous DC Present A1 Peptide to CD8 T cell




FIG 11

Secretion of IFN γ Was Blocked by Anti-A1 Antibody




FIG 12

Comparison of Peptides Binding Affinity to HLA A2
by Binding Assay




FIG 13

Comparison of Peptides Binding Affinity to HLA A2
by Binding Assay






FIG 14



Graphs show lysis of unpulsed EL4 cells (open circles) and EL4 cells pulsed with gp33 peptide (solid triangles). Symbols represent individual mice and one of three similar experiments is shown.

FIG 15



Mean tumor volumes $\pm 1SD$ are shown for mice immunized with pEFGPL33A DNA (solid circles) or control pEGFP-N3 DNA (open triangles). Numbers in brackets indicate number of mice with tumors / total number of mice in group.. One of two similar experiments is shown.

FIG 16




FIG 17

CORRECTED VERSION

(19) World Intellectual Property
Organization
International Bureau








(43) International Publication Date
30 January 2003 (30.01.2003)

PCT

(10) International Publication Number
WO 2003/008537 A2

- (51) International Patent Classification⁷: C12N Hills, CA 91367 (US). XIE, Zhidong; 22228 Victory Blvd., H-111, Woodland Hills, CA 91367 (US).
- (21) International Application Number: PCT/US2002/010189 (74) Agent: TAHMASSEBI, Sam; Knobbe, Martens, Olson & Bear, LLP, 16th Floor, 620 Newport Center Drive, Newport Beach, CA 92660 (US).
- (22) International Filing Date: 29 March 2002 (29.03.2002)
- (25) Filing Language: English (81) Designated State (national): AU.
- (26) Publication Language: English Published:
— without international search report and to be republished upon receipt of that report
- (30) Priority Data:
60/282,211 6 April 2001 (06.04.2001) US
60/337,017 7 November 2001 (07.11.2001) US
60/363,210 7 March 2002 (07.03.2002) US (48) Date of publication of this corrected version:
19 February 2004
- (71) Applicant: MANNKIND CORPORATION [US/US]; 28903 North Avenue Paine, Valencia, CA 91355 (US). (15) Information about Correction:
see PCT Gazette No. 08/2004 of 19 February 2004, Section II
- (72) Inventors: SIMARD, John, J.L.; 11918 Laughton Way, Northridge, CA 91326 (US). DIAMOND, David, C.; 23135 Schoenborn Street, West Hills, CA 91304 (US). LIU, Liping; 22228 Victory Blvd., H-111, Woodland Hills, CA 91367 (US). For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.
- (54) Title: EPITOPE SEQUENCES

281 310
RGIAEAVGLPSIPVHPIGYYDAQKLLLEKMG



Pool sequencing of PSMA_281_310 Digested for 60 min by Proteasome

WO 2003/008537 A2

(57) Abstract: Disclosed herein are polypeptides, including epitopes, clusters, and antigens. Also disclosed are compositions including said polypeptides and methods for their use.

EPITOPE SEQUENCES

Field of the Invention

5 The present invention generally relates to peptides, and nucleic acids encoding peptides, that are useful epitopes of target-associated antigens. More specifically, the invention relates to epitopes that have a high affinity for MHC class I and that are produced by target-specific proteasomes.

Description of the Related Art

Neoplasia and the Immune System

10 The neoplastic disease state commonly known as cancer is thought to result generally from a single cell growing out of control. The uncontrolled growth state typically results from a multi-step process in which a series of cellular systems fail, resulting in the genesis of a neoplastic cell. The resulting neoplastic cell rapidly reproduces itself, forms one or more tumors, and eventually may cause the death of the host.

15 Because the progenitor of the neoplastic cell shares the host's genetic material, neoplastic cells are largely unassailed by the host's immune system. During immune surveillance, the process in which the host's immune system surveys and localizes foreign materials, a neoplastic cell will appear to the host's immune surveillance machinery as a "self" cell.

Viruses and the Immune System

20 In contrast to cancer cells, virus infection involves the expression of clearly non-self antigens. As a result, many virus infections are successfully dealt with by the immune system with minimal clinical sequelae. Moreover, it has been possible to develop effective vaccines for many of those infections that do cause serious disease. A variety of vaccine approaches have been used successfully to combat various diseases. These approaches include subunit vaccines consisting of individual proteins produced through recombinant DNA technology. Notwithstanding these
25 advances, the selection and effective administration of minimal epitopes for use as viral vaccines has remained problematic.

In addition to the difficulties involved in epitope selection stands the problem of viruses that have evolved the capability of evading a host's immune system. Many viruses, especially viruses that establish persistent infections, such as members of the herpes and pox virus families,
30 produce immunomodulatory molecules that permit the virus to evade the host's immune system. The effects of these immunomodulatory molecules on antigen presentation may be overcome by the targeting of select epitopes for administration as immunogenic compositions. To better understand the interaction of neoplastic cells and virally infected cells with the host's immune system, a discussion of the system's components follows below.

35 The immune system functions to discriminate molecules endogenous to an organism ("self" molecules) from material exogenous or foreign to the organism ("non-self" molecules). The

immune system has two types of adaptive responses to foreign bodies based on the components that mediate the response: a humoral response and a cell-mediated response. The humoral response is mediated by antibodies, while the cell-mediated response involves cells classified as lymphocytes. Recent anticancer and antiviral strategies have focused on mobilizing the host immune system as a means of anticancer or antiviral treatment or therapy.

The immune system functions in three phases to protect the host from foreign bodies: the cognitive phase, the activation phase, and the effector phase. In the cognitive phase, the immune system recognizes and signals the presence of a foreign antigen or invader in the body. The foreign antigen can be, for example, a cell surface marker from a neoplastic cell or a viral protein. Once the system is aware of an invading body, antigen specific cells of the immune system proliferate and differentiate in response to the invader-triggered signals. The last stage is the effector stage in which the effector cells of the immune system respond to and neutralize the detected invader.

An array of effector cells implements an immune response to an invader. One type of effector cell, the B cell, generates antibodies targeted against foreign antigens encountered by the host. In combination with the complement system, antibodies direct the destruction of cells or organisms bearing the targeted antigen. Another type of effector cell is the natural killer cell (NK cell), a type of lymphocyte having the capacity to spontaneously recognize and destroy a variety of virus infected cells as well as malignant cell types. The method used by NK cells to recognize target cells is poorly understood.

Another type of effector cell, the T cell, has members classified into three subcategories, each playing a different role in the immune response. Helper T cells secrete cytokines which stimulate the proliferation of other cells necessary for mounting an effective immune response, while suppressor T cells down-regulate the immune response. A third category of T cell, the cytotoxic T cell (CTL), is capable of directly lysing a targeted cell presenting a foreign antigen on its surface.

The Major Histocompatibility Complex and T Cell Target Recognition

T cells are antigen-specific immune cells that function in response to specific antigen signals. B lymphocytes and the antibodies they produce are also antigen-specific entities. However, unlike B lymphocytes, T cells do not respond to antigens in a free or soluble form. For a T cell to respond to an antigen, it requires the antigen to be processed to peptides which are then bound to a presenting structure encoded in the major histocompatibility complex (MHC). This requirement is called "MHC restriction" and it is the mechanism by which T cells differentiate "self" from "non-self" cells. If an antigen is not displayed by a recognizable MHC molecule, the T cell will not recognize and act on the antigen signal. T cells specific for a peptide bound to a recognizable MHC molecule bind to these MHC-peptide complexes and proceed to the next stages of the immune response.

There are two types of MHC, class I MHC and class II MHC. T Helper cells (CD4⁺) predominately interact with class II MHC proteins while cytolytic T cells (CD8⁺) predominately

interact with class I MHC proteins. Both classes of MHC protein are transmembrane proteins with a majority of their structure on the external surface of the cell. Additionally, both classes of MHC proteins have a peptide binding cleft on their external portions. It is in this cleft that small fragments of proteins, endogenous or foreign, are bound and presented to the extracellular environment.

5 Cells called "professional antigen presenting cells" (pAPCs) display antigens to T cells using the MHC proteins but additionally express various co-stimulatory molecules depending on the particular state of differentiation/activation of the pAPC. When T cells, specific for the peptide bound to a recognizable MHC protein, bind to these MHC-peptide complexes on pAPCs, the specific co-stimulatory molecules that act upon the T cell direct the path of differentiation/activation taken by the
10 T cell. That is, the co-stimulation molecules affect how the T cell will act on antigenic signals in future encounters as it proceeds to the next stages of the immune response.

 As discussed above, neoplastic cells are largely ignored by the immune system. A great deal of effort is now being expended in an attempt to harness a host's immune system to aid in combating the presence of neoplastic cells in a host. One such area of research involves the formulation of
15 anticancer vaccines.

Anticancer Vaccines

 Among the various weapons available to an oncologist in the battle against cancer is the immune system of the patient. Work has been done in various attempts to cause the immune system to combat cancer or neoplastic diseases. Unfortunately, the results to date have been
20 largely disappointing. One area of particular interest involves the generation and use of anticancer vaccines.

 To generate a vaccine or other immunogenic composition, it is necessary to introduce to a subject an antigen or epitope against which an immune response may be mounted. Although neoplastic cells are derived from and therefore are substantially identical to normal cells on a genetic level, many neoplastic cells are known to present tumor-associated antigens (TuAAs). In
25 theory, these antigens could be used by a subject's immune system to recognize these antigens and attack the neoplastic cells. In reality, however, neoplastic cells generally appear to be ignored by the host's immune system.

 A number of different strategies have been developed in an attempt to generate vaccines with activity against neoplastic cells. These strategies include the use of tumor-associated antigens as immunogens. For example, U.S. Patent No. 5,993,828, describes a method for producing an
30 immune response against a particular subunit of the Urinary Tumor Associated Antigen by administering to a subject an effective dose of a composition comprising inactivated tumor cells having the Urinary Tumor Associated Antigen on the cell surface and at least one tumor associated antigen selected from the group consisting of GM-2, GD-2, Fetal Antigen and Melanoma
35

Associated Antigen. Accordingly, this patent describes using whole, inactivated tumor cells as the immunogen in an anticancer vaccine.

Another strategy used with anticancer vaccines involves administering a composition containing isolated tumor antigens. In one approach, MAGE-A1 antigenic peptides were used as an immunogen. (See Chaux, P., *et al.*, "Identification of Five MAGE-A1 Epitopes Recognized by Cytolytic T Lymphocytes Obtained by *In Vitro* Stimulation with Dendritic Cells Transduced with MAGE-A1," J. Immunol., 163(5):2928-2936 (1999)). There have been several therapeutic trials using MAGE-A1 peptides for vaccination, although the effectiveness of the vaccination regimes was limited. The results of some of these trials are discussed in Vose, J.M., "Tumor Antigens Recognized by T Lymphocytes," 10th European Cancer Conference, Day 2, Sept. 14, 1999.

In another example of tumor associated antigens used as vaccines, Scheinberg, *et al.* treated 12 chronic myelogenous leukemia (CML) patients already receiving interferon (IFN) or hydroxyurea with 5 injections of class I-associated bcr-abl peptides with a helper peptide plus the adjuvant QS-21. Scheinberg, D.A., *et al.*, "BCR-ABL Breakpoint Derived Oncogene Fusion Peptide Vaccines Generate Specific Immune Responses in Patients with Chronic Myelogenous Leukemia (CML) [Abstract 1665], American Society of Clinical Oncology 35th Annual Meeting, Atlanta (1999). Proliferative and delayed type hypersensitivity (DTH) T cell responses indicative of T-helper activity were elicited, but no cytolytic killer T cell activity was observed within the fresh blood samples.

Additional examples of attempts to identify TuAAs for use as vaccines are seen in the recent work of Cebon, *et al.* and Scheibenbogen, *et al.* Cebon, *et al.* immunized patients with metastatic melanoma using intradermally administered MART-1₂₆₋₃₅ peptide with IL-12 in increasing doses given either subcutaneously or intravenously. Of the first 15 patients, 1 complete remission, 1 partial remission, and 1 mixed response were noted. Immune assays for T cell generation included DTH, which was seen in patients with or without IL-12. Positive CTL assays were seen in patients with evidence of clinical benefit, but not in patients without tumor regression. Cebon, *et al.*, "Phase I Studies of Immunization with Melan-A and IL-12 in HLA A2+ Positive Patients with Stage III and IV Malignant Melanoma," [Abstract 1671], American Society of Clinical Oncology 35th Annual Meeting, Atlanta (1999).

Scheibenbogen, *et al.* immunized 18 patients with 4 HLA class I restricted tyrosinase peptides, 16 with metastatic melanoma and 2 adjuvant patients. Scheibenbogen, *et al.*, "Vaccination with Tyrosinase peptides and GM-CSF in Metastatic Melanoma: a Phase II Trial," [Abstract 1680], American Society of Clinical Oncology 35th Annual Meeting, Atlanta (1999). Increased CTL activity was observed in 4/15 patients, 2 adjuvant patients, and 2 patients with evidence of tumor regression. As in the trial by Cebon, *et al.*, patients with progressive disease did

not show boosted immunity. In spite of the various efforts expended to date to generate efficacious anticancer vaccines, no such composition has yet been developed.

Antiviral Vaccines

5 Vaccine strategies to protect against viral diseases have had many successes. Perhaps the most notable of these is the progress that has been made against the disease small pox, which has been driven to extinction. The success of the polio vaccine is of a similar magnitude.

Viral vaccines can be grouped into three classifications: live attenuated virus vaccines, such as vaccinia for small pox, the Sabin poliovirus vaccine, and measles mumps and rubella; whole killed or inactivated virus vaccines, such as the Salk poliovirus vaccine, hepatitis A virus vaccine and the typical influenza virus vaccines; and subunit vaccines, such as hepatitis B. Due to
10 their lack of a complete viral genome, subunit vaccines offer a greater degree of safety than those based on whole viruses.

The paradigm of a successful subunit vaccine is the recombinant hepatitis B vaccine based on the viruses envelope protein. Despite much academic interest in pushing the reductionist subunit concept beyond single proteins to individual epitopes, the efforts have yet to bear much
15 fruit. Viral vaccine research has also concentrated on the induction of an antibody response although cellular responses also occur. However, many of the subunit formulations are particularly poor at generating a CTL response.

Summary of the Invention

20 Previous methods of priming professional antigen presenting cells (pAPCs) to display target cell epitopes have relied simply on causing the pAPCs to express target-associated antigens (TAAs), or epitopes of those antigens which are thought to have a high affinity for MHC I molecules. However, the proteasomal processing of such antigens results in presentation of epitopes on the pAPC that do not correspond to the epitopes present on the target cells.

25 Using the knowledge that an effective cellular immune response requires that pAPCs present the same epitope that is presented by the target cells, the present invention provides epitopes that have a high affinity for MHC I, and that correspond to the processing specificity of the housekeeping proteasome, which is active in peripheral cells. These epitopes thus correspond to those presented on target cells. The use of such epitopes in vaccines can activate the cellular
30 immune response to recognize the correctly processed TAA and can result in removal of target cells that present such epitopes. In some embodiments, the housekeeping epitopes provided herein can be used in combination with immune epitopes, generating a cellular immune response that is competent to attack target cells both before and after interferon induction. In other embodiments the epitopes are useful in the diagnosis and monitoring of the target-associated disease and in the
35 generation of immunological reagents for such purposes.

Embodiments of the invention relate to isolated epitopes and antigens or polypeptides that comprise the epitopes. Preferred embodiments include an epitope or antigen having the sequence as disclosed in TABLE 1. Other embodiments can include an epitope cluster comprising a polypeptide from Table 1. Further, embodiments include a polypeptide having substantial similarity to the already mentioned epitopes, antigens, or clusters. Other preferred embodiments include a polypeptide having functional similarity to any of the above. Still further embodiments relate to a nucleic acid encoding the polypeptide of any of the epitopes, clusters, antigens, and polypeptides from Table 1 and mentioned herein.

The epitope can be immunologically active. The polypeptide comprising the epitope can be less than about 30 amino acids in length, more preferably, the polypeptide is 8 to 10 amino acids in length, for example. Substantial or functional similarity can include addition of at least one amino acid, for example, and the at least one additional amino acid can be at an N-terminus of the polypeptide. The substantial or functional similarity can include a substitution of at least one amino acid.

The epitope, cluster, or polypeptide comprising the same can have affinity to an HLA-A2 molecule. The affinity can be determined by an assay of binding, by an assay of restriction of epitope recognition, by a prediction algorithm, and the like. The epitope, cluster, or polypeptide comprising the same can have affinity to an HLA-B7, HLA-B51 molecule, and the like.

In preferred embodiments the polypeptide can be a housekeeping epitope. The epitope or polypeptide can correspond to an epitope displayed on a tumor cell, to an epitope displayed on a neovasculature cell, and the like. The epitope or polypeptide can be an immune epitope. The epitope, cluster and/or polypeptide can be a nucleic acid.

Other embodiments relate to pharmaceutical compositions comprising the polypeptides, including an epitope from Table 1, a cluster, or a polypeptide comprising the same and a pharmaceutically acceptable adjuvant, carrier, diluent, excipient, and the like. The adjuvant can be a polynucleotide. The polynucleotide can include a dinucleotide. The dinucleotide can be CpG, for example. The adjuvant can be encoded by a polynucleotide. The adjuvant can be a cytokine and the cytokine can be, for example, GM-CSF.

The pharmaceutical compositions can further include a professional antigen-presenting cell (pAPC). The pAPC can be a dendritic cell, for example. The pharmaceutical composition can further include a second epitope. The second epitope can be a polypeptide. The second epitope can be a nucleic acid. The second epitope can be a housekeeping epitope, an immune epitope, and the like.

Still further embodiments relate to pharmaceutical compositions that include any of the nucleic acids discussed herein, including those that encode polypeptides that comprise epitopes or

antigens from Table 1. Such compositions can include a pharmaceutically acceptable adjuvant, carrier, diluent, excipient, and the like.

5 Other embodiments relate to recombinant constructs that include such a nucleic acid as described herein, including those that encode polypeptides that comprise epitopes or antigens from Table 1. The constructs can further include a plasmid, a viral vector, an artificial chromosome, and the like. The construct can further include a sequence encoding at least one feature, such as for example, a second epitope, an IRES, an ISS, an NIS, ubiquitin.

10 Further embodiments relate to purified antibodies that specifically bind to at least one of the epitopes in Table 1A. Other embodiments relate to purified antibodies that specifically bind to a peptide-MHC protein complex comprising an epitope disclosed in Table 1A or any other suitable epitope. The antibody from any embodiment can be a monoclonal antibody.

Still other embodiments relate to multimeric MHC-peptide complexes that include an epitope, such as, for example, an epitope disclosed in Table 1.

15 Embodiments relate to isolated T cells expressing a T cell receptor specific for an MHC-peptide complex. The complex can include an epitope, such as, for example, an epitope disclosed in Table 1 of claim 1. The T cell can be produced by an *in vitro* immunization. The T cell can be isolated from an immunized animal. Embodiments relate to T cell clones, including cloned T cells, such as those discussed above. Embodiments also relate to polyclonal population of T cells. Such populations can include a T cell, as described above, for example.

20 Still further embodiments relate to pharmaceutical compositions that include a T cell, such as those described above, for example, and a pharmaceutically acceptable adjuvant, carrier, diluent, excipient, and the like.

25 Embodiments of the invention relate to isolated protein molecules comprising the binding domain of a T cell receptor specific for an MHC-peptide complex. The complex can include an epitope disclosed in Table 1. The protein can be multivalent. Other embodiments relate to isolated nucleic acids encoding such proteins. Still further embodiments relate to recombinant constructs that include such nucleic acids.

30 Other embodiments of the invention relate to host cells expressing the recombinant construct described herein, including constructs encoding an epitope, cluster or polypeptide comprising the same, disclosed in Table 1, for example. The host cell can be a dendritic cell, macrophage, tumor cell, tumor-derived cell, and the like. The host cell can be a bacterium, fungus, protozoan and the like. Embodiments also relate to pharmaceutical compositions that include a host cell, such as those discussed herein, and a pharmaceutically acceptable adjuvant, carrier, diluent, excipient, and the like.

35 Still other embodiments relate to vaccines or immunotherapeutic compositions that include at least one component, such as, for example, an epitope disclosed in Table 1 or otherwise

described herein; a cluster that includes such an epitope, an antigen or polypeptide that includes such an epitope; a composition described above and herein; a construct, a T cell, or a host cell as described above and herein.

Further embodiments relate to methods of treating an animal. The methods can include administering to an animal a vaccine or immunotherapeutic composition, including those disclosed above and herein. The administering step can include a mode of delivery, such as, for example, transdermal, intranodal, perinodal, oral, intravenous, intradermal, intramuscular, intraperitoneal, mucosal, aerosol inhalation, instillation, and the like. The method can further include a step of assaying to determine a characteristic indicative of a state of a target cell or target cells. The method can include a first assaying step and a second assaying step, wherein the first assaying step precedes the administering step, and wherein the second assaying step follows the administering step. The method can further include a step of comparing the characteristic determined in the first assaying step with the characteristic determined in the second assaying step to obtain a result. The result can be for example, evidence of an immune response, a diminution in number of target cells, a loss of mass or size of a tumor comprising target cells, a decrease in number or concentration of an intracellular parasite infecting target cells, and the like.

Embodiments relate to methods of evaluating immunogenicity of a vaccine or immunotherapeutic composition. The methods can include administering to an animal a vaccine or immunotherapeutic, such as those described above and elsewhere herein, and evaluating immunogenicity based on a characteristic of the animal. The animal can be HLA-transgenic.

Other embodiments relate to methods of evaluating immunogenicity that include *in vitro* stimulation of a T cell with the vaccine or immunotherapeutic composition, such as those described above and elsewhere herein, and evaluating immunogenicity based on a characteristic of the T cell. The stimulation can be a primary stimulation.

Still further embodiments relate to methods of making a passive/adoptive immunotherapeutic. The methods can include combining a T cell or a host cell, such as those described above and elsewhere herein, with a pharmaceutically acceptable adjuvant, carrier, diluent, excipient, and the like.

Other embodiments relate to methods of determining specific T cell frequency, and can include the step of contacting T cells with a MHC-peptide complex comprising an epitope disclosed in Table 1, or a complex comprising a cluster or antigen comprising such an epitope. The contacting step can include at least one feature, such as, for example, immunization, restimulation, detection, enumeration, and the like. The method can further include ELISPOT analysis, limiting dilution analysis, flow cytometry, *in situ* hybridization, the polymerase chain reaction, any combination thereof, and the like.

Embodiments relate to methods of evaluating immunologic response. The methods can include the above-described methods determining specific T cell frequency carried out prior to and subsequent to an immunization step.

Another embodiment relates to methods of evaluating immunologic response. The methods can include determining frequency, cytokine production, or cytolytic activity of T cells, prior to and subsequent to a step of stimulation with MHC-peptide complexes comprising an epitope, such as, for example an epitope from Table 1, a cluster or a polypeptide comprising such an epitope.

Further embodiments relate to methods of diagnosing a disease. The methods can include contacting a subject tissue with at least one component, including, for example, a T cell, a host cell, an antibody, a protein, including those described above and elsewhere herein; and diagnosing the disease based on a characteristic of the tissue or of the component. The contacting step can take place *in vivo*. The contacting step can take place *in vitro*.

Still other embodiments relate to methods of making a vaccine. The methods can include combining at least one component, an epitope, a composition, a construct, a T cell, a host cell; including any of those described above and elsewhere herein, with a pharmaceutically acceptable adjuvant, carrier, diluent, excipient, and the like.

Embodiments relate to computer readable media having recorded thereon the sequence of any one of SEQ ID NOS: 1 -602, in a machine having a hardware or software that calculates the physical, biochemical, immunologic, or molecular genetic properties of a molecule embodying said sequence.

Still other embodiments relate to methods of treating an animal. The methods can include combining the method of treating an animal that includes administering to the animal a vaccine or immunotherapeutic composition, such as described above and elsewhere herein, combined with at least one mode of treatment, including, for example, radiation therapy, chemotherapy, biochemotherapy, surgery, and the like.

Further embodiments relate to isolated polypeptides that include an epitope cluster from a target-associated antigen having the sequence as disclosed in any one of Tables 25-44, wherein the amino acid sequence includes not more than about 80% of the amino acid sequence of the antigen.

Other embodiments relate to vaccines or immunotherapeutic products that include an isolated peptide as described above and elsewhere herein. Still other embodiments relate to isolated polynucleotides encoding a polypeptide as described above and elsewhere herein. Other embodiments relate vaccines or immunotherapeutic products that include these polynucleotides. The polynucleotide can be DNA or RNA.

Brief Description of the Drawings

Figure 1 is a sequence alignment of NY-ESO-1 and several similar protein sequences.

Figure 2 graphically represents a plasmid vaccine backbone useful for delivering nucleic acid-encoded epitopes.

5 Figures 3A and 3B are FACS profiles showing results of HLA-A2 binding assays for tyrosinase₂₀₇₋₂₁₅ and tyrosinase₂₀₈₋₂₁₆.

Figure 4 is a T=120 min. time point mass spectrum of the fragments produced by proteasomal cleavage of SSX-2₃₁₋₆₈.

Figure 5 shows a binding curve for HLA-A2:SSX-2₄₁₋₄₉ with controls.

10 Figure 6 shows specific lysis of SSX-2₄₁₋₄₉-pulsed targets by CTL from SSX-2₄₁₋₄₉-immunized HLA-A2 transgenic mice.

Figure 7A, B, and C show results of N-terminal pool sequencing of a T=60 min. time point aliquot of the PSMA₁₆₃₋₁₉₂ proteasomal digest.

15 Figure 8 shows binding curves for HLA-A2:PSMA₁₆₈₋₁₇₇ and HLA-A2:PSMA₂₈₈₋₂₉₇ with controls.

Figure 9 shows results of N-terminal pool sequencing of a T=60 min. time point aliquot of the PSMA₂₈₁₋₃₁₀ proteasomal digest.

Figure 10 shows binding curves for HLA-A2:PSMA₄₆₁₋₄₆₉, HLA-A2:PSMA₄₆₀₋₄₆₉, and HLA-A2:PSMA₆₆₃₋₆₇₁, with controls.

20 Figure 11 shows the results of a γ -IFN-based ELISPOT assay detecting PSMA₄₆₃₋₄₇₁-reactive HLA-A1⁺ CD8⁺ T cells.

Figure 12 shows blocking of reactivity of the T cells used in figure 10 by anti-HLA-A1 mAb, demonstrating HLA-A1-restricted recognition.

Figure 13 shows a binding curve for HLA-A2:PSMA₆₆₃₋₆₇₁, with controls.

25 Figure 14 shows a binding curve for HLA-A2:PSMA₆₆₂₋₆₇₁, with controls.

Figure 15. Comparison of anti-peptide CTL responses following immunization with various doses of DNA by different routes of injection.

Figure 16. Growth of transplanted gp33 expressing tumor in mice immunized by i.ln. injection of gp33 epitope-expressing, or control, plasmid.

30 Figure 17. Amount of plasmid DNA detected by real-time PCR in injected or draining lymph nodes at various times after i.ln. of i.m. injection, respectively.

Detailed Description of the Preferred Embodiment

Definitions

35 Unless otherwise clear from the context of the use of a term herein, the following listed terms shall generally have the indicated meanings for purposes of this description.

PROFESSIONAL ANTIGEN-PRESENTING CELL (pAPC) – a cell that possesses T cell costimulatory molecules and is able to induce a T cell response. Well characterized pAPCs include dendritic cells, B cells, and macrophages.

PERIPHERAL CELL – a cell that is not a pAPC.

5 HOUSEKEEPING PROTEASOME – a proteasome normally active in peripheral cells, and generally not present or not strongly active in pAPCs.

IMMUNE PROTEASOME – a proteasome normally active in pAPCs; the immune proteasome is also active in some peripheral cells in infected tissues.

10 EPITOPE – a molecule or substance capable of stimulating an immune response. In preferred embodiments, epitopes according to this definition include but are not necessarily limited to a polypeptide and a nucleic acid encoding a polypeptide, wherein the polypeptide is capable of stimulating an immune response. In other preferred embodiments, epitopes according to this definition include but are not necessarily limited to peptides presented on the surface of cells, the peptides being non-covalently bound to the binding cleft of class I MHC, such that they can
15 interact with T cell receptors.

MHC EPITOPE – a polypeptide having a known or predicted binding affinity for a mammalian class I or class II major histocompatibility complex (MHC) molecule.

HOUSEKEEPING EPITOPE – In a preferred embodiment, a housekeeping epitope is defined as a polypeptide fragment that is an MHC epitope, and that is displayed on a cell in which
20 housekeeping proteasomes are predominantly active. In another preferred embodiment, a housekeeping epitope is defined as a polypeptide containing a housekeeping epitope according to the foregoing definition, that is flanked by one to several additional amino acids. In another preferred embodiment, a housekeeping epitope is defined as a nucleic acid that encodes a housekeeping epitope according to the foregoing definitions.

25 IMMUNE EPITOPE – In a preferred embodiment, an immune epitope is defined as a polypeptide fragment that is an MHC epitope, and that is displayed on a cell in which immune proteasomes are predominantly active. In another preferred embodiment, an immune epitope is defined as a polypeptide containing an immune epitope according to the foregoing definition, that is flanked by one to several additional amino acids. In another preferred embodiment, an immune
30 epitope is defined as a polypeptide including an epitope cluster sequence, having at least two polypeptide sequences having a known or predicted affinity for a class I MHC. In yet another preferred embodiment, an immune epitope is defined as a nucleic acid that encodes an immune epitope according to any of the foregoing definitions.

TARGET CELL – a cell to be targeted by the vaccines and methods of the invention.
35 Examples of target cells according to this definition include but are not necessarily limited to: a

neoplastic cell and a cell harboring an intracellular parasite, such as, for example, a virus, a bacterium, or a protozoan.

TARGET-ASSOCIATED ANTIGEN (TAA) – a protein or polypeptide present in a target cell.

5 TUMOR-ASSOCIATED ANTIGENS (TuAA) – a TAA, wherein the target cell is a neoplastic cell.

HLA EPITOPE – a polypeptide having a known or predicted binding affinity for a human class I or class II HLA complex molecule.

10 ANTIBODY – a natural immunoglobulin (Ig), poly- or monoclonal, or any molecule composed in whole or in part of an Ig binding domain, whether derived biochemically or by use of recombinant DNA. Examples include *inter alia*, F(ab), single chain Fv, and Ig variable region-phage coat protein fusions.

15 ENCODE – an open-ended term such that a nucleic acid encoding a particular amino acid sequence can consist of codons specifying that (poly)peptide, but can also comprise additional sequences either translatable, or for the control of transcription, translation, or replication, or to facilitate manipulation of some host nucleic acid construct.

20 SUBSTANTIAL SIMILARITY – this term is used to refer to sequences that differ from a reference sequence in an inconsequential way as judged by examination of the sequence. Nucleic acid sequences encoding the same amino acid sequence are substantially similar despite differences in degenerate positions or modest differences in length or composition of any non-coding regions. Amino acid sequences differing only by conservative substitution or minor length variations are substantially similar. Additionally, amino acid sequences comprising housekeeping epitopes that differ in the number of N-terminal flanking residues, or immune epitopes and epitope clusters that differ in the number of flanking residues at either terminus, are substantially similar. Nucleic acids
25 that encode substantially similar amino acid sequences are themselves also substantially similar.

30 FUNCTIONAL SIMILARITY – this term is used to refer to sequences that differ from a reference sequence in an inconsequential way as judged by examination of a biological or biochemical property, although the sequences may not be substantially similar. For example, two nucleic acids can be useful as hybridization probes for the same sequence but encode differing amino acid sequences. Two peptides that induce cross-reactive CTL responses are functionally similar even if they differ by non-conservative amino acid substitutions (and thus do not meet the substantial similarity definition). Pairs of antibodies, or TCRs, that recognize the same epitope can be functionally similar to each other despite whatever structural differences exist. In testing for functional similarity of immunogenicity one would generally immunize with the “altered” antigen
35 and test the ability of the elicited response (Ab, CTL, cytokine production, etc.) to recognize the target antigen. Accordingly, two sequences may be designed to differ in certain respects while

retaining the same function. Such designed sequence variants are among the embodiments of the present invention.

Table 1A. SEQ ID NOS.* including epitopes in Examples 1-7, 13.

SEQ ID NO	IDENTITY	SEQUENCE
1	Tyr 207-216	FLPWHRLFLL
2	Tyrosinase protein	Accession number**: P14679
3	SSX-2 protein	Accession number: NP_003138
4	PSMA protein	Accession number: NP_004467
5	Tyrosinase cDNA	Accession number: NM_000372
6	SSX-2 cDNA	Accession number: NM_003147
7	PSMA cDNA	Accession number: NM_004476
8	Tyr 207-215	FLPWHRLFL
9	Tyr 208-216	LPWHRLFLL
10	SSX-2 31-68	YFSKEEWEKMKASEKIFYVYMKRKYEAMTKLGFKATLP
11	SSX-2 32-40	FSKEEWEKM
12	SSX-2 39-47	KMKASEKIF
13	SSX-2 40-48	MKASEKIFY
14	SSX-2 39-48	KMKASEKIFY
15	SSX-2 41-49	KASEKIFYV
16	SSX-2 40-49	MKASEKIFYV
17	SSX-2 41-50	KASEKIFYVY
18	SSX-2 42-49	ASEKIFYVY
19	SSX-2 53-61	RKYEAMTKL
20	SSX-2 52-61	KRKYEAMTKL
21	SSX-2 54-63	KYEAMTKLGF
22	SSX-2 55-63	YEAMTKLGF
23	SSX-2 56-63	EAMTKLGF
24	HBV18-27	FLPSDYFPSV
25	HLA-B44 binder	AEMGKYSFY
26	SSX-1 41-49	KYSEKISYV
27	SSX-3 41-49	KVSEKIVYV
28	SSX-4 41-49	KSSEKIVYV
29	SSX-5 41-49	KASEKIIYV
30	PSMA163-192	AFSPQGMPEGDLVYVNYARTEDFFKLERDM
31	PSMA 168-190	GMPEGDLVYVNYARTEDFFKLER
32	PSMA 169-177	MPEGDLVYV
33	PSMA 168-177	GMPEGDLVYV
34	PSMA 168-176	GMPEGDLVY
35	PSMA 167-176	QGMPEGDLVY
36	PSMA 169-176	MPEGDLVY

37	PSMA 171-179	EGDLVYVNY
38	PSMA 170-179	PEGDLVYVNY
39	PSMA 174-183	LVYVNYARTE
40	PSMA 177-185	VNYARTEDF
41	PSMA 176-185	YVNYARTEDF
42	PSMA 178-186	NYARTEDFF
43	PSMA 179-186	YARTEDFF
44	PSMA 181-189	RTEDFFKLE
45	PSMA 281-310	RGIAEAVGLPSIPVHPIGYYDAQKLEKMG
46	PSMA 283-307	IAEAVGLPSIPVHPIGYYDAQKLE
47	PSMA 289-297	LPSIPVHPI
48	PSMA 288-297	GLPSIPVHPI
49	PSMA 297-305	IGYYDAQKL
50	PSMA 296-305	PIGYYDAQKL
51	PSMA 291-299	SIPVHPIGY
52	PSMA 290-299	PSIPVHPIGY
53	PSMA 292-299	IPVHPIGY
54	PSMA 299-307	YYDAQKLE
55	PSMA454-481	SSIEGNYTLRVDCTPLMYSLVHLTKEL
56	PSMA 456-464	IEGNYTLRV
57	PSMA 455-464	SIEGNYTLRV
58	PSMA 457-464	EGNYTLRV
59	PSMA 461-469	TLRVDCTPL
60	PSMA 460-469	YTLRVDCTPL
61	PSMA 462-470	LRVDCTPLM
62	PSMA 463-471	RVDCTPLMY
63	PSMA 462-471	LRVDCTPLMY
64	PSMA653-687	FDKSNPIVLRMMNDQLMFLERAFIDPLGLPDRPFY
65	PSMA 660-681	VLRMMNDQLMFLERAFIDPLGL
66	PSMA 663-671	MMNDQLMFL
67	PSMA 662-671	RMMNDQLMFL
68	PSMA 662-670	RMMNDQLMF
69	Tyr 1-17	MLLAVLVYCLLWSFQTSA

Table 1B. SEQ ID NOS.* including epitopes in Examples 14 and 15.

SEQ ID NO	IDENTITY	SEQUENCE
70	GP100 protein	**Accession number: P40967
71	MAGE-1 protein	Accession number: P43355
72	MAGE-2 protein	Accession number: P43356
73	MAGE-3 protein	Accession number: P43357
74	NY-ESO-1 protein	Accession number: P78358
75	LAGE-1a protein	Accession number: CAA11116

76	LAGE-1b protein	Accession number: CAA11117
77	PRAME protein	Accession number: NP 006106
78	PSA protein	Accession number: P07288
79	PSCA protein	Accession number: O43653
80	GP100 cds	Accession number: U20093
81	MAGE-1 cds	Accession number: M77481
82	MAGE-2 cds	Accession number: L18920
83	MAGE-3 cds	Accession number: U03735
84	NY-ESO-1 cDNA	Accession number: U87459
85	PRAME cDNA	Accession number: NM 006115
86	PSA cDNA	Accession number: NM 001648
87	PSCA cDNA	Accession number: AF043498
88	GP100 630-638	LPHSSSHWL
89	GP100 629-638	QLPHSSSHWL
90	GP100 614-622	LIYRRRLMK
91	GP100 613-622	SLIYRRRLMK
92	GP100 615-622	IYRRRLMK
93	GP100 630-638	LPHSSSHWL
94	GP100 629-638	QLPHSSSHWL
95	MAGE-1 95-102	ESLFRAVI
96	MAGE-1 93-102	ILESFRVI
97	MAGE-1 93-101	ILESFRV
98	MAGE-1 92-101	CILESFRV
99	MAGE-1 92-100	CILESFR
100	MAGE-1 263-271	EFLWGPRAL
101	MAGE-1 264-271	FLWGPRAL
102	MAGE-1 264-273	FLWGPRALAE
103	MAGE-1 265-274	LWGPRALAE
104	MAGE-1 268-276	PRALAETSY
105	MAGE-1 267-276	GPRALAETSY
106	MAGE-1 269-277	RALAETSYV
107	MAGE-1 271-279	LAETSYVKV
108	MAGE-1 270-279	ALAETSYVKV
109	MAGE-1 272-280	AETSYVKVL
110	MAGE-1 271-280	LAETSYVKVL
111	MAGE-1 274-282	TSYVKVLEY
112	MAGE-1 273-282	ETSYVKVLEY
113	MAGE-1 278-286	KVLEYVIKV
114	MAGE-1 168-177	SYVLVTCLGL
115	MAGE-1 169-177	YVLVTCLGL
116	MAGE-1 170-177	VLVTCLGL
117	MAGE-1 240-248	TQDLVQEKY

118	MAGE-1 239-248	LTQDLVQEKY
119	MAGE-1 232-240	YGEPRKLLT
120	MAGE-1 243-251	LVQEKYLEY
121	MAGE-1 242-251	DLVQEKYLEY
122	MAGE-1 230-238	SAYGEPRKL
123	MAGE-1 278-286	KVLEYVIKV
124	MAGE-1 277-286	VKVLEYVIKV
125	MAGE-1 276-284	YVKVLEYVI
126	MAGE-1 274-282	TSYVKVLEY
127	MAGE-1 273-282	ETSYVKVLEY
128	MAGE-1 283-291	VIKVSARVR
129	MAGE-1 282-291	YVIKVSARVR
130	MAGE-2 115-122	ELVHFLLL
131	MAGE-2 113-122	MVELVHFLLL
132	MAGE-2 109-116	ISRKMVEL
133	MAGE-2 108-116	AIKRKMVEL
134	MAGE-2 107-116	AAISRKMVEL
135	MAGE-2 112-120	KMVELVHFL
136	MAGE-2 109-117	ISRKMVELV
137	MAGE-2 108-117	AIKRKMVELV
138	MAGE-2 116-124	LVHFLLLKY
139	MAGE-2 115-124	ELVHFLLLKY
140	MAGE-2 111-119	RKMVELVHF
141	MAGE-2 158-166	LQLVFGIEV
142	MAGE-2 157-166	YLQLVFGIEV
143	MAGE-2 159-167	QLVFGIEVV
144	MAGE-2 158-167	LQLVFGIEVV
145	MAGE-2 164-172	IEVVEVVPI
146	MAGE-2 163-172	GIEVVEVVPI
147	MAGE-2 162-170	FGIEVVEVV
148	MAGE-2 154-162	ASEYLQLVF
149	MAGE-2 153-162	KASEYLQLVF
150	MAGE-2 218-225	EEKIWEEL
151	MAGE-2 216-225	APEEKIWEEL
152	MAGE-2 216-223	APEEKIWE
153	MAGE-2 220-228	KIWEELSML
154	MAGE-2 219-228	EKIWEELSML
155	MAGE-2 271-278	FLWGPRAL
156	MAGE-2 271-279	FLWGPRALI
157	MAGE-2 278-286	LIETSYVKV
158	MAGE-2 277-286	ALIETSYVKV
159	MAGE-2 276-284	RALIETSYV

160	MAGE-2 279-287	LIETSYVKVL
161	MAGE-2 278-287	LIETSYVKVL
162	MAGE-3 271-278	FLWGPRAL
163	MAGE-3 270-278	EFLWGPRAL
164	MAGE-3 271-279	FLWGPRALV
165	MAGE-3 276-284	RALVETSYV
166	MAGE-3 272-280	LWGPRALVE
167	MAGE-3 271-280	FLWGPRALVE
168	MAGE-3 272-281	LWGPRALVET
169	NY-ESO-1 82-90	GPESRLLEF
170	NY-ESO-1 83-91	PESRLLEFY
171	NY-ESO-1 82-91	GPESRLLEFY
172	NY-ESO-1 84-92	ESRLLEFYL
173	NY-ESO-1 86-94	RLLEFYLAM
174	NY-ESO-1 88-96	LEFYLAMPF
175	NY-ESO-1 87-96	LLEFYLAMPF
176	NY-ESO-1 93-102	AMPFATPMEA
177	NY-ESO-1 94-102	MPFATPMEA
178	NY-ESO-1 115-123	PLPVPGVLL
179	NY-ESO-1 114-123	PPLPVPGVLL
180	NY-ESO-1 116-123	LPVPGVLL
181	NY-ESO-1 103-112	ELARRSLAQD
182	NY-ESO-1 118-126	VPGVLLKEF
183	NY-ESO-1 117-126	PVPGVLLKEF
184	NY-ESO-1 116-123	LPVPGVLL
185	NY-ESO-1 127-135	TVSGNILT
186	NY-ESO-1 126-135	FTVSGNILT
187	NY-ESO-1 120-128	GVLLKEFTV
188	NY-ESO-1 121-130	VLLKEFTVSG
189	NY-ESO-1 122-130	LLKEFTVSG
190	NY-ESO-1 118-126	VPGVLLKEF
191	NY-ESO-1 117-126	PVPGVLLKEF
192	NY-ESO-1 139-147	AADHRQLQL
193	NY-ESO-1 148-156	SISSCLQQL
194	NY-ESO-1 147-156	LSISSCLQQL
195	NY-ESO-1 138-147	TAADHRQLQL
196	NY-ESO-1 161-169	WITQCFLPV
197	NY-ESO-1 157-165	SLLMWITQC
198	NY-ESO-1 150-158	SSCLQQLSL
199	NY-ESO-1 154-162	QQLSLLMWI
200	NY-ESO-1 151-159	SCLQQLSLL
201	NY-ESO-1 150-159	SSCLQQLSLL

202	NY-ESO-1 163-171	TQCFLPVFL
203	NY-ESO-1 162-171	TTQCFLPVFL
204	PRAME 219-227	PMQDIKMIL
205	PRAME 218-227	MPMQDIKMIL
206	PRAME 428-436	QHLIGLSNL
207	PRAME 427-436	LQHLIGLSNL
208	PRAME 429-436	HLIGLSNL
209	PRAME 431-439	IGLSNLTHV
210	PRAME 430-439	LIGLSNLTHV
211	PSA 53-61	VLVHPQWVL
212	PSA 52-61	GVLVHPQWVL
213	PSA 52-60	GVLVHPQWV
214	PSA 59-67	WVLTAACHCI
215	PSA 54-63	LVHPQWVLTA
216	PSA 53-62	VLVHPQWVLT
217	PSA 54-62	LVHPQWVLT
218	PSA 66-73	CIRNKSVI
219	PSA 65-73	HCIRNKSVI
220	PSA 56-64	HPQWVLTA
221	PSA 63-72	AAHCIRNKSV
222	PSCA 116-123	LLWGPGQL
223	PSCA 115-123	LLLWGPGQL
224	PSCA 114-123	GLLLWGPGQL
225	PSCA 99-107	ALQPAAAIL
226	PSCA 98-107	HALQPAAAIL
227	Tyr 128-137	APEKDKFFAY
228	Tyr 129-137	PEKDKFFAY
229	Tyr 130-138	EKDKFFAYL
230	Tyr 131-138	KDKFFAYL
231	Tyr 205-213	PAFLPWHRL
232	Tyr 204-213	APAFLPWHRL
233	Tyr 214-223	FLLRWEQEIQ
234	Tyr 212-220	RLFLLRWEQ
235	Tyr 191-200	GSEIWRDIDF
236	Tyr 192-200	SEIWRDIDF
237	Tyr 473-481	RIWSWLLGA
238	Tyr 476-484	SWLLGAAMV
239	Tyr 477-486	WLLGAAMVGA
240	Tyr 478-486	LLGAAMVGA
241	PSMA 4-12	LLHETDSAV
242	PSMA 13-21	ATARRPRWL
243	PSMA 53-61	TPKHNMKAF

244	PSMA 64-73	ELKAENKKF
245	PSMA 69-77	NIKKFLH ¹ NF
246	PSMA 68-77	ENIKKFLH ¹ NF
247	PSMA 220-228	AGAKGVILY
248	PSMA 468-477	PLMYSLVHNL
249	PSMA 469-477	LMYSLVHNL
250	PSMA 463-471	RVDCTPLMY
251	PSMA 465-473	DCTPLMYSL
252	PSMA 507-515	SGMPRISKL
253	PSMA 506-515	FSGMPRISKL
254	NY-ESO-1 136-163	RLTAADHRQLQLSISSCLQQLSLLMWIT
255	NY-ESO-1 150-177	SSCLQQLSLLMWITQCFLPVFLAQPPSG

¹ This H was reported as Y in the SWISSPROT database.

Table 1C. SEQ ID NOS.* including epitopes in Example 14.

SEQ ID NO.	IDENTITY	SEQUENCE
256	Mage-1 125-132	KAEMLESV
257	Mage-1 124-132	TKAEMLESV
258	Mage-1 123-132	VTKAEMLESV
259	Mage-1 128-136	MLESVIKNY
260	Mage-1 127-136	EMLESVIKNY
261	Mage-1 125-133	KAEMLESVI
262	Mage-1 146-153	KASESLQL
263	Mage-1 145-153	GKASESLQL
264	Mage-1 147-155	ASESLQLVF
265	Mage-1 153-161	LVFGIDVKE
266	Mage-1 114-121	LLKYRARE
267	Mage-1 106-113	VADLVGFL
268	Mage-1 105-113	KVADLVGFL
269	Mage-1 107-115	ADLVGFLLL
270	Mage-1 106-115	VADLVGFLLL
271	Mage-1 114-123	LLKYRAREPV
272	Mage-3 278-286	LVETSYVKV
273	Mage-3 277-286	ALVETSYVKV
274	Mage-3 285-293	KVLHHMVKI
275	Mage-3 283-291	YVKVLHHMV
276	Mage-3 275-283	PRALVETSY
277	Mage-3 274-283	GPRALVETSY
278	Mage-3 278-287	LVETSYVKVL
279	ED-B 4'-5	TIPEVPQL
280	ED-B 5'-5	DTIPEVPQL
281	ED-B 1-10	EVPQLTDLF
282	ED-B 23-30	TPLNSSTI
283	ED-B 18-25	IGLRWTPL
284	ED-B 17-25	SIGLRWTPL
285	ED-B 25-33	LNSSTIIGY
286	ED-B 24-33	PLNSSTIIGY

287	ED-B 23-31	TPLNSSTII
288	ED-B 31-38	IGYRITVV
289	ED-B 30-38	IIGYRITVV
290	ED-B 29-38	TIIGYRITVV
291	ED-B 31-39	IGYRITVVA
292	ED-B 30-39	IIGYRITVVA
293	CEA 184-191	SLPVSPRL
294	CEA 183-191	QSLPVSPRL
295	CEA 186-193	PVSPRLQL
296	CEA 185-193	LPVSPRLQL
297	CEA 184-193	SLPVSPRLQL
298	CEA 185-192	LPVSPRLQ
299	CEA 192-200	QLSNGNRTL
300	CEA 191-200	LQLSNGNRTL
301	CEA 179-187	WVNNQSLPV
302	CEA 186-194	PVSPRLQLS
303	CEA 362-369	SLPVSPRL
304	CEA 361-369	QSLPVSPRL
305	CEA 364-371	PVSPRLQL
306	CEA 363-371	LPVSPRLQL
307	CEA 362-371	SLPVSPRLQL
308	CEA 363-370	LPVSPRLQ
309	CEA 370-378	QLSNDNRTL
310	CEA 369-378	LQLSNDNRTL
311	CEA 357-365	WVNNQSLPV
312	CEA 360-368	NQSLPVSPR
313	CEA 540-547	SLPVSPRL
314	CEA 539-547	QSLPVSPRL
315	CEA 542-549	PVSPRLQL
316	CEA 541-549	LPVSPRLQL
317	CEA 540-549	SLPVSPRLQL
318	CEA 541-548	LPVSPRLQ
319	CEA 548-556	QLSNGNRTL
320	CEA 547-556	LQLSNGNRTL
321	CEA 535-543	WVNGQSLPV
322	CEA 533-541	LWWVNGQSL
323	CEA 532-541	YLWWVNGQSL
324	CEA 538-546	GQSLPVSPR
325	Her-2 30-37	DMKLRLPA
326	Her-2 28-37	GTDMLRLPA
327	Her-2 42-49	HLDMLRHL
328	Her-2 41-49	THLDMLRHL
329	Her-2 40-49	ETHLDMLRHL
330	Her-2 36-43	PASPETHL
331	Her-2 35-43	LPASPETHL
332	Her-2 34-43	RLPASPETHL
333	Her-2 38-46	SPETHDML
334	Her-2 37-46	ASPETHDML
335	Her-2 42-50	HLDMLRHLY
336	Her-2 41-50	THLDMLRHLY
337	Her-2 719-726	ELRKVKVL

338	Her-2 718-726	TELKVKVL
339	Her-2 717-726	ETELRKVKVL
340	Her-2 715-723	LKETELRKV
341	Her-2 714-723	ILKETELRKV
342	Her-2 712-720	MRILKETEL
343	Her-2 711-720	QMRILKETEL
344	Her-2 717-725	ETELRKVKV
345	Her-2 716-725	KETELRKVKV
346	Her-2 706-714	MPNQAQMRI
347	Her-2 705-714	AMPNQAQMRI
348	Her-2 706-715	MPNQAQMRL
349	HER-2 966-973	RPRFRELV
350	HER-2 965-973	CRPRFRELV
351	HER-2 968-976	RFRELVSEF
352	HER-2 967-976	PRFRELVSEF
353	HER-2 964-972	ECRPRFREL
354	NY-ESO-1 67-75	GAASGLNGC
355	NY-ESO-1 52-60	RASGPGGGA
356	NY-ESO-1 64-72	PHGGAASGL
357	NY-ESO-1 63-72	GPHGGAASGL
358	NY-ESO-1 60-69	APRGPHGGAA
359	PRAME 112-119	VRPRRWKL
360	PRAME 111-119	EVRPRRWKL
361	PRAME 113-121	RPRRWKLQV
362	PRAME 114-122	PRRWKLQVL
363	PRAME 113-122	RPRRWKLQVL
364	PRAME 116-124	RWKLQVLDL
365	PRAME 115-124	RRWKLQVLDL
366	PRAME 174-182	PVEVLVDF
367	PRAME 199-206	VKRKKNVL
368	PRAME 198-206	KVKRKKKNVL
369	PRAME 197-206	EKVKRKKKNVL
370	PRAME 198-205	KVKRKKKNV
371	PRAME 201-208	RKKNVLRL
372	PRAME 200-208	KRKKKNVLRL
373	PRAME 199-208	VKRKKKNVLRL
374	PRAME 189-196	DELFSLI
375	PRAME 205-213	VLRLCCKKL
376	PRAME 204-213	NVLRCLCCKKL
377	PRAME 194-202	YLIEKVKRK
378	PRAME 74-81	QAWPFTCL
379	PRAME 73-81	VQAWPFTCL
380	PRAME 72-81	MVQAWPFTCL
381	PRAME 81-88	LPLGVLMK
382	PRAME 80-88	CLPLGVLMK
383	PRAME 79-88	TCLPLGVLMK
384	PRAME 84-92	GVLKMGQHL
385	PRAME 81-89	LPLGVLMKG
386	PRAME 80-89	CLPLGVLMKG
387	PRAME 76-85	WPFTCLPLGV
388	PRAME 51-59	ELFPPLFMA

389	PRAME 49-57	PRELFPPLF
390	PRAME 48-57	LPRELFPPLF
391	PRAME 50-58	RELFPPLFM
392	PRAME 49-58	PRELFPPLFM
393	PSA 239-246	RPSLYTKV
394	PSA 238-246	ERPSLYTKV
395	PSA 236-243	LPERPSLY
396	PSA 235-243	ALPERPSLY
397	PSA 241-249	SLYTKVVHY
398	PSA 240-249	PSLYTKVVHY
399	PSA 239-247	RPSLYTKVV
400	PSMA 211-218	GNKVKNAQ
401	PSMA 202-209	IARYGKVF
402	PSMA 217-225	AQLAGAKGV
403	PSMA 207-215	KVFRGNKVK
404	PSMA 211-219	GNKVKNAQL
405	PSMA 269-277	TPGYPANEY
406	PSMA 268-277	LTPGYPANEY
407	PSMA 271-279	GYPANEYAY
408	PSMA 270-279	PGYPANEYAY
409	PSMA 266-274	DPLTPGYPA
410	PSMA 492-500	SLYESWTKK
411	PSMA 491-500	KSLYESWTKK
412	PSMA 486-494	EGFEGKSLY
413	PSMA 485-494	DEGFEGKSLY
414	PSMA 498-506	TKKSPSPEF
415	PSMA 497-506	WTKKSPSPEF
416	PSMA 492-501	SLYESWTKKS
417	PSMA 725-732	WGEVKRQI
418	PSMA 724-732	AWGEVKRQI
419	PSMA 723-732	KAWGEVKRQI
420	PSMA 723-730	KAWGEVKR
421	PSMA 722-730	SKAWGEVKR
422	PSMA 731-739	QIYVAAFTV
423	PSMA 733-741	YVAAFTVQA
424	PSMA 725-733	WGEVKRQIY
425	PSMA 727-735	EVKRQIYVA
426	PSMA 738-746	TVQAAAETL
427	PSMA 737-746	FTVQAAAETL
428	PSMA 729-737	KRQIYVAAF
429	PSMA 721-729	PSKAWGEVK
430	PSMA 723-731	KAWGEVKRQ
431	PSMA 100-108	WKEFGLDSV
432	PSMA 99-108	QWKEFGLDSV
433	PSMA 102-111	BFGLDSVELA
434	SCP-1 126-134	ELRQKESKL
435	SCP-1 125-134	AELRQKESKL
436	SCP-1 133-141	KLQENRKII
437	SCP-1 298-305	QLEEKTKL
438	SCP-1 297-305	NQLEEKTKL
439	SCP-1 288-296	LLEESRDKV

440	SCP-1 287-296	FLLEESRDKV
441	SCP-1 291-299	ESRDKVNQL
442	SCP-1 290-299	EESRDKVNQL
443	SCP-1 475-483	EKEVHDLEY
444	SCP-1 474-483	REKEVHDLEY
445	SCP-1 480-488	DLEYSYCHY
446	SCP-1 477-485	EVHDLEYSY
447	SCP-1 477-486	EVHDLEYSYC
448	SCP-1 502-509	KLSSKREL
449	SCP-1 508-515	ELKNTEYF
450	SCP-1 507-515	RELKNTEYF
451	SCP-1 496-503	KRGQRPKL
452	SCP-1 494-503	LPKRGQRPKL
453	SCP-1 509-517	LKNTEYFTL
454	SCP-1 508-517	ELKNTEYFTL
455	SCP-1 506-514	KRELKNTEY
456	SCP-1 502-510	KLSSKRELK
457	SCP-1 498-506	GQRPKLSSK
458	SCP-1 497-506	RGQRPKLSSK
459	SCP-1 500-508	RPKLSSKRE
460	SCP-1 573-580	LEYVREEL
461	SCP-1 572-580	ELEYVREEL
462	SCP-1 571-580	NELEYVREEL
463	SCP-1 579-587	ELKQKREDEV
464	SCP-1 575-583	YVREELKQK
465	SCP-1 632-640	QLNVYEIKV
466	SCP-1 630-638	SKQLNVYEI
467	SCP-1 628-636	AESKQLNVY
468	SCP-1 627-636	TAESKQLNVY
469	SCP-1 638-645	IKVKNLEL
470	SCP-1 637-645	EIKVKNLEL
471	SCP-1 636-645	YEIKVKNLEL
472	SCP-1 642-650	KLELELESA
473	SCP-1 635-643	VYEIKVNKL
474	SCP-1 634-643	NVYEIKVNKL
475	SCP-1 646-654	ELESAKQKF
476	SCP-1 642-650	KLELELESA
477	SCP-1 646-654	ELESAKQKF
478	SCP-1 771-778	KEKCLKREA
479	SCP-1 777-785	EAKENTATL
480	SCP-1 776-785	REAKENTATL
481	SCP-1 773-782	KLKREAKENT
482	SCP-1 112-119	EAEKIKKW
483	SCP-1 101-109	GLSRVYSKL
484	SCP-1 100-109	EGLSRVYSKL
485	SCP-1 108-116	KLYKEAEKI
486	SCP-1 98-106	NSEGLSRVY
487	SCP-1 97-106	ENSEGLSRVY
488	SCP-1 102-110	LSRVYSKLY
489	SCP-1 101-110	GLSRVYSKLY
490	SCP-1 96-105	LENSEGLSRV

491	SCP-1 108-117	KLYKEAEKIK
492	SCP-1 949-956	REDRWAVI
493	SCP-1 948-956	MRREDRWAVI
494	SCP-1 947-956	KMREDRWAVI
495	SCP-1 947-955	KMREDRWAV
496	SCP-1 934-942	TTPGSTLKF
497	SCP-1 933-942	LTPGSTLKF
498	SCP-1 937-945	GSTLKGAI
499	SCP-1 945-953	IRKMREDRW
500	SCP-1 236-243	RLEMHFKL
501	SCP-1 235-243	SRLEMHFKL
502	SCP-1 242-250	KLKEDYEKI
503	SCP-1 249-257	KIQHLEQEY
504	SCP-1 248-257	EKIQHLEQEY
505	SCP-1 233-242	ENSRLEMHF
506	SCP-1 236-245	RLEMHFKLKE
507	SCP-1 324-331	LEDIKVSL
508	SCP-1 323-331	ELEDIKVSL
509	SCP-1 322-331	KELEDIKVSL
510	SCP-1 320-327	LTKLEDI
511	SCP-1 319-327	HLTKLEDI
512	SCP-1 330-338	SLQRSVSTQ
513	SCP-1 321-329	TKELEDIKV
514	SCP-1 320-329	LTKLEEDIKV
515	SCP-1 326-335	DIKVSLSQSV
516	SCP-1 281-288	KMKDLTFL
517	SCP-1 280-288	NKMKDLTFL
518	SCP-1 279-288	ENKMKDLTFL
519	SCP-1 288-296	LLEESRDKV
520	SCP-1 287-296	FLLEESRDKV
521	SCP-1 291-299	ESRDKVNQL
522	SCP-1 290-299	EESRDKVNQL
523	SCP-1 277-285	EKENKMKDL
524	SCP-1 276-285	TEKENKMKDL
525	SCP-1 279-287	ENKMKDLTF
526	SCP-1 218-225	IEKMITAF
527	SCP-1 217-225	NIEKMITAF
528	SCP-1 216-225	SNIEKMITAF
529	SCP-1 223-230	TAFEELRV
530	SCP-1 222-230	ITAFEELRV
531	SCP-1 221-230	MITAFEELRV
532	SCP-1 220-228	KMITAFEEL
533	SCP-1 219-228	EKMITAFEEL
534	SCP-1 227-235	ELRVQAENS
535	SCP-1 213-222	DLNSNIEKMI
536	SCP-1 837-844	WTSAKNTL
537	SCP-1 846-854	TPLPKAYTV
538	SCP-1 845-854	STPLPKAYTV
539	SCP-1 844-852	LSTPLPKAY
540	SCP-1 843-852	TLSTPLPKAY
541	SCP-1 842-850	NTLSTPLPK

542	SCP-1 841-850	KNTLSTPLPK
543	SCP-1 828-835	ISKDKRDY
544	SCP-1 826-835	HGISKDKRDY
545	SCP-1 832-840	KRDYLWTS
546	SCP-1 829-838	SKDKRDYLWT
547	SCP-1 279-286	ENKMKDLT
548	SCP-1 260-268	EINDKEKQV
549	SCP-1 274-282	QITEKENKM
550	SCP-1 269-277	SLLLIQITE
551	SCP-1 453-460	FEKIAEEL
552	SCP-1 452-460	QFEKIAEEL
553	SCP-1 451-460	KQFEKIAEEL
554	SCP-1 449-456	DNKQFEKI
555	SCP-1 448-456	YDNKQFEKI
556	SCP-1 447-456	LYDNKQFEKI
557	SCP-1 440-447	LGEKETLL
558	SCP-1 439-447	VLGEKETLL
559	SCP-1 438-447	KVLGEKETLL
560	SCP-1 390-398	LLRTEQQRL
561	SCP-1 389-398	ELLRTEQQRL
562	SCP-1 393-401	TEQQRL
563	SCP-1 392-401	RTEQQRL
564	SCP-1 402-410	EDQLILTM
565	SCP-1 397-406	RLENYEDQLI
566	SCP-1 368-375	KARAAHSF
567	SCP-1 376-384	VVTEFETTV
568	SCP-1 375-384	FVVTEFETTV
569	SCP-1 377-385	VTEFETTV
570	SCP-1 376-385	VVTEFETTV
571	SCP-1 344-352	DLQIATNTI
572	SCP-1 347-355	IATNTICQL
573	SCP-1 346-355	QIATNTICQL
574	SSX4 57-65	VMTKLGFKY
575	SSX4 53-61	LNVEVMTKL
576	SSX4 52-61	KLNVEMTKL
577	SSX4 66-74	TLPPFMRSK
578	SSX4 110-118	KIMPKKPAE
579	SSX4 103-112	SLQRIFPKIM
580	Tyr 463-471	YIKSYLEQA
581	Tyr 459-467	SFQDYIKSY
582	Tyr 458-467	DSFQDYIKSY
583	Tyr 507-514	LPEEKQPL
584	Tyr 506-514	QLPEEKQPL
585	Tyr 505-514	KQLPEEKQPL
586	Tyr 507-515	LPEEKQPLL
587	Tyr 506-515	QLPEEKQPLL
588	Tyr 497-505	SLLCRHKRK
589	ED-B domain of Fibronectin	EVQQLTDLFSVDITDSSIGLRWTPLNSSITIGYRI TVVAAGEGIPIFEDFVDSSVGYTGTGLEPGID YDISVITLINGGESAPTTLTQQT
590	ED-B domain of	CTFDNLSPGLEYNVSVYTVKDDKESVPISDTIP

	Fibronectin with flanking sequence from Fribronectin	EVPQLTDLFSFVDITDSSIGLRWTPPLNSSTIIGYRI TVVAAGEGIPIFEDFVDSSVGYYTGTGLEPGID YDISVITLINGGESAPTTLTQQT AVPPPTDLRFTNIGPDTMRVTW
591	ED-B domain of Fibronectin cds	Accession number: X07717
592	CEA protein	Accession number: P06731
593	CEA cDNA	Accession number: NM_004363
594	Her2/Neu protein	Accession number: P04626
595	Her2/Neu cDNA	Accession number: M11730
596	SCP-1 protein	Accession number: Q15431
597	SCP-1 cDNA	Accession number: X95654
598	SSX-4 protein	Accession number: O60224
599	SSX-4 cDNA	Accession number: NM_005636

*Any of SEQ ID NOS. 1, 8, 9, 11-23, 26-29, 32-44, 47-54, 56-63, 66-68 88-253, and 256-588 can be useful as epitopes in any of the various embodiments of the invention. Any of SEQ ID NOS. 10, 30, 31, 45, 46, 55, 64, 65, 69, 254, and 255 can be useful as sequences containing epitopes or epitope clusters, as described in various embodiments of the invention.

**All accession numbers used here and throughout can be accessed through the NCBI databases, for example, through the Entrez seek and retrieval system on the world wide web.

Note that the following discussion sets forth the inventors' understanding of the operation of the invention. However, it is not intended that this discussion limit the patent to any particular theory of operation not set forth in the claims.

In pursuing the development of epitope vaccines others have generated lists of predicted epitopes based on MHC binding motifs. Such peptides can be immunogenic, but may not correspond to any naturally produced antigenic fragment so that whole antigen will not elicit a similar response or sensitize a target cell to cytolysis by CTL. Therefore such lists do not differentiate between those sequences that can be useful as vaccines and those that cannot. Efforts to determine which of these predicted epitopes are in fact naturally produced have often relied on screening their reactivity with tumor infiltrating lymphocytes (TIL). However, TIL are strongly biased to recognize immune epitopes whereas tumors (and chronically infected cells) will generally present housekeeping epitopes. Thus, unless the epitope is produced by both the housekeeping and immuno- proteasomes, the target cell will generally not be recognized by CTL induced with TIL-identified epitopes. The epitopes of the present invention, in contrast, are generated by the action a specified proteasome, indicating that they can be naturally produced, and enabling their appropriate use. The importance of the distinction between housekeeping and immune epitopes to vaccine design in more fully set forth in PCT publication WO 01/82963A2.

The epitopes of the invention include or encode polypeptide fragments of TAAs that are precursors or products of proteasomal cleavage by a housekeeping or immune proteasome, and that have known or predicted affinity for at least one allele of MHC I. In some embodiments, the epitopes include or encode a polypeptide of about 6 to 25 amino acids in length, preferably about 7

to 20 amino acids in length, more preferably about 8 to 15 amino acids in length, and still more preferably 9 or 10 amino acids in length. However, it is understood that the polypeptides can be larger as long as they do not contain sequences that cause the polypeptides to be directed away from the proteasome or to be destroyed by the proteasome. For immune epitopes, if the larger peptides do not contain such sequences, they can be processed in the pAPC by the immune proteasome. Housekeeping epitopes may also be embedded in longer sequences provided that the sequence is adapted to facilitate liberation of the epitope's C-terminus by action of the immunoproteasome. The sequences of these epitopes can be subjected to computer analysis in order to calculate physical, biochemical, immunologic, or molecular genetic properties such as mass, isoelectric point, predicted mobility in electrophoresis, predicted binding to other MHC molecules, melting temperature of nucleic acid probes, reverse translations, similarity or homology to other sequences, and the like.

In constructing the polynucleotides encoding the polypeptide epitopes of the invention, the gene sequence of the associated TAA can be used, or the polynucleotide can be assembled from any of the corresponding codons. For a 10 amino acid epitope this can constitute on the order of 10^6 different sequences, depending on the particular amino acid composition. While large, this is a distinct and readily definable set representing a miniscule fraction of the $>10^{18}$ possible polynucleotides of this length, and thus in some embodiments, equivalents of a particular sequence disclosed herein encompass such distinct and readily definable variations on the listed sequence. In choosing a particular one of these sequences to use in a vaccine, considerations such as codon usage, self-complementarity, restriction sites, chemical stability, etc. can be used as will be apparent to one skilled in the art.

The invention contemplates producing peptide epitopes. Specifically these epitopes are derived from the sequence of a TAA, and have known or predicted affinity for at least one allele of MHC I. Such epitopes are typically identical to those produced on target cells or pAPCs.

Compositions Containing Active Epitopes

Embodiments of the present invention provide polypeptide compositions, including vaccines, therapeutics, diagnostics, pharmacological and pharmaceutical compositions. The various compositions include newly identified epitopes of TAAs, as well as variants of these epitopes. Other embodiments of the invention provide polynucleotides encoding the polypeptide epitopes of the invention. The invention further provides vectors for expression of the polypeptide epitopes for purification. In addition, the invention provides vectors for the expression of the polypeptide epitopes in an APC for use as an anti-tumor vaccine. Any of the epitopes or antigens, or nucleic acids encoding the same, from Table 1A can be used. Other embodiments relate to methods of making and using the various compositions.

A general architecture for a class I MHC-binding epitope can be described, and has been reviewed more extensively in Madden, D.R. *Annu. Rev. Immunol.* 13:587-622, 1995. Much of the binding energy arises from main chain contacts between conserved residues in the MHC molecule and the N- and C-termini of the peptide. Additional main chain contacts are made but vary among MHC alleles. Sequence specificity is conferred by side chain contacts of so-called anchor residues with pockets that, again, vary among MHC alleles. Anchor residues can be divided into primary and secondary. Primary anchor positions exhibit strong preferences for relatively well-defined sets of amino acid residues. Secondary positions show weaker and/or less well-defined preferences that can often be better described in terms of less favored, rather than more favored, residues. Additionally, residues in some secondary anchor positions are not always positioned to contact the pocket on the MHC molecule at all. Thus, a subset of peptides exists that bind to a particular MHC molecule and have a side chain-pocket contact at the position in question and another subset exists that show binding to the same MHC molecule that does not depend on the conformation the peptide assumes in the peptide-binding groove of the MHC molecule. The C-terminal residue (P₉) is preferably a primary anchor residue. For many of the better studied HLA molecules (e.g. A2, A68, B27, B7, B35, and B53) the second position (P2) is also an anchor residue. However, central anchor residues have also been observed including P3 and P5 in HLA-B8, as well as P5 and P₋₃ in the murine MHC molecules H-2D^b and H-2K^b, respectively. Since more stable binding will generally improve immunogenicity, anchor residues are preferably conserved or optimized in the design of variants, regardless of their position.

Because the anchor residues are generally located near the ends of the epitope, the peptide can buckle upward out of the peptide-binding groove allowing some variation in length. Epitopes ranging from 8-11 amino acids have been found for HLA-A68, and up to 13 amino acids for HLA-A2. In addition to length variation between the anchor positions, single residue truncations and extensions have been reported and the N- and C-termini, respectively. Of the non-anchor residues, some point up out of the groove, making no contact with the MHC molecule but being available to contact the TCR, very often P1, P4, and P₋₁ for HLA-A2. Others of the non-anchor residues can become interposed between the upper edges of the peptide-binding groove and the TCR, contacting both. The exact positioning of these side chain residues, and thus their effects on binding, MHC fine conformation, and ultimately immunogenicity, are highly sequence dependent. For an epitope to be highly immunogenic it must not only promote stable enough TCR binding for activation to occur, but the TCR must also have a high enough off-rate that multiple TCR molecules can interact sequentially with the same peptide-MHC complex (Kalergis, A.M. et al., *Nature Immunol.* 2:229-234, 2001). Thus without further information about the ternary complex, both conservative and non-conservative substitutions at these positions merit consideration when designing variants.

The polypeptide epitope variants can be made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations. Variants can be derived from substitution, deletion or insertion of one or more amino acids as compared with the native sequence. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a threonine with a serine. Such replacements are referred to as conservative amino acid replacements, and all appropriate conservative amino acid replacements are considered to be embodiments of one invention. Insertions or deletions can optionally be in the range of about 1 to 4, preferably 1 to 2, amino acids. It is generally preferable to maintain the "anchor positions" of the peptide which are responsible for binding to the MHC molecule in question. Indeed, immunogenicity of peptides can be improved in many cases by substituting more preferred residues at the anchor positions (Franco, et al., *Nature Immunology*, 1(2):145-150, 2000). Immunogenicity of a peptide can also often be improved by substituting bulkier amino acids for small amino acids found in non-anchor positions while maintaining sufficient cross-reactivity with the original epitope to constitute a useful vaccine. The variation allowed can be determined by routine insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the polypeptide epitope. Because the polypeptide epitope is often 9 amino acids, the substitutions preferably are made to the shortest active epitope, for example, an epitope of 9 amino acids.

Variants can also be made by adding any sequence onto the N-terminus of the polypeptide epitope variant. Such N-terminal additions can be from 1 amino acid up to at least 25 amino acids. Because peptide epitopes are often trimmed by N-terminal exopeptidases active in the pAPC, it is understood that variations in the added sequence can have no effect on the activity of the epitope. In preferred embodiments, the amino acid residues between the last upstream proteasomal cleavage site and the N-terminus of the MHC epitope do not include a proline residue. Serwold, T. et al., *Nature Immunol.* 2:644-651, 2001. Accordingly, effective epitopes can be generated from precursors larger than the preferred 9-mer class I motif.

Peptides are useful to the extent that they correspond to epitopes actually displayed by MHC I on the surface of a target cell or a pACP. A single peptide can have varying affinities for different MHC molecules, binding some well, others adequately, and still others not appreciably (Table 2). MHC alleles have traditionally been grouped according to serologic reactivity which does not reflect the structure of the peptide-binding groove, which can differ among different alleles of the same type. Similarly, binding properties can be shared across types; groups based on shared binding properties have been termed supertypes. There are numerous alleles of MHC I in the human population; epitopes specific to certain alleles can be selected based on the genotype of the patient.

Table 2.
Predicted Binding of Tyrosinase₂₀₇₋₂₁₆ (SEQ ID NO. 1) to Various MHC types

MHC I type	*Half time of dissociation (min)
A1	0.05
A*0201	1311.
A*0205	50.4
A3	2.7
A*1101 (part of the A3 supertype)	0.012
A24	6.0
B7	4.0
B8	8.0
B14 (part of the B27 supertype)	60.0
B*2702	0.9
B*2705	30.0
B*3501 (part of the B7 supertype)	2.0
B*4403	0.1
B*5101 (part of the B7 supertype)	26.0
B*5102	55.0
B*5801	0.20
B60	0.40
B62	2.0

*HLA Peptide Binding Predictions (internet [http:// access at bimas.dcrt.nih.gov/molbio/hla_bin](http://bimas.dcrt.nih.gov/molbio/hla_bin))

5

In further embodiments of the invention, the epitope, as peptide or encoding polynucleotide, can be administered as a vaccine or immunogenic composition, alone or in combination with various adjuvants, carriers, or excipients. It should be noted that although the term vaccine may be used herein, the discussion can be applied and used with any of the other compositions mentioned herein. Particularly advantageous adjuvants include various cytokines and oligonucleotides containing immunostimulatory sequences (as set forth in greater detail in the co-pending applications referenced herein). Additionally the polynucleotide encoded epitope can be contained in a virus (e.g. *vaccinia* or adenovirus) or in a microbial host cell (e.g. *Salmonella* or *Listeria monocytogenes*) which is then used as a vector for the polynucleotide (Dietrich, G. et al. Nat. Biotech. 16:181-185, 1998). Alternatively a pAPC can be transformed, *ex vivo*, to express the epitope, or pulsed with peptide epitope, to be itself administered as a vaccine. To increase efficiency of these processes, the encoded epitope can be carried by a viral or bacterial vector, or complexed with a ligand of a receptor found on pAPC. Similarly the peptide epitope can be complexed with or conjugated to a pAPC ligand. A vaccine can be composed of more than a single epitope.

10

15

20

Particularly advantageous strategies for incorporating epitopes, and combining them with epitope clusters, into a vaccine are disclosed in U.S. Patent Application No. 09/560,465 entitled "EPI TOPE SYNCHRONIZATION IN ANTIGEN PRESENTING CELLS," filed on April 28,

2000. Epitope clusters for use in connection with this invention are disclosed in U.S. Patent Application No. 09/561,571 entitled "EPITOPE CLUSTERS," filed on April 28, 2000.

Preferred embodiments of the present invention are directed to vaccines and methods for causing a pAPC or population of pAPCs to present housekeeping epitopes that correspond to the epitopes displayed on a particular target cell. Any of the epitopes or antigens in Table 1A, can be used for example. In one embodiment, the housekeeping epitope is a TuAA epitope processed by the housekeeping proteasome of a particular tumor type. In another embodiment, the housekeeping epitope is a virus-associated epitope processed by the housekeeping proteasome of a cell infected with a virus. This facilitates a specific T cell response to the target cells. Concurrent expression by the pAPCs of multiple epitopes, corresponding to different induction states (pre- and post-attack), can drive a CTL response effective against target cells as they display either housekeeping epitopes or immune epitopes.

By having both housekeeping and immune epitopes present on the pAPC, this embodiment can optimize the cytotoxic T cell response to a target cell. With dual epitope expression, the pAPCs can continue to sustain a CTL response to the immune-type epitope when the tumor cell switches from the housekeeping proteasome to the immune proteasome with induction by IFN, which, for example, may be produced by tumor-infiltrating CTLs.

In a preferred embodiment, immunization of a patient is with a vaccine that includes a housekeeping epitope. Many preferred TAAs are associated exclusively with a target cell, particularly in the case of infected cells. In another embodiment, many preferred TAAs are the result of deregulated gene expression in transformed cells, but are found also in tissues of the testis, ovaries and fetus. In another embodiment, useful TAAs are expressed at higher levels in the target cell than in other cells. In still other embodiments, TAAs are not differentially expressed in the target cell compare to other cells, but are still useful since they are involved in a particular function of the cell and differentiate the target cell from most other peripheral cells; in such embodiments, healthy cells also displaying the TAA may be collaterally attacked by the induced T cell response, but such collateral damage is considered to be far preferable to the condition caused by the target cell.

A preferred embodiment of the present invention includes a method of administering a vaccine including a housekeeping epitope to induce a therapeutic immune response. The vaccine is administered to a patient in a manner consistent with the standard vaccine delivery protocols that are well known in the art. Methods of administering epitopes of TAAs include, without limitation, transdermal, intranodal, perinodal, oral, intravenous, intradermal, intramuscular, intraperitoneal, and mucosal administration. A particularly useful method of vaccine delivery to elicit a CTL response is disclosed in PCT Publication No. WO 99/01283, entitled "A METHOD OF INDUCING A CTL RESPONSE," filed on July 10, 1998.

Because the epitope synchronization system has utility in inducing a cell mediated immune response, a vaccine to induce a specific T cell response to a target cell is likewise included in a preferred embodiment of the present invention. The vaccine contains a housekeeping epitope in a concentration effective to cause a pAPC or populations of pAPCs to display housekeeping epitopes. Advantageously, the vaccine can include a plurality of housekeeping epitopes or one or more housekeeping epitopes optionally in combination with one or more immune epitopes. Formulations of the vaccine contain peptides and/or nucleic acids in a concentration sufficient to cause pAPCs to present the epitopes. The formulations preferably contain epitopes in a total concentration of about 1µg-1mg/100µl of vaccine preparation. Conventional dosages and dosing for peptide vaccines and/or nucleic acid vaccines can be used with the present invention, and such dosing regimens are well understood in the art. In one embodiment, a single dosage for an adult human may advantageously be from about 1 to about 5000 µl of such a composition, administered one time or multiple times, e.g., in 2, 3, 4 or more dosages separated by 1 week, 2 weeks, 1 month, or more. insulin pump delivers 1 ul per hour (lowest frequency) ref intranodal method patent.

The compositions and methods of the invention disclosed herein further contemplate incorporating adjuvants into the formulations in order to enhance the performance of the vaccines. Specifically, the addition of adjuvants to the formulations is designed to enhance the delivery or uptake of the epitopes by the pAPCs. The adjuvants contemplated by the present invention are known by those of skill in the art and include, for example, GMCSF, GCSF, IL-2, IL-12, BCG, tetanus toxoid, osteopontin, and ETA-1.

In some embodiments of the invention, the vaccines can include a recombinant organism, such as a virus, bacterium or parasite, genetically engineered to express an epitope in a host. For example, *Listeria monocytogenes*, a gram-positive, facultative intracellular bacterium, is a potent vector for targeting TuAAs to the immune system. In a preferred embodiment, this vector can be engineered to express a housekeeping epitope to induce therapeutic responses. The normal route of infection of this organism is through the gut and can be delivered orally. In another embodiment, an adenovirus (Ad) vector encoding a housekeeping epitope for a TuAA can be used to induce anti-virus or anti-tumor responses. Bone marrow-derived dendritic cells can be transduced with the virus construct and then injected, or the virus can be delivered directly via subcutaneous injection into an animal to induce potent T-cell responses. Another embodiment employs a recombinant vaccinia virus engineered to encode amino acid sequences corresponding to a housekeeping epitope for a TAA. Vaccinia viruses carrying constructs with the appropriate nucleotide substitutions in the form of a minigene construct can direct the expression of a housekeeping epitope, leading to a therapeutic T cell response against the epitope.

The immunization with DNA requires that APCs take up the DNA and express the encoded proteins or peptides. It is possible to encode a discrete class I peptide on the DNA. By

immunizing with this construct, APCs can be caused to express a housekeeping epitope, which is then displayed on class I MHC on the surface of the cell for stimulating an appropriate CTL response. Constructs generally relying on termination of translation or non-proteasomal proteases for generation of proper termini of housekeeping epitopes have been described in U.S. Patent application No. 09/561,572 entitled EXPRESSION VECTORS ENCODING EPITOPES OF TARGET-ASSOCIATED ANTIGENS, filed on April 28, 2000.

It can be desirable to express housekeeping peptides in the context of a larger protein. Processing can be detected even when a small number of amino acids are present beyond the terminus of an epitope. Small peptide hormones are usually proteolytically processed from longer translation products, often in the size range of approximately 60-120 amino acids. This fact has led some to assume that this is the minimum size that can be efficiently translated. In some embodiments, the housekeeping peptide can be embedded in a translation product of at least about 60 amino acids. In other embodiments the housekeeping peptide can be embedded in a translation product of at least about 50, 30, or 15 amino acids.

Due to differential proteasomal processing, the immune proteasome of the pAPC produces peptides that are different from those produced by the housekeeping proteasome in peripheral body cells. Thus, in expressing a housekeeping peptide in the context of a larger protein, it is preferably expressed in the APC in a context other than its full length native sequence, because, as a housekeeping epitope, it is generally only efficiently processed from the native protein by the housekeeping proteasome, which is not active in the APC. In order to encode the housekeeping epitope in a DNA sequence encoding a larger protein, it is useful to find flanking areas on either side of the sequence encoding the epitope that permit appropriate cleavage by the immune proteasome in order to liberate that housekeeping epitope. Such a sequence ensuring epitope synchronization is referred to hereinafter as a SYNCHROTOPE™. Altering flanking amino acid residues at the N-terminus and C-terminus of the desired housekeeping epitope can facilitate appropriate cleavage and generation of the housekeeping epitope in the APC. Sequences embedding housekeeping epitopes can be designed *de novo* and screened to determine which can be successfully processed by immune proteasomes to liberate housekeeping epitopes.

Alternatively, another strategy is very effective for identifying sequences allowing production of housekeeping epitopes in APC. A contiguous sequence of amino acids can be generated from head to tail arrangement of one or more housekeeping epitopes. A construct expressing this sequence is used to immunize an animal, and the resulting T cell response is evaluated to determine its specificity to one or more of the epitopes in the array. By definition, these immune responses indicate housekeeping epitopes that are processed in the pAPC effectively. The necessary flanking areas around this epitope are thereby defined. The use of flanking regions of about 4-6 amino acids on either side of the desired peptide can provide the necessary

information to facilitate proteasome processing of the housekeeping epitope by the immune proteasome. Therefore, a SYNCHROTOPE™ of approximately 16-22 amino acids can be inserted into, or fused to, any protein sequence effectively to result in that housekeeping epitope being produced in an APC. In alternate embodiments the whole head-to-tail array of epitopes, or just the epitopes immediately adjacent to the correctly processed housekeeping epitope can be similarly transferred from a test construct to a vaccine vector.

In a preferred embodiment, the housekeeping epitopes can be embedded between known immune epitopes, or segments of such, thereby providing an appropriate context for processing. The abutment of housekeeping and immune epitopes can generate the necessary context to enable the immune proteasome to liberate the housekeeping epitope, or a larger fragment, preferably including a correct C-terminus. It can be useful to screen constructs to verify that the desired epitope is produced. The abutment of housekeeping epitopes can generate a site cleavable by the immune proteasome. Some embodiments of the invention employ known epitopes to flank housekeeping epitopes in test substrates; in others, screening as described below are used whether the flanking regions are arbitrary sequences or mutants of the natural flanking sequence, and whether or not knowledge of proteasomal cleavage preferences are used in designing the substrates.

Cleavage at the mature N-terminus of the epitope, while advantageous, is not required, since a variety of N-terminal trimming activities exist in the cell that can generate the mature N-terminus of the epitope subsequent to proteasomal processing. It is preferred that such N-terminal extension be less than about 25 amino acids in length and it is further preferred that the extension have few or no proline residues. Preferably, in screening, consideration is given not only to cleavage at the ends of the epitope (or at least at its C-terminus), but consideration also can be given to ensure limited cleavage within the epitope.

Shotgun approaches can be used in designing test substrates and can increase the efficiency of screening. In one embodiment multiple epitopes can be assembled one after the other, with individual epitopes possibly appearing more than once. The substrate can be screened to determine which epitopes can be produced. In the case where a particular epitope is of concern a substrate can be designed in which it appears in multiple different contexts. When a single epitope appearing in more than one context is liberated from the substrate additional secondary test substrates, in which individual instances of the epitope are removed, disabled, or are unique, can be used to determine which are being liberated and truly constitute SYNCHROTOPE™s.

Several readily practicable screens exist. A preferred *in vitro* screen utilizes proteasomal digestion analysis, using purified immune proteasomes, to determine if the desired housekeeping epitope can be liberated from a synthetic peptide embodying the sequence in question. The position of the cleavages obtained can be determined by techniques such as mass spectrometry, HPLC, and

N-terminal pool sequencing; as described in greater detail in U. S. Patent Applications entitled METHOD OF EPITOPE DISCOVERY, EPITOPE SYNCHRONIZATION IN ANTIGEN PRESENTING CELLS, two Provisional U. S. Patent Applications entitled EPITOPE SEQUENCES, which are all cited and incorporated by reference above.

5 Alternatively, *in vivo* screens such as immunization or target sensitization can be employed. For immunization a nucleic acid construct capable of expressing the sequence in question is used. Harvested CTL can be tested for their ability to recognize target cells presenting the housekeeping epitope in question. Such targets cells are most readily obtained by pulsing cells expressing the appropriate MHC molecule with synthetic peptide embodying the mature
10 housekeeping epitope. Alternatively, cells known to express housekeeping proteasome and the antigen from which the housekeeping epitope is derived, either endogenously or through genetic engineering, can be used. To use target sensitization as a screen, CTL, or preferably a CTL clone, that recognizes the housekeeping epitope can be used. In this case it is the target cell that expresses the embedded housekeeping epitope (instead of the pAPC during immunization) and it must
15 express immune proteasome. Generally, the target cell can be transformed with an appropriate nucleic acid construct to confer expression of the embedded housekeeping epitope. Loading with a synthetic peptide embodying the embedded epitope using peptide loaded liposomes or a protein transfer reagent such as BIOPORTER™ (Gene Therapy Systems, San Diego, CA) represents an alternative.

20 Additional guidance on nucleic acid constructs useful as vaccines in accordance with the present invention are disclosed in U.S. Patent Application No. 09/561,572 entitled "EXPRESSION VECTORS ENCODING EPITOPES OF TARGET-ASSOCIATED ANTIGENS," filed on April 28, 2000. Further, expression vectors and methods for their design, which are useful in accordance with the present invention are disclosed in U.S. Patent Application No. 60/336,968 (attorney
25 docket number CTLIMM.022PR) entitled "EXPRESSION VECTORS ENCODING EPITOPES OF TARGET-ASSOCIATED ANTIGENS AND METHODS FOR THEIR DESIGN," filed on 11/7/2001, which is incorporated by reference in its entirety.

 A preferred embodiment of the present invention includes a method of administering a vaccine including an epitope (or epitopes) to induce a therapeutic immune response. The vaccine
30 is administered to a patient in a manner consistent with the standard vaccine delivery protocols that are known in the art. Methods of administering epitopes of TAAs including, without limitation, transdermal, intranodal, perinodal, oral, intravenous, intradermal, intramuscular, intraperitoneal, and mucosal administration, including delivery by injection, instillation or inhalation. A particularly useful method of vaccine delivery to elicit a CTL response is disclosed in Australian
35 Patent No. 739189 issued January 17, 2002; U.S. Patent Application No. 09/380,534, filed on

September 1, 1999; and a Continuation-in-Part thereof U.S. Patent Application No. 09/776,232 both entitled "A METHOD OF INDUCING A CTL RESPONSE," filed on February 2, 2001.

Reagents Recognizing Epitopes

In another aspect of the invention, proteins with binding specificity for the epitope and the epitope-MHC molecule complex are contemplated, as well as the isolated cells by which they can be expressed. In one set of embodiments these reagents take the form of immunoglobulins: polyclonal sera or monoclonal antibodies (mAb), methods for the generation of which are well known in the art. Generation of mAb with specificity for peptide-MHC molecule complexes is known in the art. See, for example, Aharoni et al. *Nature* 351:147-150, 1991; Andersen et al. *Proc. Natl. Acad. Sci. USA* 93:1820-1824, 1996; Dadaglio et al. *Immunity* 6:727-738, 1997; Duc et al. *Int. Immunol.* 5:427-431, 1993; Eastman et al. *Eur. J. Immunol.* 26:385-393, 1996; Engberg et al. *Immunotechnology* 4:273-278, 1999; Porgdor et al. *Immunity* 6:715-726, 1997; Puri et al. *J. Immunol.* 158:2471-2476, 1997; and Polakova, K., et al. *J. Immunol.* 165 342-348, 2000; all of which are hereby incorporated by reference in their entirety.

In other embodiments the compositions can be used to induce and generate, *in vivo* and *in vitro*, T-cells specific for any of the epitopes, including those listed in Table 1A, for example. Thus, embodiments also relate to and include isolated T cells, T cell clones, T cell hybridomas, or a protein containing the T cell receptor (TCR) binding domain derived from the cloned gene, as well as a recombinant cell expressing such a protein. Such TCR derived proteins can be simply the extra-cellular domains of the TCR, or a fusion with portions of another protein to confer a desired property or function. One example of such a fusion is the attachment of TCR binding domains to the constant regions of an antibody molecule so as to create a divalent molecule. The construction and activity of molecules following this general pattern have been reported, for example, Plaksin, D. et al. *J. Immunol.* 158:2218-2227, 1997 and Lebowitz, M.S. et al. *Cell Immunol.* 192:175-184, 1999, which are hereby incorporated by reference in their entirety. The more general construction and use of such molecules is also treated in U.S. patent 5,830,755 entitled T CELL RECEPTORS AND THEIR USE IN THERAPEUTIC AND DIAGNOSTIC METHODS, which is hereby incorporated by reference in its entirety.

The generation of such T cells can be readily accomplished by standard immunization of laboratory animals, and reactivity to human target cells can be obtained by immunizing with human target cells or by immunizing HLA-transgenic animals with the antigen/epitope. For some therapeutic approaches T cells derived from the same species are desirable. While such a cell can be created by cloning, for example, a murine TCR into a human T cell as contemplated above, *in vitro* immunization of human cells offers a potentially faster option. Techniques for *in vitro* immunization, even using naive donors, are known in the field, for example, Stauss et al., *Proc. Natl. Acad. Sci. USA* 89:7871-7875, 1992; Salgaller et al. *Cancer Res.* 55:4972-4979, 1995; Tsai et

al., *J. Immunol.* 158:1796-1802, 1997; and Chung et al., *J. Immunother.* 22:279-287, 1999; which are hereby incorporated by reference in their entirety.

Any of these molecules can be conjugated to enzymes, radiochemicals, fluorescent tags, and toxins, so as to be used in the diagnosis (imaging or other detection), monitoring, and treatment of the pathogenic condition associated with the epitope. Thus a toxin conjugate can be administered to kill tumor cells, radiolabeling can facilitate imaging of epitope positive tumor, an enzyme conjugate can be used in an ELISA-like assay to diagnose cancer and confirm epitope expression in biopsied tissue. In a further embodiment, such T cells as set forth above, following expansion accomplished through stimulation with the epitope and/or cytokines, can be administered to a patient as an adoptive immunotherapy.

Reagents Comprising Epitopes

A further aspect of the invention provides isolated epitope-MHC complexes. In a particularly advantageous embodiment of this aspect of the invention, the complexes can be soluble, multimeric proteins such as those described in U. S. Patent No. 5,635,363 (tetramers) or U. S. Patent No. 6,015,884 (Ig-dimers), both of which are hereby incorporated by reference in their entirety. Such reagents are useful in detecting and monitoring specific T cell responses, and in purifying such T cells.

Isolated MHC molecules complexed with epitopic peptides can also be incorporated into planar lipid bilayers or liposomes. Such compositions can be used to stimulate T cells *in vitro* or, in the case of liposomes, *in vivo*. Co-stimulatory molecules (e.g. B7, CD40, LFA-3) can be incorporated into the same compositions or, especially for *in vitro* work, co-stimulation can be provided by anti-co-receptor antibodies (e.g. anti-CD28, anti-CD154, anti-CD2) or cytokines (e.g. IL-2, IL-12). Such stimulation of T cells can constitute vaccination, drive expansion of T cells *in vitro* for subsequent infusion in an immunotherapy, or constitute a step in an assay of T cell function.

The epitope, or more directly its complex with an MHC molecule, can be an important constituent of functional assays of antigen-specific T cells at either an activation or readout step or both. Of the many assays of T cell function current in the art (detailed procedures can be found in standard immunological references such as *Current Protocols in Immunology* 1999 John Wiley & Sons Inc., N.Y., which is hereby incorporated by reference in its entirety) two broad classes can be defined, those that measure the response of a pool of cells and those that measure the response of individual cells. Whereas the former conveys a global measure of the strength of a response, the latter allows determination of the relative frequency of responding cells. Examples of assays measuring global response are cytotoxicity assays, ELISA, and proliferation assays detecting cytokine secretion. Assays measuring the responses of individual cells (or small clones derived from them) include limiting dilution analysis (LDA), ELISPOT, flow cytometric detection of

unsecreted cytokine (described in U.S. Patent No. 5,445,939, entitled "METHOD FOR ASSESSMENT OF THE MONONUCLEAR LEUKOCYTE IMMUNE SYSTEM" and U.S. Patent Nos 5,656,446; and 5,843,689, both entitled "METHOD FOR THE ASSESSMENT OF THE MONONUCLEAR LEUKOCYTE IMMUNE SYSTEM," reagents for which are sold by Becton, Dickinson & Company under the tradename 'FASTIMMUNE', which patents are hereby incorporated by reference in their entirety) and detection of specific TCR with tetramers or Ig-dimers as stated and referenced above. The comparative virtues of these techniques have been reviewed in Yee, C. et al. *Current Opinion in Immunology*, 13:141-146, 2001, which is hereby incorporated by reference in its entirety. Additionally detection of a specific TCR rearrangement or expression can be accomplished through a variety of established nucleic acid based techniques, particularly in situ and single-cell PCR techniques, as will be apparent to one of skill in the art.

These functional assays are used to assess endogenous levels of immunity, response to an immunologic stimulus (e.g. a vaccine), and to monitor immune status through the course of a disease and treatment. Except when measuring endogenous levels of immunity, any of these assays presume a preliminary step of immunization, whether *in vivo* or *in vitro* depending on the nature of the issue being addressed. Such immunization can be carried out with the various embodiments of the invention described above or with other forms of immunogen (e.g., pAPC-tumor cell fusions) that can provoke similar immunity. With the exception of PCR and tetramer/Ig-dimer type analyses which can detect expression of the cognate TCR, these assays generally benefit from a step of *in vitro* antigenic stimulation which can advantageously use various embodiments of the invention as described above in order to detect the particular functional activity (highly cytolytic responses can sometimes be detected directly). Finally, detection of cytolytic activity requires epitope-displaying target cells, which can be generated using various embodiments of the invention. The particular embodiment chosen for any particular step depends on the question to be addressed, ease of use, cost, and the like, but the advantages of one embodiment over another for any particular set of circumstances will be apparent to one of skill in the art.

Tumor Associated Antigens

Epitopes of the present invention are derived from the TuAAs tyrosinase (SEQ ID NO. 2), SSX-2, (SEQ ID NO. 3), PSMA (prostate-specific membrane antigen) (SEQ ID NO. 4), GP100, (SEQ ID NO. 70), MAGE-1, (SEQ ID NO. 71), MAGE-2, (SEQ ID NO. 72), MAGE-3, (SEQ ID NO. 73), NY-ESO-1, (SEQ ID NO. 74), PRAME, (SEQ ID NO. 77), PSA, (SEQ ID NO. 78), and PSCA, (SEQ ID NO. 79). The natural coding sequences for these eleven proteins, or any segments within them, can be determined from their cDNA or complete coding (cds) sequences, SEQ ID NOS. 5-7, and 80-87, respectively.

Tyrosinase is a melanin biosynthetic enzyme that is considered one of the most specific markers of melanocytic differentiation. Tyrosinase is expressed in few cell types, primarily in

melanocytes, and high levels are often found in melanomas. The usefulness of tyrosinase as a TuAA is taught in U.S. Patent 5,747,271 entitled "METHOD FOR IDENTIFYING INDIVIDUALS SUFFERING FROM A CELLULAR ABNORMALITY SOME OF WHOSE ABNORMAL CELLS PRESENT COMPLEXES OF HLA-A2/TYROSINASE DERIVED PEPTIDES, AND METHODS FOR TREATING SAID INDIVIDUALS" which is hereby incorporated by reference in its entirety.

GP100, also known as PMel17, also is a melanin biosynthetic protein expressed at high levels in melanomas. GP100 as a TuAA is disclosed in U.S. Patent 5,844,075 entitled "MELANOMA ANTIGENS AND THEIR USE IN DIAGNOSTIC AND THERAPEUTIC METHODS," which is hereby incorporated by reference in its entirety.

SSX-2, also known as Hom-Mel-40, is a member of a family of highly conserved cancer-testis antigens (Gure, A.O. et al. *Int. J. Cancer* 72:965-971, 1997, which is hereby incorporated by reference in its entirety). Its identification as a TuAA is taught in U.S. Patent 6,025,191 entitled "ISOLATED NUCLEIC ACID MOLECULES WHICH ENCODE A MELANOMA SPECIFIC ANTIGEN AND USES THEREOF," which is hereby incorporated by reference in its entirety. Cancer-testis antigens are found in a variety of tumors, but are generally absent from normal adult tissues except testis. Expression of different members of the SSX family have been found variously in tumor cell lines. Due to the high degree of sequence identity among SSX family members, similar epitopes from more than one member of the family will be generated and able to bind to an MHC molecule, so that some vaccines directed against one member of this family can cross-react and be effective against other members of this family (see example 3 below).

MAGE-1, MAGE-2, and MAGE-3 are members of another family of cancer-testis antigens originally discovered in melanoma (MAGE is a contraction of melanoma-associated antigen) but found in a variety of tumors. The identification of MAGE proteins as TuAAs is taught in U.S. Patent 5,342,774 entitled NUCLEOTIDE SEQUENCE ENCODING THE TUMOR REJECTION ANTIGEN PRECURSOR, MAGE-1, which is hereby incorporated by reference in its entirety, and in numerous subsequent patents. Currently there are 17 entries for (human) MAGE in the SWISS Protein database. There is extensive similarity among these proteins so in many cases, an epitope from one can induce a cross-reactive response to other members of the family. A few of these have not been observed in tumors, most notably MAGE-H1 and MAGE-D1, which are expressed in testes and brain, and bone marrow stromal cells, respectively. The possibility of cross-reactivity on normal tissue is ameliorated by the fact that they are among the least similar to the other MAGE proteins.

NY-ESO-1, is a cancer-testis antigen found in a wide variety of tumors, also known as CTAG-1 (Cancer-Testis Antigen-1) and CAG-3 (Cancer Antigen-3). NY-ESO-1 as a TuAA is disclosed in U.S. Patent 5,804,381 entitled ISOLATED NUCLEIC ACID MOLECULE

ENCODING AN ESOPHAGEAL CANCER ASSOCIATED ANTIGEN, THE ANTIGEN ITSELF,
AND USES THEREOF which is hereby incorporated by reference in its entirety. A paralogous
locus encoding antigens with extensive sequence identity, LAGE-1a/s (SEQ ID NO. 75) and
LAGE-1b/L (SEQ ID NO. 76), have been disclosed in publicly available assemblies of the human
5 genome, and have been concluded to arise through alternate splicing. Additionally, CT-2 (or
CTAG-2, Cancer-Testis Antigen-2) appears to be either an allele, a mutant, or a sequencing
discrepancy of LAGE-1b/L. Due to the extensive sequence identity, many epitopes from NY-ESO-
1 can also induce immunity to tumors expressing these other antigens. See figure 1. The proteins
are virtually identical through amino acid 70. From 71-134 the longest run of identities between
10 NY-ESO-1 and LAGE is 6 residues, but potentially cross-reactive sequences are present. And
from 135-180 NY-ESO and LAGE-1a/s are identical except for a single residue, but LAGE-1b/L is
unrelated due to the alternate splice. The CAMEL and LAGE-2 antigens appear to derive from the
LAGE-1 mRNA, but from alternate reading frames, thus giving rise to unrelated protein sequences.
More recently, GenBank Accession AF277315.5, Homo sapiens chromosome X clone RP5-
15 865E18, RP5-1087L19, complete sequence, reports three independent loci in this region which are
labeled as LAGE1 (corresponding to CTAG-2 in the genome assemblies), plus LAGE2-A and
LAGE2-B (both corresponding to CTAG-1 in the genome assemblies).

PSMA (prostate-specific membranes antigen), a TuAA described in U.S. Patent 5,538,866
entitled "PROSTATE-SPECIFIC MEMBRANES ANTIGEN" which is hereby incorporated by
20 reference in its entirety, is expressed by normal prostate epithelium and, at a higher level, in
prostatic cancer. It has also been found in the neovasculature of non-prostatic tumors. PSMA can
thus form the basis for vaccines directed to both prostate cancer and to the neovasculature of other
tumors. This later concept is more fully described in a provisional U.S. Patent application No.
60/274,063 entitled ANTI-NEOVASCULAR VACCINES FOR CANCER, filed March 7, 2001,
25 and U.S. Application No. __/_____, attorney docket number CTLIMM.015A, filed on March 7,
2002, entitled "ANTI-NEOVASCULAR PREPARATIONS FOR CANCER," both of which are
hereby incorporated by reference in their entirety. Alternate splicing of the PSMA mRNA also
leads to a protein with an apparent start at Met₅₈, thereby deleting the putative membrane anchor
region of PSMA as described in U.S. Patent 5,935,818 entitled "ISOLATED NUCLEIC ACID
30 MOLECULE ENCODING ALTERNATIVELY SPLICED PROSTATE-SPECIFIC
MEMBRANES ANTIGEN AND USES THEREOF" which is hereby incorporated by reference in
its entirety. A protein termed PSMA-like protein, Genbank accession number AF261715, is nearly
identical to amino acids 309-750 of PSMA and has a different expression profile. Thus the most
preferred epitopes are those with an N-terminus located from amino acid 58 to 308.

35 PRAME, also known as MAPE, DAGE, and OIP4, was originally observed as a melanoma
antigen. Subsequently, it has been recognized as a CT antigen, but unlike many CT antigens (e.g.,

MAGE, GAGE, and BAGE) it is expressed in acute myeloid leukemias. PRAME is a member of the MAPE family which consists largely of hypothetical proteins with which it shares limited sequence similarity. The usefulness of PRAME as a TuAA is taught in U.S. Patent 5,830,753 entitled "ISOLATED NUCLEIC ACID MOLECULES CODING FOR TUMOR REJECTION
5 ANTIGEN PRECURSOR DAGE AND USES THEREOF" which is hereby incorporated by reference in its entirety.

PSA, prostate specific antigen, is a peptidase of the kallikrein family and a differentiation antigen of the prostate. Expression in breast tissue has also been reported. Alternate names include gamma-seminoprotein, kallikrein 3, seminogelase, seminin, and P-30 antigen. PSA has a high
10 degree of sequence identity with the various alternate splicing products prostatic/glandular kallikrein-1 and -2, as well as kallikrein 4, which is also expressed in prostate and breast tissue. Other kallikreins generally share less sequence identity and have different expression profiles. Nonetheless, cross-reactivity that might be provoked by any particular epitope, along with the likelihood that that epitope would be liberated by processing in non-target tissues (most generally
15 by the housekeeping proteasome), should be considered in designing a vaccine.

PSCA, prostate stem cell antigen, and also known as SCAH-2, is a differentiation antigen preferentially expressed in prostate epithelial cells, and overexpressed in prostate cancers. Lower level expression is seen in some normal tissues including neuroendocrine cells of the digestive tract and collecting ducts of the kidney. PSCA is described in U.S. Patent 5,856,136 entitled "HUMAN
20 STEM CELL ANTIGENS" which is hereby incorporated by reference in its entirety.

Synaptonemal complex protein 1 (SCP-1), also known as HOM-TES-14, is a meiosis-associated protein and also a cancer-testis antigen (Tureci, O., et al. *Proc. Natl. Acad. Sci. USA* 95:5211-5216, 1998). As a cancer antigen its expression is not cell-cycle regulated and it is found frequently in gliomas, breast, renal cell, and ovarian carcinomas. It has some similarity to myosins,
25 but with few enough identities that cross-reactive epitopes are not an immediate prospect.

The ED-B domain of fibronectin is also a potential target. Fibronectin is subject to developmentally regulated alternative splicing, with the ED-B domain being encoded by a single exon that is used primarily in oncofetal tissues (Matsuura, H. and S. Hakomori *Proc. Natl. Acad. Sci. USA* 82:6517-6521, 1985; Carnemolla, B. et al. *J. Cell Biol.* 108:1139-1148, 1989; Loridon-Rosa, B. et al. *Cancer Res.* 50:1608-1612, 1990; Nicolo, G. et al. *Cell Differ. Dev.* 32:401-408,
30 1990; Borsi, L. et al. *Exp. Cell Res.* 199:98-105, 1992; Oyama, F. et al. *Cancer Res.* 53:2005-2011, 1993; Mandel, U. et al. *APMIS* 102:695-702, 1994; Farnoud, M.R. et al. *Int. J. Cancer* 61:27-34, 1995; Pujuguet, P. et al. *Am. J. Pathol.* 148:579-592, 1996; Gabler, U. et al. *Heart* 75:358-362, 1996; Chevalier, X. *Br. J. Rheumatol.* 35:407-415, 1996; Midulla, M. *Cancer Res.* 60:164-169, 2000).
35

The ED-B domain is also expressed in fibronectin of the neovasculature (Kaczmarek, J. et al. *Int. J. Cancer* 59:11-16, 1994; Castellani, P. et al. *Int. J. Cancer* 59:612-618, 1994; Neri, D. et al. *Nat. Biotech.* 15:1271-1275, 1997; Karelina, T.V. and A.Z. Eisen *Cancer Detect. Prev.* 22:438-444, 1998; Tarli, L. et al. *Blood* 94:192-198, 1999; Castellani, P. et al. *Acta Neurochir. (Wien)* 142:277-282, 2000). As an oncofetal domain, the ED-B domain is commonly found in the fibronectin expressed by neoplastic cells in addition to being expressed by the neovasculature. Thus, CTL-inducing vaccines targeting the ED-B domain can exhibit two mechanisms of action: direct lysis of tumor cells, and disruption of the tumor's blood supply through destruction of the tumor-associated neovasculature. As CTL activity can decay rapidly after withdrawal of vaccine, interference with normal angiogenesis can be minimal. The design and testing of vaccines targeted to neovasculature is described in Provisional U.S. Patent Application No. 60/274,063 entitled "ANTI-NEOVASCULATURE VACCINES FOR CANCER" and in U.S. Patent Application No. ____/____, attorney docket number CTLIMM.015A, entitled "ANTI-NEOVASCULATURE PREPARATIONS FOR CANCER, filed on date even with this application (March 7, 2002). A tumor cell line is disclosed in Provisional U.S. Application No. ____/____, filed on March 7, 2002, attorney docket number CTLIMM.028PR, entitled "HLA-TRANSGENIC MURINE TUMOR CELL LINE," which is hereby incorporated by reference in its entirety.

Carcinoembryonic antigen (CEA) is a paradigmatic oncofetal protein first described in 1965 (Gold and Freedman, *J. Exp. Med.* 121: 439-462, 1965. Fuller references can be found in the Online Medelian Inheritance in Man; record *114890). It has officially been renamed carcinoembryonic antigen-related cell adhesion molecule 5 (CEACAM5). Its expression is most strongly associated with adenocarcinomas of the epithelial lining of the digestive tract and in fetal colon. CEA is a member of the immunoglobulin supergene family and the defining member of the CEA subfamily.

HER2/NEU is an oncogene related to the epidermal growth factor receptor (van de Vijver, et al., *New Eng. J. Med.* 319:1239-1245, 1988), and apparently identical to the c-ERBB2 oncogene (Di Fiore, et al., *Science* 237: 178-182, 1987). The over-expression of ERBB2 has been implicated in the neoplastic transformation of prostate cancer. As HER2 it is amplified and over-expressed in 25-30% of breast cancers among other tumors where expression level is correlated with the aggressiveness of the tumor (Slamon, et al., *New Eng. J. Med.* 344:783-792, 2001). A more detailed description is available in the Online Medelian Inheritance in Man; record *164870.

All references mentioned herein are hereby incorporated by reference in their entirety. Further, incorporated by reference in its entirety is U.S. Patent Application No. 10/005,905 (attorney docket number CTLIMM.021CP1) entitled "EPITOPE SYNCHRONIZATION IN ANTIGEN PRESENTING CELLS," filed on November 7, 2001 and a continuation thereof, U.S. Application No. ____/____, filed on December 7, 2000, attorney docket number

CTLIMM.21CP1C, also entitled "EPITOPE SYNCHRONIZATION IN ANTIGEN PRESENTING CELLS."

Useful epitopes were identified and tested as described in the following examples. However, these examples are intended for illustration purposes only, and should not be construed as limiting the scope of the invention in any way.

EXAMPLES

Sequences of Specific Preferred Epitopes

Example 1

Manufacture of tyrosinase epitopes.

A. Synthetic production of epitopes

Peptides having an amino acid sequence of any of SEQ ID NO: 1, 8, 9, 11-23, 2-29, 32-44, 47-54, 56-63, 66-68 88-253, or 256-588 are synthesized using either FMOC or tBOC solid phase synthesis methodologies. After synthesis, the peptides are cleaved from their supports with either trifluoroacetic acid or hydrogen fluoride, respectively, in the presence of appropriate protective scavengers. After removing the acid by evaporation, the peptides are extracted with ether to remove the scavengers and the crude, precipitated peptide is then lyophilized. Purity of the crude peptides is determined by HPLC, sequence analysis, amino acid analysis, counterion content analysis and other suitable means. If the crude peptides are pure enough (greater than or equal to about 90% pure), they can be used as is. If purification is required to meet drug substance specifications, the peptides are purified using one or a combination of the following: re-precipitation; reverse-phase, ion exchange, size exclusion or hydrophobic interaction chromatography; or counter-current distribution.

Drug product formulation

GMP-grade peptides are formulated in a parenterally acceptable aqueous, organic, or aqueous-organic buffer or solvent system in which they remain both physically and chemically stable and biologically potent. Generally, buffers or combinations of buffers or combinations of buffers and organic solvents are appropriate. The pH range is typically between 6 and 9. Organic modifiers or other excipients can be added to help solubilize and stabilize the peptides. These include detergents, lipids, co-solvents, antioxidants, chelators and reducing agents. In the case of a lyophilized product, sucrose or mannitol or other lyophilization aids can be added. Peptide solutions are sterilized by membrane filtration into their final container-closure system and either lyophilized for dissolution in the clinic, or stored until use.

B. Construction of expression vectors for use as nucleic acid vaccines

The construction of three generic epitope expression vectors is presented below. The particular advantages of these designs are set forth in U.S. Patent Application No. 09/561,572

entitled "EXPRESSION VECTORS ENCODING EPITOPES OF TARGET-ASSOCIATED ANTIGENS," which has been incorporated by reference in its entirety above.

A suitable *E. coli* strain was then transfected with the plasmid and plated out onto a selective medium. Several colonies were grown up in suspension culture and positive clones were identified by restriction mapping. The positive clone was then grown up and aliquotted into storage vials and stored at -70°C.

A mini-prep (QIAprep Spin Mini-prep: Qiagen, Valencia, CA) of the plasmid was then made from a sample of these cells and automated fluorescent dideoxy sequence analysis was used to confirm that the construct had the desired sequence.

B.1 Construction of pVAX-EP1-IRES-EP2

Overview:

The starting plasmid for this construct is pVAX1 purchased from Invitrogen (Carlsbad, CA). Epitopes EP1 and EP2 were synthesized by GIBCO BRL (Rockville, MD). The IRES was excised from pIRES purchased from Clontech (Palo Alto, CA).

Procedure:

1 pIRES was digested with EcoRI and NotI. The digested fragments were separated by agarose gel electrophoresis, and the IRES fragment was purified from the excised band.

2 pVAX1 was digested with EcoRI and NotI, and the pVAX1 fragment was gel-purified.

3 The purified pVAX1 and IRES fragments were then ligated together.

4 Competent *E. coli* of strain DH5 α were transformed with the ligation mixture.

5 Minipreps were made from 4 of the resultant colonies.

6 Restriction enzyme digestion analysis was performed on the miniprep DNA. One recombinant colony having the IRES insert was used for further insertion of EP1 and EP2. This intermediate construct was called pVAX-IRES.

7 Oligonucleotides encoding EP1 and EP2 were synthesized.

8 EP1 was subcloned into pVAX-IRES between AflII and EcoRI sites, to make pVAX-EP1-IRES;

9 EP2 was subcloned into pVAX-EP1-IRES between SalI and NotI sites, to make the final construct pVAX-EP1-IRES-EP2.

10 The sequence of the EP1-IRES-EP2 insert was confirmed by DNA sequencing.

B.2. Construction of pVAX-EP1-IRES-EP2-ISS-NIS

Overview:

The starting plasmid for this construct was pVAX-EP1-IRES-EP2 (Example 1). The ISS (immunostimulatory sequence) introduced into this construct is AACGTT, and the NIS (standing for nuclear import sequence) used is the SV40 72bp repeat sequence. ISS-NIS was synthesized by GIBCO BRL. See Figure 2.

Procedure:

- 1 pVAX-EP1-IRES-EP2 was digested with NruI; the linearized plasmid was gel-purified.
- 2 ISS-NIS oligonucleotide was synthesized.
- 3 The purified linearized pVAX-EP1-IRES-EP2 and synthesized ISS-NIS were ligated
5 together.
- 4 Competent E. coli of strain DH5 α were transformed with the ligation product.
- 5 Minipreps were made from resultant colonies.
- 6 Restriction enzyme digestions of the minipreps were carried out.
- 7 The plasmid with the insert was sequenced.

10 **B3. Construction of pVAX-EP2-UB-EP1**

Overview:

The starting plasmid for this construct was pVAX1 (Invitrogen). EP2 and EP1 were synthesized by GIBCO BRL. Wild type Ubiquitin cDNA encoding the 76 amino acids in the construct was cloned from yeast.

15 **Procedure:**

- 1 RT-PCR was performed using yeast mRNA. Primers were designed to amplify the complete coding sequence of yeast Ubiquitin.
- 2 The RT-PCR products were analyzed using agarose gel electrophoresis. A band with the predicted size was gel-purified.
- 20 3 The purified DNA band was subcloned into pZERO1 at EcoRV site. The resulting clone was named pZERO-UB.
- 4 Several clones of pZERO-UB were sequenced to confirm the Ubiquitin sequence before further manipulations.
- 5 EP1 and EP2 were synthesized.
- 25 6 EP2, Ubiquitin and EP1 were ligated and the insert cloned into pVAX1 between BamHI and EcoRI, putting it under control of the CMV promoter.
- 7 The sequence of the insert EP2-UB-EP1 was confirmed by DNA sequencing.

Example 2**Identification of useful epitope variants.**

- 30 The 10-mer FLPWHRLFLL (SEQ ID NO. 1) is identified as a useful epitope. Based on this sequence, numerous variants are made. Variants exhibiting activity in HLA binding assays (see Example 3, section 6) are identified as useful, and are subsequently incorporated into vaccines.

- The HLA-A2 binding of length variants of FLPWHRLFLL have been evaluated. Proteasomal digestion analysis indicates that the C-terminus of the 9-mer FLPWHRLFL (SEQ ID
35 NO. 8) is also produced. Additionally the 9-mer LPWHRLFLL (SEQ ID NO. 9) can result from N-terminal trimming of the 10-mer. Both are predicted to bind to the HLA-A*0201 molecule,

however of these two 9-mers, FLPWHRLFL displayed more significant binding and is preferred (see Figs. 3A and B).

Sequence variants of FLPWHRLFL are constructed as follow. Consistent with the binding coefficient table (see Table 3) from the NIH/BIMAS MHC binding prediction program (see reference in example 3 below), binding can be improved by changing the L at position 9, an anchor position, to V. Binding can also be altered, though generally to a lesser extent, by changes at non-anchor positions. Referring generally to Table 3, binding can be increased by employing residues with relatively larger coefficients. Changes in sequence can also alter immunogenicity independently of their effect on binding to MHC. Thus binding and/or immunogenicity can be improved as follows:

By substituting F, L, M, W, or Y for P at position 3; these are all bulkier residues that can also improve immunogenicity independent of the effect on binding. The amine and hydroxyl-bearing residues, Q and N; and S and T; respectively, can also provoke a stronger, cross-reactive response.

By substituting D or E for W at position 4 to improve binding; this addition of a negative charge can also make the epitope more immunogenic, while in some cases reducing cross-reactivity with the natural epitope. Alternatively the conservative substitutions of F or Y can provoke a cross-reactive response.

By substituting F for H at position 5 to improve binding. H can be viewed as partially charged, thus in some cases the loss of charge can hinder cross-reactivity. Substitution of the fully charged residues R or K at this position can enhance immunogenicity without disrupting charge-dependent cross-reactivity.

By substituting I, L, M, V, F, W, or Y for R at position 6. The same caveats and alternatives apply here as at position 5.

By substituting W or F for L at position 7 to improve binding. Substitution of V, I, S, T, Q, or N at this position are not generally predicted to reduce binding affinity by this model (the NIH algorithm), yet can be advantageous as discussed above.

Y and W, which are equally preferred as the Fs at positions 1 and 8, can provoke a useful cross-reactivity. Finally, while substitutions in the direction of bulkiness are generally favored to improve immunogenicity, the substitution of smaller residues such as A, S, and C, at positions 3-7 can be useful according to the theory that contrast in size, rather than bulkiness per se, is an important factor in immunogenicity. The reactivity of the thiol group in C can introduce other properties as discussed in Chen, J.-L., et al. *J. Immunol.* 165:948-955, 2000.

Table 3. 9-mer Coefficient Table for HLA-A*0201*

HLA Coefficient table for file "A_0201_standard"									
Amino Acid Type	1 st	2 nd	3 rd	4 th	5 th	6 th	7 th	8 th	9 th
A	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.00
C	1.000	0.470	1.000	1.000	1.000	1.000	1.000	1.000	1.00
D	0.075	0.100	0.400	4.100	1.000	1.000	0.490	1.000	0.00
E	0.075	1.400	0.064	4.100	1.000	1.000	0.490	1.000	0.00
F	4.600	0.050	3.700	1.000	3.800	1.900	5.800	5.500	0.01
G	1.000	0.470	1.000	1.000	1.000	1.000	0.130	1.000	0.01
H	0.034	0.050	1.000	1.000	1.000	1.000	1.000	1.000	0.01
I	1.700	9.900	1.000	1.000	1.000	2.300	1.000	0.410	2.10
K	3.500	0.100	0.035	1.000	1.000	1.000	1.000	1.000	0.00
L	1.700	72.000	3.700	1.000	1.000	2.300	1.000	1.000	4.30
M	1.700	52.000	3.700	1.000	1.000	2.300	1.000	1.000	1.00
N	1.000	0.470	1.000	1.000	1.000	1.000	1.000	1.000	0.01
P	0.022	0.470	1.000	1.000	1.000	1.000	1.000	1.000	0.00
Q	1.000	7.300	1.000	1.000	1.000	1.000	1.000	1.000	0.00
R	1.000	0.010	0.076	1.000	1.000	1.000	0.200	1.000	0.00
S	1.000	0.470	1.000	1.000	1.000	1.000	1.000	1.000	0.01
T	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.50
V	1.700	6.300	1.000	1.000	1.000	2.300	1.000	0.410	14.00
W	4.600	0.010	8.300	1.000	1.000	1.700	7.500	5.500	0.01
Y	4.600	0.010	3.200	1.000	1.000	1.500	1.000	5.500	0.01

*This table and other comparable data that are publicly available are useful in designing epitope variants and in determining whether a particular variant is substantially similar, or is functionally similar.

Example 3

Cluster Analysis (SSX-2₃₁₋₆₈).

1. Epitope cluster region prediction:

The computer algorithms: SYFPEITHI (internet [http:// access at syfpeithi.bmi-heidelberg.com/Scripts/MHCServer.dll/EpPredict.htm](http://access.at.syfpeithi.bmi-heidelberg.com/Scripts/MHCServer.dll/EpPredict.htm)), based on the book "MHC Ligands and Peptide Motifs" by H.G.Rammensee, J.Bachmann and S.Stevanovic; and HLA Peptide Binding Predictions (NIH) (internet [http:// access at bimas.dcrf.nih.gov/molbio/hla_bin](http://access.at.bimas.dcrf.nih.gov/molbio/hla_bin)), described in Parker, K. C., et al., *J. Immunol.* 152:163, 1994; were used to analyze the protein sequence of SSX-2 (GI:10337583). Epitope clusters (regions with higher than average density of peptide fragments with high predicted MHC affinity) were defined as described fully in U.S. Patent Application No. 09/561,571 entitled "EPITOPE CLUSTERS," filed on April 28, 2000. Using a epitope density ratio cutoff of 2, five and two clusters were defined using the SYFPEITHI and NIH algorithms, respectively, and peptides score cutoffs of 16 (SYFPEITHI) and 5 (NIH). The highest scoring peptide with the NIH algorithm, SSX-2₄₁₋₄₉, with an estimated halftime of dissociation of

>1000 min., does not overlap any other predicted epitope but does cluster with SSX-2₅₇₋₆₅ in the NIH analysis.

2. Peptide synthesis and characterization:

SSX-2₃₁₋₆₈, YFSKEEWEKMKASEKIFYVYMKRKYEAMTKLGFKATLP (SEQ ID NO. 10) was synthesized by MPS (Multiple Peptide Systems, San Diego, CA 92121) using standard solid phase chemistry. According to the provided 'Certificate of Analysis', the purity of this peptide was 95%.

3. Proteasome digestion:

Proteasome was isolated from human red blood cells using the proteasome isolation protocol described in U.S. Patent Application No. 09/561,074 entitled "METHOD OF EPITOPE DISCOVERY," filed on April 28, 2000. SDS-PAGE, western-blotting, and ELISA were used as quality control assays. The final concentration of proteasome was 4 mg/ml, which was determined by non-interfering protein assay (Geno Technologies Inc.). Proteasomes were stored at -70°C in 25 µl aliquots.

SSX-2₃₁₋₆₈ was dissolved in Milli-Q water, and a 2 mM stock solution prepared and 20µL aliquots stored at -20°C.

1 tube of proteasome (25 µL) was removed from storage at -70°C and thawed on ice. It was then mixed thoroughly with 12.5µL of 2mM peptide by repipetting (samples were kept on ice). A 5µL sample was immediately removed after mixing and transferred to a tube containing 1.25µL 10%TFA (final concentration of TFA was 2%); the T=0 min sample. The proteasome digestion reaction was then started and carried out at 37°C in a programmable thermal controller. Additional 5µL samples were taken out at 15, 30, 60, 120, 180 and 240 min respectively, the reaction was stopped by adding the sample to 1.25µL 10% TFA as before. Samples were kept on ice or frozen until being analyzed by MALDI-MS. All samples were saved and stored at -20°C for HPLC analysis and N-terminal sequencing. Peptide alone (without proteasome) was used as a blank control: 2 µL peptide + 4µL Tris buffer (20 mM, pH 7.6) + 1.5µL TFA.

4. MALDI-TOF MS measurements:

For each time point 0.3 µL of matrix solution (10mg/ml α-cyano-4-hydroxycinnamic acid in AcCN/H₂O (70:30)) was first applied on a sample slide, and then an equal volume of digested sample was mixed gently with matrix solution on the slide. The slide was allowed to dry at ambient air for 3-5 min. before acquiring the mass spectra. MS was performed on a Lasermat 2000 MALDI-TOF mass spectrometer that was calibrated with peptide/protein standards. To improve the accuracy of measurement, the molecular ion weight (MH⁺) of the peptide substrate was used as an internal calibration standard. The mass spectrum of the T=120 min. digested sample is shown in figure 4.

5. MS data analysis and epitope identification:

To assign the measured mass peaks, the computer program MS-Product, a tool from the UCSF Mass Spectrometry Facility (<http://prospector.ucsf.edu/ucsfhtml3.4/msprod.htm>), was used to generate all possible fragments (N- and C-terminal ions, and internal fragments) and their corresponding molecular weights. Due to the sensitivity of the mass spectrometer, average molecular weight was used. The mass peaks observed over the course of the digestion were identified as summarized in Table 4.

Fragments co-C-terminal with 8-10 amino acid long sequences predicted to bind HLA by the SYFPEITHI or NIH algorithms were chosen for further study. The digestion and prediction steps of the procedure can be usefully practiced in any order. Although the substrate peptide used in proteasomal digest described here was specifically designed to include predicted HLA-A2.1 binding sequences, the actual products of digestion can be checked after the fact for actual or predicted binding to other MHC molecules. Selected results are shown in Table 5.

Table 4. SSX-2₃₁₋₆₈ Mass Peak Identification.

MS PEAK (measured)	PEPTIDE	SEQUENCE	CALCULATED MASS (MH ⁺)
988.23	31-37	YFSKBEW	989.08
1377.68±2.3 8	31-40	YFSKEEWEKM	1377.68
1662.45±1.3 0	31-43	YFSKEEWEKMKAS	1663.90
2181.72±0.8 5	31-47	YFSKEEWEKMKASEKIF	2181.52
2346.6	31-48	YFSKEEWEKMKASEKIFY	2344.71
1472.16±1.5 4	38-49	EKMKASEKIFYV	1473.77
2445.78±1.1 8	31-49*	YFSKEEWEKMKASEKIFYV	2443.84
2607.	31-50	YFSKEEWEKMKASEKIFYVY	2607.02
1563.3	50-61	YMKRKYEAMTKL	1562.93
3989.9	31-61	YFSKEEWEKMKASEKIFYVYMKRKYEAMTKL	3987.77
1603.74±1.5 3	51-63	MKRKYEAMTKLGF	1603.98
1766.45±1.5	50-63	YMKRKYEAMTKLGF	1767.16
1866.32±1.2 2	49-63	VYMKRKYEAMTKLGF	1866.29
4192.6	31-63	YFSKEEWEKMKASEKIFYVYMKRKYEAMTKLGF	4192.00
4392.1	31-65**	FKA	4391.25

Boldface sequence correspond to peptides predicted to bind to MHC.

* On the basis of mass alone this peak could also have been assigned to the peptide 32-50, however proteasomal removal of just the N-terminal amino acid is unlikely. N-terminal sequencing (below) verifies the assignment to 31-49.

** On the basis of mass this fragment might also represent 33-68. N-terminal sequencing below is consistent with the assignment to 31-65.

Table 5. Predicted HLA binding by proteasomally generated fragments

<u>SEQ ID NO.</u>	<u>PEPTIDE</u>	<u>HLA</u>	<u>SYFPEITHI</u>	<u>NIH</u>
11	FSKEEWEKM	B*3501	NP†	90
12	KMKASEKIF	B*08	17	<5
13 & (14)	(K) MKASEKIFY	A1	19 (19)	<5
15 & (16)	(M) KASEKIFYV	A*0201	22 (16)	1017
		B*08	17	<5
		B*5101	22 (13)	60
		B*5102	NP	133
		B*5103	NP	121
17 & (18)	(K) ASEKIFYVY	A1	34 (19)	14
19 & (20)	(K) RKYEAMTKL	A*0201	15	<5
		A26	15	NP
		B14	NP	45 (60)
		B*2705	21	15
		B*2709	16	NP
		B*5101	15	<5
21	KYEAMTKLGF	A1	16	<5
		A24	NP	300
22	YEAMTKLGF	B*4403	NP	80
23	EAMTKLGF	B*08	22	<5

†No prediction

5

As seen in Table 5, N-terminal addition of authentic sequence to epitopes can generate epitopes for the same or different MHC restriction elements. Note in particular the pairing of (K)RKYEAMTKL (SEQ ID NOS 19 and (20)) with HLA-B14, where the 10-mer has a longer predicted halftime of dissociation than the co-C-terminal 9-mer. Also note the case of the 10-mer KYEAMTKLGF (SEQ ID NO. 21) which can be used as a vaccine useful with several MHC types by relying on N-terminal trimming to create the epitopes for HLA-B*4403 and -B*08.

10

6. HLA-A0201 binding assay:

15

Binding of the candidate epitope KASEKIFYV, SSX-2₄₁₋₄₉, (SEQ ID NO. 15) to HLA-A2.1 was assayed using a modification of the method of Stauss et al., (Proc Natl Acad Sci USA 89(17):7871-5 (1992)). Specifically, T2 cells, which express empty or unstable MHC molecules on their surface, were washed twice with Iscove's modified Dulbecco's medium (IMDM) and cultured overnight in serum-free AIM-V medium (Life Technologies, Inc., Rockville, MD) supplemented with human β 2-microglobulin at 3 μ g/ml (Sigma, St. Louis, MO) and added peptide,

at 800, 400, 200, 100, 50, 25, 12.5, and 6.25 µg/ml in a 96-well flat-bottom plate at 3×10^5 cells/200 µl/well. Peptide was mixed with the cells by repipetting before distributing to the plate (alternatively peptide can be added to individual wells), and the plate was rocked gently for 2 minutes. Incubation was in a 5% CO₂ incubator at 37°C. The next day the unbound peptide was removed by washing twice with serum free RPMI medium and a saturating amount of anti-class I HLA monoclonal antibody, fluorescein isothiocyanate (FITC)-conjugated anti-HLA A2, A28 (One Lambda, Canoga Park, CA) was added. After incubation for 30 minutes at 4°C, cells were washed 3 times with PBS supplemented with 0.5% BSA, 0.05%(w/v) sodium azide, pH 7.4-7.6 (staining buffer). (Alternatively W6/32 (Sigma) can be used as the anti-class I HLA monoclonal antibody the cells washed with staining buffer and then incubated with fluorescein isothiocyanate (FITC)-conjugated goat F(ab') antimouse-IgG (Sigma) for 30 min at 4°C and washed 3 times as before.) The cells were resuspended in 0.5 ml staining buffer. The analysis of surface HLA-A2.1 molecules stabilized by peptide binding was performed by flow cytometry using a FACScan (Becton Dickinson, San Jose, CA). If flow cytometry is not to be performed immediately the cells can be fixed by adding a quarter volume of 2% paraformaldehyde and storing in the dark at 4 °C.

The results of the experiment are shown in Figure 5. SSX-2₄₁₋₄₉ (SEQ ID NO. 15) was found to bind HLA-A2.1 to a similar extent as the known A2.1 binder FLPSDYFPSV (HBV₁₈₋₂₇; SEQ ID NO: 24) used as a positive control. An HLA-B44 binding peptide, AEMGKYSFY (SEQ ID NO: 25), was used as a negative control. The fluorescence obtained from the negative control was similar to the signal obtained when no peptide was used in the assay. Positive and negative control peptides were chosen from Table 18.3.1 in *Current Protocols in Immunology* p. 18.3.2, John Wiley and Sons, New York, 1998.

7. Immunogenicity:

A. In vivo immunization of mice.

HHD1 transgenic A*0201 mice (Pascolo, S., et al. *J. Exp. Med.* 185:2043-2051, 1997) were anesthetized and injected subcutaneously at the base of the tail, avoiding lateral tail veins, using 100 µl containing 100 nmol of SSX-2₄₁₋₄₉ (SEQ ID NO. 15) and 20 µg of HTL epitope peptide in PBS emulsified with 50 µl of IFA (incomplete Freund's adjuvant).

B. Preparation of stimulating cells (LPS blasts).

Using spleens from 2 naive mice for each group of immunized mice, un-immunized mice were sacrificed and the carcasses were placed in alcohol. Using sterile instruments, the top dermal layer of skin on the mouse's left side (lower mid-section) was cut through, exposing the peritoneum. The peritoneum was saturated with alcohol, and the spleen was aseptically extracted. The spleen was placed in a petri dish with serum-free media. Splenocytes were isolated by using sterile plungers from 3 ml syringes to mash the spleens. Cells were collected in a 50 ml conical tubes in serum-free media, rinsing dish well. Cells were centrifuged (12000 rpm, 7 min) and

washed one time with RPMI. Fresh spleen cells were resuspended to a concentration of 1×10^6 cells per ml in RPMI-10%FCS (fetal calf serum). 25g/ml lipopolysaccharide and 7 μ g/ml Dextran Sulfate were added. Cell were incubated for 3 days in T-75 flasks at 37°C, with 5% CO₂. Splenic blasts were collected in 50 ml tubes pelleted (12000 rpm, 7 min) and resuspended to 3×10^7 /ml in RPMI. The blasts were pulsed with the priming peptide at 50 μ g/ml, RT 4hr. mitomycin C-treated at 25 μ g/ml, 37°C, 20 min and washed three times with DMEM.

C. In vitro stimulation.

3 days after LPS stimulation of the blast cells and the same day as peptide loading, the primed mice were sacrificed (at 14 days post immunization) to remove spleens as above. 3×10^6 splenocytes were co-cultured with 1×10^6 LPS blasts/well in 24-well plates at 37°C, with 5% CO₂ in DMEM media supplemented with 10% FCS, 5×10^{-5} M β -mercaptoethanol, 100 μ g/ml streptomycin and 100 IU/ml penicillin. Cultures were fed 5% (vol/vol) ConA supernatant on day 3 and assayed for cytolytic activity on day 7 in a ⁵¹Cr-release assay.

D. Chromium-release assay measuring CTL activity.

To assess peptide specific lysis, 2×10^6 T2 cells were incubated with 100 μ Ci sodium chromate together with 50 μ g/ml peptide at 37°C for 1 hour. During incubation they were gently shaken every 15 minutes. After labeling and loading, cells were washed three times with 10 ml of DMEM-10% FCS, wiping each tube with a fresh Kimwipe after pouring off the supernatant. Target cells were resuspended in DMEM-10% FBS 1×10^5 /ml. Effector cells were adjusted to 1×10^7 /ml in DMEM-10% FCS and 100 μ l serial 3-fold dilutions of effectors were prepared in U-bottom 96-well plates. 100 μ l of target cells were added per well. In order to determine spontaneous release and maximum release, six additional wells containing 100 μ l of target cells were prepared for each target. Spontaneous release was revealed by incubating the target cells with 100 μ l medium; maximum release was revealed by incubating the target cells with 100 μ l of 2% SDS. Plates were then centrifuged for 5 min at 600 rpm and incubated for 4 hours at 37°C in 5% CO₂ and 80% humidity. After the incubation, plates were then centrifuged for 5 min at 1200 rpm. Supernatants were harvested and counted using a gamma counter. Specific lysis was determined as follows: % specific release = [(experimental release - spontaneous release)/(maximum release - spontaneous release)] x 100.

Results of the chromium release assay demonstrating specific lysis of peptide pulsed target cells are shown in figure 6.

8. Cross-reactivity with other SSX proteins:

SSX-2₄₁₋₄₉ (SEQ ID NO. 15) shares a high degree of sequence identity with the same region of the other SSX proteins. The surrounding regions have also been generally well conserved. Thus the housekeeping proteasome can cleave following V₄₉ in all five sequences. Moreover, SSX₄₁₋₄₉ is

predicted to bind HLA-A*0201 (see Table 6). CTL generated by immunization with SSX-2₄₁₋₄₉ cross-react with tumor cells expressing other SSX proteins.

Table 6. SSX₄₁₋₄₉ - A*0201 Predicted Binding

SEQ ID NO.	Family Member	Sequence	SYFPEITHI Score	NIH Score
15	SSX-2	KASEKIFYV	22	1017
26	SSX-1	KYSEKISYV	18	1.7
27	SSX-3	KVSEKIVYV	24	1105
28	SSX-4	KSSEKIVYV	20	82
29	SSX-5	KASEKIYV	22	175

5 Example 4

Cluster Analysis (PSMA₁₆₃₋₁₉₂).

A peptide, AFSPQGMPEGDLVYVNYARTEDFFKLERDM, PSMA₁₆₃₋₁₉₂, (SEQ ID NO. 30), containing an A1 epitope cluster from prostate specific membrane antigen, PSMA₁₆₈₋₁₉₀ (SEQ ID NO. 31) was synthesized using standard solid-phase F-moc chemistry on a 433A ABI Peptide synthesizer. After side chain deprotection and cleavage from the resin, peptide first dissolved in formic acid and then diluted into 30% Acetic acid, was run on a reverse-phase preparative HPLC C4 column at following conditions: linear AB gradient (5% B/min) at a flow rate of 4 ml/min, where eluent A is 0.1% aqueous TFA and eluent B is 0.1% TFA in acetonitrile. A fraction at time 16.642 min containing the expected peptide, as judged by mass spectrometry, was pooled and lyophilized. The peptide was then subjected to proteasome digestion and mass spectrum analysis essentially as described above. Prominent peaks from the mass spectra are summarized in Table 7.

Table 7. PSMA₁₆₃₋₁₉₂ Mass Peak Identification.

PEPTIDE	SEQUENCE	CALCULATE D MASS (MH ⁺)
163-177	AFSPQGMPEGDLVYV	1610.0
178-189	NYARTEDFFKLE	1533.68
170-189	PEGDLVYVNYARTEDFFKLE	2406.66
178-191	NYARTEDFFKLERD	1804.95
170-191	PEGDLVYVNYARTEDFFKLERD	2677.93
178-192	NYARTEDFFKLERDM	1936.17
163-176	AFSPQGMPEGDLVY	1511.70
177-192	VNYARTEDFFKLERDM	2035.30
163-179	AFSPQGMPEGDLVYVNY	1888.12

180-192	ARTEDFFKLERDM	1658.89
163-183	AFSPQGMPEGDLVYVNYARTE	2345.61
184-192	DFFKLERDM	1201.40
176-192	YVNYARTEDFFKLERDM	2198.48
167-185	QGMPEGDLVYVNYARTEDF	2205.41
178-186	NYARTEDFF	1163.22

Boldface sequences correspond to peptides predicted to bind to MHC, see Table 8.

N-terminal Pool Sequence Analysis

One aliquot at one hour of the proteasomal digestion (see Example 3 part 3 above) was subjected to N-terminal amino acid sequence analysis by an ABI 473A Protein Sequencer (Applied Biosystems, Foster City, CA). Determination of the sites and efficiencies of cleavage was based on consideration of the sequence cycle, the repetitive yield of the protein sequencer, and the relative yields of amino acids unique in the analyzed sequence. That is if the unique (in the analyzed sequence) residue X appears only in the nth cycle a cleavage site exists n-1 residues before it in the N-terminal direction. In addition to helping resolve any ambiguity in the assignment of mass to sequences, these data also provide a more reliable indication of the relative yield of the various fragments than does mass spectrometry.

For PSMA₁₆₃₋₁₉₂ (SEQ ID NO. 30) this pool sequencing supports a single major cleavage site after V₁₇₇ and several minor cleavage sites, particularly one after Y₁₇₉. Reviewing the results presented in figures 7A-C reveals the following:

- S at the 3rd cycle indicating presence of the N-terminus of the substrate.
 - Q at the 5th cycle indicating presence of the N-terminus of the substrate.
 - N at the 1st cycle indicating cleavage after V₁₇₇.
 - N at the 3rd cycle indicating cleavage after V₁₇₅. Note the fragment 176-192 in Table 7.
 - T at the 5th cycle indicating cleavage after V₁₇₇.
 - T at the 1st - 3rd cycles, indicating increasingly common cleavages after R₁₈₁, A₁₈₀ and Y₁₇₉.
- Only the last of these correspond to peaks detected by mass spectrometry; 163-179 and 180-192, see Table 7. The absence of the others can indicate that they are on fragments smaller than were examined in the mass spectrum.
- K at the 4th, 8th, and 10th cycles indicating cleavages after E₁₈₃, Y₁₇₉, and V₁₇₇, respectively, all of which correspond to fragments observed by mass spectroscopy. See Table 7.
 - A at the 1st and 3rd cycles indicating presence of the N-terminus of the substrate and cleavage after V₁₇₇, respectively.
 - P at the 4th and 8th cycles indicating presence of the N-terminus of the substrate.

G at the 6th and 10th cycles indicating presence of the N-terminus of the substrate.

M at the 7th cycle indicating presence of the N-terminus of the substrate and/or cleavage after F₁₈₅.

M at the 15th cycle indicating cleavage after V₁₇₇.

5 The 1st cycle can indicate cleavage after D₁₉₁, see Table 7.

R at the 4th and 13th cycle indicating cleavage after V₁₇₇.

R at the 2nd and 11th cycle indicating cleavage after Y₁₇₉.

V at the 2nd, 6th, and 13th cycle indicating cleavage after V₁₇₅, M₁₆₉ and presence of the N-terminus of the substrate, respectively. Note fragments beginning at 176 and 170 in

10 Table 7.

Y at the 1st, 2nd, and 14th cycles indicating cleavage after V₁₇₅, V₁₇₇, and presence of the N-terminus of the substrate, respectively.

L at the 11th and 12th cycles indicating cleavage after V₁₇₇, and presence of the N-terminus of the substrate, respectively, is the interpretation most consistent with the other data.

15 Comparing to the mass spectrometry results we see that L at the 2nd, 5th, and 9th cycles is consistent with cleavage after F₁₈₆, E₁₈₃ or M₁₆₉, and Y₁₇₉, respectively. See Table 7.

Epitope Identification

20 Fragments co-C-terminal with 8-10 amino acid long sequences predicted to bind HLA by the SYFPEITHI or NIH algorithms were chosen for further analysis. The digestion and prediction steps of the procedure can be usefully practiced in any order. Although the substrate peptide used in proteasomal digest described here was specifically designed to include a predicted HLA-A1 binding sequence, the actual products of digestion can be checked after the fact for actual or predicted binding to other MHC molecules. Selected results are shown in Table 8.

Table 8. Predicted HLA binding by proteasomally generated fragments

SEQ ID NO	PEPTIDE	HLA	SYFPEITHI	NIH
32 & (33)	(G)MPEGDLVY V	A*0201	17 (27)	(2605)
		B*0702	20	<5
		B*5101	22	314
34 & (35)	(Q)GMPEGDLV Y	A1	24 (26)	<5
		A3	16 (18)	36
		B*2705	17	25
	36 MPEGDLVY	B*5101	15	NP†
37 & (38)	(P)EGDLVYVN Y	A1	27 (15)	12
		A26	23 (17)	NP
39	LVYVNYARTE	A3	21	<5
40 & (41)	(Y)VNYARTED F	A26	(20)	NP
		B*08	15	<5
		B*2705	12	50
42	NYARTEDFF	A24	NP†	100
		Cw*0401	NP	120
43	YARTEDFF	B*08	16	<5
44	RTEDFFKLE	A1	21	<5
		A26	15	NP

†No prediction

5 HLA-A*0201 binding assay:

HLA-A*0201 binding studies were performed with PSMA₁₆₈₋₁₇₇, GMPEGDLVYV, (SEQ ID NO. 33) essentially as described in Example 3 above. As seen in figure 8, this epitope exhibits significant binding at even lower concentrations than the positive control peptides. The Melan-A peptide used as a control in this assay (and throughout this disclosure), ELAGIGILTV, is actually a variant of the natural sequence (EAAGIGILTV) and exhibits a high affinity in this assay.

10

Example 5**Cluster Analysis (PSMA₂₈₁₋₃₁₀).**

Another peptide, RGIAEAVGLPSIPVHPIGYYDAQKLEKMG, PSMA₂₈₁₋₃₁₀, (SEQ ID NO. 45), containing an A1 epitope cluster from prostate specific membrane antigen, PSMA₂₈₃₋₃₀₇ (SEQ ID NO. 46), was synthesized using standard solid-phase F-moc chemistry on a 433A ABI Peptide synthesizer. After side chain deprotection and cleavage from the resin, peptide in ddH₂O was run on a reverse-phase preparative HPLC C18 column at following conditions: linear AB gradient (5% B/min) at a flow rate of 4 ml/min, where eluent A is 0.1% aqueous TFA and eluent B is 0.1% TFA in acetonitrile. A fraction at time 17.061 min containing the expected peptide as judged by mass spectrometry, was pooled and lyophilized. The peptide was then subjected to proteasome digestion and mass spectrum analysis essentially as described above. Prominent peaks from the mass spectra are summarized in Table 9.

Table 9. PSMA₂₈₁₋₃₁₀ Mass Peak Identification.

PEPTIDE	SEQUENCE	CALCULATE D MASS (MH ⁺)
281-297	RGIAEAVGLPSIPVHPI*	1727.07
286-297	AVGLPSIPVHPI**	1200.46
287-297	VGLPSIPVHPI	1129.38
288-297	GLPSIPVHPI [†]	1030.25
298-310	GYDAQKLEKMG [‡]	1516.5
298-305	GYDAQKLS	958.05
281-305	RGIAEAVGLPSIPVHPIGYYDAQKL	2666.12
281-307	RGIAEAVGLPSIPVHPIGYYDAQKLE	2908.39
286-307	AVGLPSIPVHPIGYYDAQKLE¶	2381.78
287-307	VGLPSIPVHPIGYYDAQKLE	2310.70
288-307	GLPSIPVHPIGYYDAQKLE#	2211.57
281-299	RGIAEAVGLPSIPVHPIGY	1947
286-299	AVGLPSIPVHPIGY	1420.69
287-299	VGLPSIPVHPIGY	1349.61
288-299	GLPSIPVHPIGY	1250.48
287-310	VGLPSIPVHPIGYYDAQKLEKMG	2627.14
288-310	GLPSIPVHPIGYYDAQKLEKMG	2528.01

Boldface sequences correspond to peptides predicted to bind to MHC, see Table 10.

*By mass alone this peak could also have been 296-310 or 288-303.

**By mass alone this peak could also have been 298-307. Combination of HPLC and mass spectrometry show that at some later time points this peak is a mixture of both species.

† By mass alone this peak could also have been 289-298.

- By mass alone this peak could also have been 281-295 or 294-306.
 § By mass alone this peak could also have been 297-303.
 ¶ By mass alone this peak could also have been 285-306.
 # By mass alone this peak could also have been 288-303.
 5 None of these alternate assignments are supported N-terminal pool sequence analysis.

N-terminal Pool Sequence Analysis

One aliquot at one hour of the proteasomal digestion (see Example 3 part 3 above) was subjected to N-terminal amino acid sequence analysis by an ABI 473A Protein Sequencer (Applied Biosystems, Foster City, CA). Determination of the sites and efficiencies of cleavage was based on
 10 consideration of the sequence cycle, the repetitive yield of the protein sequencer, and the relative yields of amino acids unique in the analyzed sequence. That is if the unique (in the analyzed sequence) residue X appears only in the nth cycle a cleavage site exists n-1 residues before it in the N-terminal direction. In addition to helping resolve any ambiguity in the assignment of mass to
 15 sequences, these data also provide a more reliable indication of the relative yield of the various fragments than does mass spectrometry.

For PSMA₂₈₁₋₃₁₀ (SEQ ID NO. 45) this pool sequencing supports two major cleavage sites after V₂₈₇ and I₂₉₇ among other minor cleavage sites. Reviewing the results presented in Fig. 9 reveals the following:

- 20 S at the 4th and 11th cycles indicating cleavage after V₂₈₇ and presence of the N-terminus of the substrate, respectively.
 H at the 8th cycle indicating cleavage after V₂₈₇. The lack of decay in peak height at positions 9 and 10 versus the drop in height present going from 10 to 11 can suggest cleavage after A₂₈₆ and E₂₈₅ as well, rather than the peaks representing latency in the
 25 sequencing reaction.
 D at the 2nd, 4th, and 7th cycles indicating cleavages after Y₂₉₉, I₂₉₇, and V₂₉₄, respectively. This last cleavage is not observed in any of the fragments in Table 10 or in the alternate assignments in the notes below.
 Q at the 6th cycle indicating cleavage after I₂₉₇.
 30 M at the 10th and 12th cycle indicating cleavages after Y₂₉₉ and I₂₉₇, respectively.

Epitope Identification

Fragments co-C-terminal with 8-10 amino acid long sequences predicted to bind HLA by the SYFPEITHI or NIH algorithms were chosen for further study. The digestion and prediction steps of the procedure can be usefully practiced in any order. Although the substrate peptide used
 35 in proteasomal digest described here was specifically designed to include a predicted HLA-A1 binding sequence, the actual products of digestion can be checked after the fact for actual or predicted binding to other MHC molecules. Selected results are shown in Table 10.

Table 10.**Predicted HLA binding by proteasomally generated fragments: PSMA₂₈₁₋₃₁₀**

SEQ ID NO.	PEPTIDE	HLA	SYFPEITHI	NIH
47 & (48)	(G)LPSIPVHPI	A*0201	16 (24)	(24)
		B*0702/B7	23	12
		B*5101	24	572
		Cw*0401	NP†	20
49 & (50)	(P)IGYYDAQKL	A*0201	(16)	<5
		A26	(20)	NP
		B*2705	16	25
		B*2709	15	NP
		B*5101	21	57
		Cw*0301	NP	24
51 & (52)	(P)SIPVHPIGY	A1	21 (27)	<5
		A26	22	NP
		A3	16	<5
		B*5101	16	NP
53	IPVHPIGY			
54	YYDAQKLE	A1	22	<5

†No prediction

5

As seen in Table 10, N-terminal addition of authentic sequence to epitopes can often generate still useful, even better epitopes, for the same or different MHC restriction elements. Note for example the pairing of (G)LPSIPVHPI with HLA-A*0201, where the 10-mer can be used as a vaccine useful with several MHC types by relying on N-terminal trimming to create the epitopes for HLA-B7, -B*5101, and Cw*0401.

10

HLA-A*0201 binding assay:

HLA-A*0201 binding studies were preformed with PSMA₂₈₈₋₂₉₇, GLPSIPVHPI, (SEQ ID NO. 48) essentially as described in Examples 3 and 4 above. As seen in figure 8, this epitope exhibits significant binding at even lower concentrations than the positive control peptides.

Example 6**Cluster Analysis (PSMA₄₅₄₋₄₈₁).**

Another peptide, SSIEGNYTLRVDCPTPLMYSLVHLTKEL, PSMA₄₅₄₋₄₈₁, (SEQ ID NO. 55) containing an epitope cluster from prostate specific membrane antigen, was synthesized by MPS (purity >95%) and subjected to proteasome digestion and mass spectrum analysis as described above. Prominent peaks from the mass spectra are summarized in Table 11.

Table 11. PSMA₄₅₄₋₄₈₁ Mass Peak Identification.

MS PEAK (measured)	PEPTIDE	SEQUENCE	CALCULATED MASS (MH ⁺)
1238.5	454-464	SSIEGNYTLRV	1239.78
1768.38±0.60	454-469	SSIEGNYTLRVDCPTPL	1768.99
1899.8	454-470	SSIEGNYTLRVDCPTPLM	1900.19
1097.63±0.91	463-471	RVDCTPLMY	1098.32
2062.87±0.68	454-471*	SSIEGNYTLRVDCPTPLMY	2063.36
1153	472-481**	SLVHNLTKEL	1154.36
1449.93±1.79	470-481	MYSLVHNLTKEL	1448.73

10 **Boldface** sequence correspond to peptides predicted to bind to MHC, see Table 12.

* On the basis of mass alone this peak could equally well be assigned to the peptide 455-472 however proteasomal removal of just the N-terminal amino acid is considered unlikely. If the issue were important it could be resolved by N-terminal sequencing.

15 **On the basis of mass this fragment might also represent 455-464.

Epitope Identification

Fragments co-C-terminal with 8-10 amino acid long sequences predicted to bind HLA by the SYFPEITHI or NIH algorithms were chosen for further study. The digestion and prediction steps of the procedure can be usefully practiced in any order. Although the substrate peptide used in proteasomal digest described here was specifically designed to include predicted HLA-A2.1 binding sequences, the actual products of digestion can be checked after the fact for actual or predicted binding to other MHC molecules. Selected results are shown in Table 12.

Table 12. Predicted HLA binding by proteasomally generated fragments

<u>SEQ ID NO</u>	<u>PEPTIDE</u>	<u>HLA</u>	<u>SYFPEITHI</u>	<u>NIH</u>
56 & (57)	(S) IEGNYTLRV	A1	(19)	<5
58	EGNYTLRV	A*0201	16 (22)	<5
		B*5101	15	NP†
59 & (60)	(Y) TLRVDCTPL	A*0201	20 (18)	(5)
		A26	16 (18)	NP
		B7	14	40
		B8	23	<5
		B*2705	12	30
		Cw*0301	NP	(30)
61	LRVDCTPLM	B*2705	20	600
		B*2709	20	NP
62 & (63)	(L) RVDCTPLMY	A1	32 (22)	125 (13.5)
		A3	25	<5
		A26	22	NP
		B*2702	NP	(200)
		B*2705	13 (NP)	(1000)

†No prediction

- 5 As seen in Table 12, N-terminal addition of authentic sequence to epitopes can often generate still useful, even better epitopes, for the same or different MHC restriction elements. Note for example the pairing of (L)RVDCTPLMY (SEQ ID NOS 62 and (63)) with HLA-B*2702/5, where the 10-mer has substantial predicted halftimes of dissociation and the co-C-terminal 9-mer does not. Also note the case of SIEGNYTLRV (SEQ ID NO 57) a predicted HLA-A*0201 epitope which can be used as a vaccine useful with HLA-B*5101 by relying on N-terminal trimming to create the epitope.

HLA-A*0201 binding assay

- 15 HLA-A*0201 binding studies were preformed, essentially as described in Example 3 above, with PSMA₄₆₀₋₄₆₉, TLRVDCTPL, (SEQ ID NO. 60). As seen in figure 10, this epitope was found to bind HLA-A2.1 to a similar extent as the known A2.1 binder FLPSDYFPSV (HBV₁₈₋₂₇; SEQ ID NO: 24) used as a positive control. Additionally, PSMA₄₆₁₋₄₆₉, (SEQ ID NO. 59) binds nearly as well.

ELISPOT analysis: PSMA₄₆₃₋₄₇₁ (SEQ ID NO. 62)

- 20 The wells of a nitrocellulose-backed microtiter plate were coated with capture antibody by incubating overnight at 4°C using 50 µl/well of 4µg/ml murine anti-human γ-IFN monoclonal

antibody in coating buffer (35 mM sodium bicarbonate, 15 mM sodium carbonate, pH 9.5). Unbound antibody was removed by washing 4 times 5 min. with PBS. Unbound sites on the membrane then were blocked by adding 200µl/well of RPMI medium with 10% serum and incubating 1 hr. at room temperature. Antigen stimulated CD8⁺ T cells, in 1:3 serial dilutions, were seeded into the wells of the microtiter plate using 100µl/well, starting at 2x10⁵ cells/well. (Prior antigen stimulation was essentially as described in Scheibenbogen, C. et al. *Int. J. Cancer* 71:932-936, 1997. PSMA₄₆₂₋₄₇₁ (SEQ ID NO. 62) was added to a final concentration of 10µg/ml and IL-2 to 100 U/ml and the cells cultured at 37°C in a 5% CO₂, water-saturated atmosphere for 40 hrs. Following this incubation the plates were washed with 6 times 200 µl/well of PBS containing 0.05% Tween-20 (PBS-Tween). Detection antibody, 50µl/well of 2g/ml biotinylated murine anti-human γ-IFN monoclonal antibody in PBS+10% fetal calf serum, was added and the plate incubated at room temperature for 2 hrs. Unbound detection antibody was removed by washing with 4 times 200 µl of PBS-Tween. 100µl of avidin-conjugated horseradish peroxidase (Pharmingen, San Diego, CA) was added to each well and incubated at room temperature for 1 hr. Unbound enzyme was removed by washing with 6 times 200 µl of PBS-Tween. Substrate was prepared by dissolving a 20 mg tablet of 3-amino 9-ethylcarbasole in 2.5 ml of N, N-dimethylformamide and adding that solution to 47,5 ml of 0.05 M phosphate-citrate buffer (pH 5.0). 25 µl of 30% H₂O₂ was added to the substrate solution immediately before distributing substrate at 100 µl/well and incubating the plate at room temperature. After color development (generally 15-30 min.), the reaction was stopped by washing the plate with water. The plate was air dried and the spots counted using a stereomicroscope.

Figure 11 shows the detection of PSMA₄₆₃₋₄₇₁ (SEQ ID NO. 62)-reactive HLA-A1⁺ CD8⁺ T cells previously generated in cultures of HLA-A1⁺ CD8⁺ T cells with autologous dendritic cells plus the peptide. No reactivity is detected from cultures without peptide (data not shown). In this case it can be seen that the peptide reactive T cells are present in the culture at a frequency between 1 in 2.2x10⁴ and 1 in 6.7x10⁴. That this is truly an HLA-A1-restricted response is demonstrated by the ability of anti-HLA-A1 monoclonal antibody to block γ-IFN production; see figure 12.

Example 7

Cluster Analysis (PSMA₆₅₃₋₆₈₇).

Another peptide, FDKSNPIVLRMMNDQLMFLERAFIDPLGLPDRPFY PSMA₆₅₃₋₆₈₇, (SEQ ID NO. 64) containing an A2 epitope cluster from prostate specific membrane antigen, PSMA₆₆₀₋₆₈₁ (SEQ ID NO 65), was synthesized by MPS (purity >95%) and subjected to proteasome digestion and mass spectrum analysis as described above. Prominent peaks from the mass spectra are summarized in Table 13.

Table 13. PSMA₆₅₃₋₆₈₇ Mass Peak Identification.

MS PEAK (measured)	PEPTIDE	SEQUENCE	CALCULATED MASS (MH ⁺)
906.17±0.65	681-687**	LPDRPFY	908.05
1287.73±0.76	677-687**	DPLGLPDRPFY	1290.47
1400.3±1.79	676-687	IDPLGLPDRPFY	1403.63
1548.0±1.37	675-687	FIDPLGLPDRPFY	1550.80
1619.5±1.51	674-687**	AFIDPLGLPDRPFY	1621.88
1775.48±1.32	673-687*	RAFIDPLGLPDRPFY	1778.07
2440.2±1.3	653-672	FDKSNPIVLRMMNDQLMFLE	2442.932
1904.63±1.56	672-687*	ERAFIDPLGLPDRPFY	1907.19
2310.6±2.5	653-671	FDKSNPIVLRMMNDQLMFL	2313.82
2017.4±1.94	671-687	LERAFIDPLGLPDRPFY	2020.35
2197.43±1.78	653-670	FDKSNPIVLRMMNDQLMF	2200.66

Boldface sequence correspond to peptides predicted to bind to MHC, see Table 13.

* On the basis of mass alone this peak could equally well be assigned to a peptide beginning at 654, however proteasomal removal of just the N-terminal amino acid is considered unlikely. If the issue were important it could be resolved by N-terminal sequencing.

** On the basis of mass alone these peaks could have been assigned to internal fragments, but given the overall pattern of digestion it was considered unlikely.

Epitope Identification

Fragments co-C-terminal with 8-10 amino acid long sequences predicted to bind HLA by the SYFPEITHI or NIH algorithms were chosen for further study. The digestion and prediction steps of the procedure can be usefully practiced in any order. Although the substrate peptide used in proteasomal digest described here was specifically designed to include predicted HLA-A2.1 binding sequences, the actual products of digestion can be checked after the fact for actual or predicted binding to other MHC molecules. Selected results are shown in Table 14.

Table 14. Predicted HLA binding by proteasomally generated fragments

SEQ ID NO	PEPTIDE	HLA	SYFPEITHI	NIH
66 & (67)	(R)MMNDQLMF L	A*0201	24 (23)	1360 (722)
		A*0205	NP†	71 (42)
		A26	15	NP
		B*2705	12	50
68	RMMNDQLMF	B*2705	17	75

†No prediction

5

As seen in Table 14, N-terminal addition of authentic sequence to epitopes can generate still useful, even better epitopes, for the same or different MHC restriction elements. Note for example the pairing of (R)MMNDQLMFL (SEQ ID NOS. 66 and (67)) with HLA-A*02, where the 10-mer retains substantial predicted binding potential.

10

HLA-A*0201 binding assay

HLA-A*0201 binding studies were performed, essentially as described in Example 3 above, with PSMA₆₆₃₋₆₇₁, (SEQ ID NO. 66) and PSMA₆₆₂₋₆₇₁, RMMNDQLMFL (SEQ NO. 67). As seen in figures 10, 13 and 14, this epitope exhibits significant binding at even lower concentrations than the positive control peptide (FLPSDYFPSV (HBV₁₈₋₂₇); SEQ ID NO: 24). Though not run in parallel, comparison to the controls suggests that PSMA₆₆₂₋₆₇₁ (which approaches the Melan A peptide in affinity) has the superior binding activity of these two PSMA peptides.

15

Example 8**Vaccinating with epitope vaccines.**

20

1. Vaccination with peptide vaccines:A. Intranodal delivery

A formulation containing peptide in aqueous buffer with an antimicrobial agent, an antioxidant, and an immunomodulating cytokine, was injected continuously over several days into the inguinal lymph node using a miniature pumping system developed for insulin delivery (MiniMed; Northridge, CA). This infusion cycle was selected in order to mimic the kinetics of antigen presentation during a natural infection.

25

B. Controlled release

A peptide formulation is delivered using controlled PLGA microspheres as is known in the art, which alter the pharmacokinetics of the peptide and improve immunogenicity. This formulation is injected or taken orally.

30

C. Gene gun delivery

A peptide formulation is prepared wherein the peptide is adhered to gold microparticles as is known in the art. The particles are delivered in a gene gun, being accelerated at high speed so as to penetrate the skin, carrying the particles into dermal tissues that contain pAPCs.

D. Aerosol delivery

A peptide formulation is inhaled as an aerosol as is known in the art, for uptake into appropriate vascular or lymphatic tissue in the lungs.

2. Vaccination with nucleic acid vaccines:

A nucleic acid vaccine is injected into a lymph node using a miniature pumping system, such as the MiniMed insulin pump. A nucleic acid construct formulated in an aqueous buffered solution containing an antimicrobial agent, an antioxidant, and an immunomodulating cytokine, is delivered over a several day infusion cycle in order to mimic the kinetics of antigen presentation during a natural infection.

Optionally, the nucleic acid construct is delivered using controlled release substances, such as PLGA microspheres or other biodegradable substances. These substances are injected or taken orally. Nucleic acid vaccines are given using oral delivery, priming the immune response through uptake into GALT tissues. Alternatively, the nucleic acid vaccines are delivered using a gene gun, wherein the nucleic acid vaccine is adhered to minute gold particles. Nucleic acid constructs can also be inhaled as an aerosol, for uptake into appropriate vascular or lymphatic tissue in the lungs.

Example 9

Assays for the effectiveness of epitope vaccines.

1. Tetramer analysis:

Class I tetramer analysis is used to determine T cell frequency in an animal before and after administration of a housekeeping epitope. Clonal expansion of T cells in response to an epitope indicates that the epitope is presented to T cells by pAPCs. The specific T cell frequency is measured against the housekeeping epitope before and after administration of the epitope to an animal, to determine if the epitope is present on pAPCs. An increase in frequency of T cells specific to the epitope after administration indicates that the epitope was presented on pAPC.

2. Proliferation assay:

Approximately 24 hours after vaccination of an animal with housekeeping epitope, pAPCs are harvested from PBMCs, splenocytes, or lymph node cells, using monoclonal antibodies against specific markers present on pAPCs, fixed to magnetic beads for affinity purification. Crude blood or splenocyte preparation is enriched for pAPCs using this technique. The enriched pAPCs are then used in a proliferation assay against a T cell clone that has been generated and is specific for the housekeeping epitope of interest. The pAPCs are coincubated with the T cell clone and the T cells are monitored for proliferation activity by measuring the incorporation of radiolabeled

thymidine by T cells. Proliferation indicates that T cells specific for the housekeeping epitope are being stimulated by that epitope on the pAPCs.

3. Chromium release assay:

A human patient, or non-human animal genetically engineered to express human class I MHC, is immunized using a housekeeping epitope. T cells from the immunized subject are used in a standard chromium release assay using human tumor targets or targets engineered to express the same class I MHC. T cell killing of the targets indicates that stimulation of T cells in a patient would be effective at killing a tumor expressing a similar TuAA.

Example 10

Induction of CTL response with naked DNA is efficient by Intra-lymph node immunization.

In order to quantitatively compare the CD8⁺ CTL responses induced by different routes of immunization a plasmid DNA vaccine (pEGFP-L33A) containing a well-characterized immunodominant CTL epitope from the LCMV-glycoprotein (G) (gp33; amino acids 33-41) (Oehen, S., et al., *Immunology* 99, 163-169 2000) was used, as this system allows a comprehensive assessment of antiviral CTL responses. Groups of 2 C57BL/6 mice were immunized once with titrated doses (200-0.02µg) of pEGFP-L33A DNA or of control plasmid pEGFP-N3, administered i.m. (intramuscular), i.d. (intradermal), i.spl. (intrasplenic), or i.ln. (intra-lymph node). Positive control mice received 500 pfu LCMV i.v. (intravenous). Ten days after immunization spleen cells were isolated and gp33-specific CTL activity was determined after secondary *in vitro* restimulation. As shown in Fig. 15, i.m. or i.d. immunization induced weakly detectable CTL responses when high doses of pEGFP-L33A DNA (200µg) were administered. In contrast, potent gp33-specific CTL responses were elicited by immunization with only 2µg pEGFP-L33A DNA i.spl. and with as little as 0.2µg pEGFP-L33A DNA given i.ln. (figure 15; symbols represent individual mice and one of three similar experiments is shown). Immunization with the control pEGFP-N3 DNA did not elicit any detectable gp33-specific CTL responses (data not shown).

Example 11

Intra-lymph node DNA immunization elicits anti-tumor immunity.

To examine whether the potent CTL responses elicited following i.ln. immunization were able to confer protection against peripheral tumors, groups of 6 C57BL/6 mice were immunized three times at 6-day intervals with 10µg of pEGFP-L33A DNA or control pEGFP-N3 DNA. Five days after the last immunization small pieces of solid tumors expressing the gp33 epitope (EL4-33) were transplanted s.c. into both flanks and tumor growth was measured every 3-4d. Although the EL4-33 tumors grew well in mice that had been repetitively immunized with control pEGFP-N3 DNA (figure 16), mice which were immunized with pEGFP-L33A DNA i.ln. rapidly eradicated the peripheral EL4-33 tumors (figure 16).

Example 12

Differences in lymph node DNA content mirrors differences in CTL response following intra-lymph node and intramuscular injection.

pEFGPL33A DNA was injected i.ln. or i.m. and plasmid content of the injected or draining lymph node was assessed by real time PCR after 6, 12, 24, 48 hours, and 4 and 30 days. At 6, 12, and 24 hours the plasmid DNA content of the injected lymph nodes was approximately three orders of magnitude greater than that of the draining lymph nodes following i.m. injection. No plasmid DNA was detectable in the draining lymph node at subsequent time points (Fig. 17). This is consonant with the three orders of magnitude greater dose needed using i.m. as compared to i.ln. injections to achieve a similar levels of CTL activity. CD8^{-/-} knockout mice, which do not develop a CTL response to this epitope, were also injected i.ln. showing clearance of DNA from the lymph node is not due to CD8⁺ CTL killing of cells in the lymph node. This observation also supports the conclusion that i.ln. administration will not provoke immunopathological damage to the lymph node.

Example 13

Administration of a DNA plasmid formulation of a therapeutic vaccine for melanoma to humans.

SYNCHROTOPE TA2M, a melanoma vaccine, encoding the HLA-A2-restricted tyrosinase epitope SEQ ID NO. 1 and epitope cluster SEQ ID NO. 69, was formulated in 1% Benzyl alcohol, 1% ethyl alcohol, 0.5mM EDTA, citrate-phosphate, pH 7.6. Aliquots of 80, 160, and 320 µg DNA/ml were prepared for loading into MINIMED 407C infusion pumps. The catheter of a SILHOUETTE infusion set was placed into an inguinal lymph node visualized by ultrasound imaging. The assembly of pump and infusion set was originally designed for the delivery of insulin to diabetics and the usual 17mm catheter was substituted with a 31mm catheter for this application. The infusion set was kept patent for 4 days (approximately 96 hours) with an infusion rate of about 25 µl/hour resulting in a total infused volume of approximately 2.4 ml. Thus the total administered dose per infusion was approximately 200, and 400 µg; and can be 800 µg, respectively, for the three concentrations described above. Following an infusion subjects were given a 10 day rest period before starting a subsequent infusion. Given the continued residency of plasmid DNA in the lymph node after administration (as in example 12) and the usual kinetics of CTL response following disappearance of antigen, this schedule will be sufficient to maintain the immunologic CTL response.

Example 14**Additional Epitopes.**

The methodologies described above, and in particular in examples 3-7, have been applied to additional synthetic peptide substrates, leading to the identification of further epitopes as set for the in tables 15-36 below. The substrates used here were designed to identify products of housekeeping proteasomal processing that give rise to HLA-A*0201 binding epitopes, but additional MHC-binding reactivities can be predicted, as discussed above. Many such reactivities are disclosed, however, these listings are meant to be exemplary, not exhaustive or limiting. As also discussed above, individual components of the analyses can be used in varying combinations and orders. The digests of the NY-ESO-1 substrates 136-163 and 150-177 (SEQ ID NOS. 254 and 255, respectively) yielded fragments that did not fly well in MALDI-TOF mass spectrometry. However, they were quite amenable to N-terminal peptide pool sequencing, thereby allowing identification of cleavage sites. Not all of the substrates necessarily meet the formal definition of an epitope cluster as referenced in example 3. Some clusters are so large, e.g. NY-ESO-1₈₆₋₁₇₁, that it was more convenient to use substrates spanning only a portion of this cluster. In other cases, substrates were extended beyond clusters meeting the formal definition to include neighboring predicted epitopes. In some instances, actual binding activity may have dictated what substrate was made, as with for example the MAGE epitopes reported here, where HLA binding activity was determined for a selection of peptides with predicted affinity, before synthetic substrates were designed.

Table 15
GP100: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion
 †Scores are given from the two binding prediction programs referenced above (see example 3).

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI/NIH)†					Comments
				A*0201	A1	A3	B7	B8	
609-644	630-638*	LPHSSSHWL	88				20/80	16/<5	*The digestion of 609-644 and 622-650 have generated the same epitopes.
	629-638*	QLPHSSSHWL	89	21/117					
	614-622	LIYRRRLMK	90			32/20			
	613-622	SLTYRRRLMK	91	14/<5		29/60			
	615-622	IYRRRLMK	92					15/<5	
622-650	630-638*	LPHSSSHWL	93				20/80	16/<5	
	629-638*	QLPHSSSHWL	94	21/117					

Table 16A

MAGE-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI/NIH) [†]						
				A*0201	A1	A3	B7	B8	Other	
86-109	95-102	ESLFRAVI	95					16/<5		
	93-102	ILESFRVI	96	21/<5		20/<5				
	93-101	ILESFRV	97	23/<5						
	92-101	CILESFRV	98	23/55						
	92-100	CILESFR	99	20/138						
	263-271	EFLWGPRAL	100						A26 (R 21), A24 (NIH 30)	
263-292	264-271	FLWGPRAL	101					17/<5		
	264-273	FLWGPRALAE	102	16/<5		19/<5				
	265-274	LWGPRALAE	103	16/<5						
	268-276	PRALAEYS	104	15/<5						
	267-276	GPRALAEYS	105	15/<5			<15/<5		B4403 (NIH 7); B3501 (NIH 120)	
	269-277	RALAEYSV	106	18/20						
	271-279	LAETSYYKV	107	19/<5						
	270-279	ALAETSYYKV	108	30/427		19/<5<5				
	272-280	AETSYYKVL	109	15/<5					B4403 (NIH 36)	
	271-280	LAETSYYKVL	110	18/<5			<15/<5			
	274-282	TSYYVKVLEY	111		26/<5				B4403 (NIH 14)	
	273-282	ETSYVKVLEY	112		28/6				A26 (R 31), B4403 (NIH 14)	
	278-286	KVLEYVVKV	113	26/743		16/<5				

Table 16B
MAGE-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI/NIH) [†]					
				A*0201	A1	A3	B7	B8	Other
168-193	168-177	SYVLVTCLGL	114						A24 (NIH 300)
	169-177	YVLVTCLGL	115	20/32		15/<5	<15/20		
	170-177	VLVTCLGL	116					17/<5	
	240-248	TQDLVQEKY	117		29/<5				
229-258	239-248	LTQDLVQEKY	118		23/<5				A26 (R 22)
	232-240	YGEPRKLLT	119		24/11				
	243-251	LVQEKYLEY	120		21/<5	21/<5			A26 (R 28)
	242-251	DLVQEKYLEY	121		22/<5	19/<5			A26 (R 30)
	230-238	SAYGEPRKL	122	21/<5					B5101 (25/121)
	278-286	KVLEYVIVK	123	26/743		16/<5			
	277-286	VKVLEYVIVK	124	17/<5					
272-297	276-284	YVKVLEYVI	125	15/<5		15/<5		17/<5	
	274-282	TSYVKVLEY	126		26/<5				
	273-282	ETSYVKVLEY	127		28/6				
	283-291	VIKVSARVR	128			20/<5			
	282-291	YVIKVSARVR	129			24/<5			

[†]Scores are given from the two binding prediction programs referenced above (see example 3). R indicates a SYFPEITHI score.

Table 17A

MAGE-2: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

MAGE-2: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion									
Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI /NIH)†					
				A*0201	A1	A3	B7	B8	Other
107-126	115-122	ELVHFLLL	130					18/<5	
	113-122	MVELVHFLLL	131		21/<5				A26 (R 22)
	109-116	ISRKMVEL	132					17/<5	
	108-116	AISRKMVEL	133	25/7		19/<5	16/12	26/<5	
	107-116	AAISRKMVEL	134	22/<5			14/36	n.p./16	
	112-120	KMVELVHFL	135	27/2800					
	109-117	ISRKMVELV	136	16/<5					
	108-117	AISRKMVELV	137	24/11					
	116-124	LVHFLLLY	138		23/<5	19/<5			A26 (R 26)
	115-124	ELVHFLLLY	139		24/<5	19/5			A26 (R 29)
	111-119	RKMVELVHF	140						
	158-166	LQLVFGIEV	141	17/168					
	157-166	YLQLVFGIEV	142	24/1215					
	159-167	QLVFGIEVV	143	25/32		18/<5			
145-175	158-167	LQLVFGIEVV	144	18/20					
	164-172	IEVVEVYPI	145	16/<5					
	163-172	GIEVVEVYPI	146	22/<5					
	162-170	FGIEVVEVV	147	19/<5					B5101(24/69.212)
	154-162	ASEYLQLVF	148		22/68				
	153-162	KASEYLQLVF	149			15/<5			

[†]Scores are given from the two binding prediction programs referenced above (see example 3). R indicates a SYFPEITHI score.

Table 17B
MAGE-2: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence		HLA Binding Predictions (SYFPEITHI /NIH) [†]					
				A*0201	A1	A3	B7	B8	Other
213-233	218-225	EEKIWEEL	150					22/<5	
	216-225	APEEKIWEEL	151	15/<5			22/72		
	216-223	APEEKIWE	152					18/<5	
	220-228	KIWEEL SML	153	26/804		16/<5		16/<5	A26 (R 26)
	219-228	EKIWEEL SML	154						A26 (R 22)
271-291	271-278	FLWGPRAL	155					17/<5	
	271-279	FLWGPRALI	156	25/398		16/7			
	278-286	LIETSYVKV	157	23/<5					
	277-286	ALIETSYVKV	158	30/427		21/<5			
	276-284	RALIETSYV	159	18/19					B5101 (20/55)
	279-287	IETSYVKVL	160	15/<5					
	278-287	LIETSYVKVL	161	22/<5					A26 (R 22)

[†]Scores are given from the two binding prediction programs referenced above (see example 3). R indicates a SYFPEITHI score.

Table 18
MAGE-3: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI / NIH) [†]					
				A*0201	A1	A3	B7	B8	Other
267-286	271-278	FLWGPRAL	162					17/<5	
	270-278	EFLWGPRAL	163						A26 (R 21); A24 (NIH 30)
	271-279	FLWGPRALV	164	27/2655		16/<5			
	276-284	RALVETSYV	165	18/19					B5101 (20/55)
	272-280	LWGPRALVE	166			15/<5			
	271-280	FLWGPRALVE	167	15/<5		22/<5			
	272-281	LWGPRALVET	168	16/<5					

[†]Scores are given from the two binding prediction programs referenced above (see example 3). R indicates a SYFPEITHI score.

Table 19A
NY-ESO-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI /NIH) [†]							
				A*0201	A1	A3	B7	B8	Other	Comment	
81-113	82-90	GPESRLLEF	169		16/11		18/<5	22/<5			
	83-91	PESRLLEFY	170		15/<5				B4403 (NIH 18)		
	82-91	GPESRLLEFY	171		25/11						
	84-92	ESRLLEFY	172					19/8			
	86-94	RLLEFYLAM	173	21/430		21/<5					
	88-96	LEFYLAMPF	174						B4403 (NIH 60)		
	87-96	LEFYLAMPF	175		<15/45	18/<5					
	93-102	AMPEATPMEA	176	15/<5							
	94-102	MPFATPMEA	177				17/<5				
	115-123	PLPVPGVLL	178	20/<5		17/<5	16/<5	18/<5			
101-133	114-123	PPLPVPGVLL	179				23/12			*Evidence of the same epitope obtained from two digests.	
	116-123*	LPVPGVLL	180					16/<5			
	103-112	ELARRSLAQD	181	15/<5		20/<5					
	118-126*	VPGVLLKEF	182				17/<5	16/<5			
	117-126*	PVPGVLLKEF	183			16/<5					
116-145	116-123*	LPVPGVLL	184					16/<5			
	127-135	TVSGNLT	185	21/<5		19/<5					
	126-135	FTVSGNLT	186	20/<5							
	120-128	GVLLKEFTV	187	20/130		18/<5					
	121-130	VLLKEFTVSG	188	17/<5		18/<5					
	122-130	LLKEFTVSG	189	20/<5		18/<5					
	118-126*	VPGVLLKEF	190				17/<5	16/<5			
	117-126*	PVPGVLLKEF	191			16/<5					

[†]Scores are given from the two binding prediction programs referenced above (see example 3).

Table 19B
 NY-ESO-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI/NIH) [†]					
				A*0201	A1	A3	B7	B8	Other
136-163 (SEQ ID NO 254)	139-147	AADHRQLQL	192	17/<5	17/<5			22/<5	
	148-156	SISSCLQQL	193	24/7					A26 (R 25)
	147-156	LSISSCLQQL	194	18/<5					
	138-147	TAADHRQLQL	195	18/<5					
	161-169	WITQCFLPV	196	18/84					
150-177 (SEQ ID NO 255)	157-165	SLLMWITQC	197	18/42		17/<5			
	150-158	SSCLQQLSL	198	15/<5					
	154-162	QQLSLLMWI	199	15/50					
	151-159	SCLQQLSLL	200	18/<5					
	150-159	SSCLQQLSLL	201	16/<5					
	163-171	TQCFLPVFL	202	<15/12					
	162-171	ITQCFLPVFL	203	18/<5					A26 (R 19)

[†]Scores are given from the two binding prediction programs referenced above (see example 3). R indicates a SYFPEITHI score

Table 20
PRAME: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI /NIH) [†]					
				A*0201	A1	A3	B7	B8	Other
211-245	219-227	PMQDIKMIL	204	16/<5				16/n.d.	A26 (R 20)
	218-227	MPMQDIKMIL	205				<15/240		
411-446	428-436	QHLIGLSNL	206	18/<5					
	427-436	LQHLIGLSNL	207	16/8					
	429-436	HLIGLSNL	208					17/<5	B15 (R 21)
	431-439	IGLSNLTHV	209	18/7					B*5101 (R 22)
	430-439	LIGLSNLTHV	210	24/37					

[†]Scores are given from the two binding prediction programs referenced above (see example 3). R indicates a SYFPEITHI score.

Table 21
PSA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI /NID) [†]					
				A*0201	A1	A3	B7	B8	Other
42-77	53-61	VLVHPQWVL	211	22/112			<15/6	17/<5	
	52-61	GVLVHPQWVL	212	17/21		16/<5	<15/30		A26 (R 18)
	52-60	GVLVHPQWV	213	17/124					
	59-67	WVLTAAHCI	214	15/16					
	54-63	LVHPQWVLTA	215	19/<5		20/<5			A26 (R 16)
	53-62	VLVHPQWVLT	216	17/22					
	54-62	LVHPQWVLT	217			17/n.d.			
55-95	66-73	CIRNKSVI	218					26/20	
	65-73	HCIRNKSVI	219					<15/16	
	56-64	HPQWVLTA	220				18/<5		
	63-72	AAHCIRNKSV	221	17/<5					

[†]Scores are given from the two binding prediction programs referenced above (see example 3). R indicates a SYFPEITHI score.

Table 22
PSCA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI /NIEH) [†]					
				A*0201	A1	A3	B7	B8	Other
93-123*	116-123	LLWGPQQL	222					16/<5	
	115-123	LLLWGPQQL	223	<15/18					
	114-123	GLLWGPQQL	224	<15/10					
	99-107	ALQPAAAIL	225	26/9		22/<5	<15/12	16/<5	A26 (R 19)
	98-107	HALQPAAAIL	226	18/<5			<15/12		

*L123 is the C-terminus of the natural protein.

[†]Scores are given from the two binding prediction programs referenced above (see example 3).

Table 23
Tyrosinase: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI /NIH) [†]					
				A*0201	A1	A3	B7	B8	Other
128-157	128-137	APEKDKFFAY	227		29/6		15/<5		B4403 (NIH 14)
	129-137	PEKDKFFAY	228		18/<5			21/<5	
	130-138	EKDKFFAYL	229				15/<5		
	131-138	KDKFFAYL	230					20/<5	
	205-213	PAFLPWHL	231					15/<5	
197-228	204-213	APAFLPWHL	232				23/360		
	207-216	FLPWHLFL	1	25/1310				<15/8	
	208-216	LPWHLFL	9	17/26			20/80	24/16	
	214-223	FLLRWEQEI	233			15/<5			
	212-220	RLFLLRWEQ	234			16/<5			
191-211	191-200	GSEIWRDIDF	235		18/68				
	192-200	SEIWRDIDF	236					16/<5	B4403 (NIH 400)
	207-215	FLWHLRLFL	8	22/540			<15/6	17/<5	
207-230	473-481	RIWSWLIGA	237	19/13		15/<5			
466-484	476-484	SWLLGAAMV	238	18/<5					
	477-486	WLLGAAMVGA	239	21/194		18/<5			
	478-486	LLGAAMVGA	240	19/19		16/<5			

[†]Scores are given from the two binding prediction programs referenced above (see example 3).

Table 24
PSMA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	SEQ ID NO	HLA Binding Predictions (SYFPEITHI /NIH)†					
				A*0201	A1	A3	B7	B8	Other
1-30	4-12	LLHETDSAV	241	25/485		15/<5			
	13-21	ATARRPRWL	242	18/<5				18/<5	A26 (R 19)
	53-61	TPKHNMKAF	243					24/<5	
53-80	64-73	ELKAENIKKF	244			17/<5			A26 (R 30)
	69-77	NIKKFLH'NF	245						A26 (R 27)
	68-77	ENIKKFLH'NF	246						A26 (R 24)
	220-228	AGAKGVILY	247		25/<5				
457-489	468-477	PLMYSLVHNL	248	22/<5					
	469-477	LMYSLVHNL	249	27/193		<15/9			
	463-471	RVDCITPLMY	250		32/125	25/<5			A26 (R 22)
	465-473	DCTPLMYSL	251						A26 (R 22)
503-533	507-515	SGMPRISKL	252	21/<5				21/<5	
	506-515	FSGMPRISKL	253	17/<5					

[†] This H was reported as Y in the SWISSPROT database.

+Scores are given from the two binding prediction programs referenced above (see example 3).

Table 25A
MAGE-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
Mage-1 119-146	125-132	KAEMLESV	256	B5101	19	n.a.
	124-132	TKAEMLESV	257	A0201	20	<5
	123-132	VTKAEMLESV	258	A0201	20	<5
				A1	28	45
	128-136	MLESVIKNY	259	A26	24	n.a.
				A3	17	5
	127-136	EMLESVIKNY	260	A1	15	<1.0
				A26	23	<1.0
	125-133	KAEMLESVI	261	B5101	23	100
				A24	N.A.	4
Mage-1 143-170	146-153	KASESLQL	262	B08	16	<1.0
				B5101	17	N.A.
	145-153	GKASESLQL	263	B2705	17	1
	147-155	ASESLQLVF	264	B2709	16	N.A.
				A1	22	68
	153-161	LVFGIDVKE	265	A26	16	N.A.
				A3	16	<1.0

Table 25B MAGE-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion						
Substrate	Epitope	Sequence	Seq. ID No.	Blinding Prediction		
				HLA type	SYFPEITHI	NIH
Mage-1 99-125	114-121	LLKYRARE	266	B8	25	<1.0
	106-113	VADLVGFL	267	B8	16	<1.0
				B5101	21	N.A.
				A0201	23	44
	105-113	KVADLVGFL	268	A26	25	N.A.
				A3	16	<5
				B0702	14	20
	107-115	ADLVGFLLL	269	B2705	14	30
				A0201	17	<5
				B0702	15	<5
	106-115	VADLVGFLLL	270	B2705	16	1
				A0201	16	<5
				A1	22	3
114-123	LLKYRAREPV	271	A0201	20	2	

Table 26
MAGE-3: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
Mage-3 267-295	271-278	FLWGPRAL	162	B08	17	<5
	270-278	EFLWGPRAL	163	A26	21	N.A.
				A24	N.A.	30
				B1510	16	N.A.
	271-279	FLWGPRALV	164	A0201	27	2655
				A3	16	2
				A0201	19	<1.0
	278-286	LVETSYVKV	272	A26	17	N.A.
				A0201	28	428
				A26	16	<5
	277-286	ALVETSYVKV	273	A3	18	<5
				A0201	19	27
				A3	19	<5
	285-293	KVLHLMVKI	274	A0201	18	20
	276-284	RALVETSYV	165	A0201	17	<1.0
	283-291	YVKVLHLMV	275	A0201	17	<1.0
	275-283	PRALVETSY	276	A1	15	<1.0
	274-283	GPRALVETSY	277	A1	18	<1.0
	278-287	LVETSYVKVL	278	A0201	16	<1.0
	272-281	LWGPRALVET	168	A0201	22	<5
	271-280	FLWGPRALVE	167	A3		

Table 27A
Fibronectin ED-B: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
ED-B 14'-21*	4'-5**	<i>TI</i> PEVPQL [†]	279	A0201	27	7
				A26	28	N.A.
				A3	17	<5
				B8	15	<5
				B1510	15	N.A.
				B2705	17	10
				B2709	15	N.A.
				A0201	20	<5
	5'-5**	<i>DTI</i> PEVPQL [†]	280	A26	32	N.A.
				A26	29	N.A.
	1-10	EVPQLTDLSE	281			

*This substrate contains the 14 amino acids from fibronectin flanking ED-B to the N-terminal side.

**These peptides span the junction between the N-terminus of the ED-B domain and the rest of fibronectin.

† The *italicized* lettering indicates sequence outside the ED-B domain.

Table 27B
Fibronectin ED-B: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
ED-B 8-35	23-30	TPLNSSTI	282	B5101	22	N.A.
	18-25	IGLRWTPL	283	B5101	18	N.A.
	17-25	SIGLRWTPL	284	A0201	20	5
				A26	18	N.A.
				B08	25	<5
				A1	19	<5
	25-33	LNSSTIIGY	285	A26	16	<5
				A1	20	<5
				A26	24	N.A.
	24-33	PLNSSTIIGY	286	A3	16	<5
				B0702	17	8
	23-31	TPLNSSTI	287	B5101	25	440

Table 27C
Fibronectin ED-B: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
ED-B 20-49	31-38	IGYRITVV	288	B5101	25	N.A.
				A0201	23	15
				A3	17	<1.0
	30-38	IIGYRITVV	289	B08	15	<1.0
				B5101	15	3
				A0201	26	9
	29-38	TIIGYRITVV	290	A26	18	N.A.
				A3	18	<5
	23-30	TPLNSSTI	282	B5101	22	N.A.
				A1	19	<5
	25-33	LNSSTIIGY	285	A26	16	N.A.
				A26	24	N.A.
	24-33	PLNSSTIIGY	286	A3	16	<5
	31-39	IGYRITVVA	291	A3	17	<5
				A0201	15	<5
	30-39	IIGYRITVVA	292	A3	18	<5
				B0702	17	8
	23-31	TPLNSSTII	287	B5101	25	440

Table 28A CEA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion						
Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
CEA 176-202	184-191	SLPVSPRL	293	B08	19	<5
				A0201	15	<5
	183-191	QSLPVSPRL	294	B1510	15	
				B2705	18	10
				B2709	15	
	186-193	PVSPRLQL	295	B08	18	<5
				B0702	26	180
	185-193	LPVSPRLQL	296	B08	16	<5
				B5101	19	130
	184-193	SLPVSPRLQL	297	A0201	23	21
				A26	18	N.A.
				A3	18	<5
	185-192	LPVSPRLQ	298	B5101	17	N.A.
				A0201	21	4
	192-200	QLSNGNRTL	299	A26	16	N.A.
				A3	19	<5
				B08	17	<5
				B1510	15	
	191-200	LQLSNGNRTL	300	A0201	16	3
	179-187	WVNNQSLPV	301	A0201	16	28
	186-194	PVSPRLQLS	302	A26	17	N.A.
A3				15	<5	

Table 28B
CEA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYTPEITHI	NIH
CEA 354-380	362-369	SLPVSRL	303	B08	19	<1.0
	361-369	QSLPVSRL	304	A0201	15	<1.0
				B2705	18	10
	364-371	PVSRLQL	305	B2709	15	
				B08	18	<1.0
	363-371	LPVSPRLQL	306	B0702	26	180
				B08	16	<1.0
				B5101	19	130
	362-371	SLPVSRLQL	307	A0201	23	21
				A26	18	N.A.
				A24	N.A.	6
				A3	18	<5
	363-370	LPVSPRLQ	308	B5101	17	N.A.
	370-378	QLSNDNRIL	309	A0201	22	4
				A26	16	N.A.
				A3	17	<1.0
				B08	17	<1.0
	369-378	LQLSNDNRIL	310	A0201	16	3
	357-365	WVNNQSLPV	311	A0201	16	28
	360-368	NQSLPVSPR	312	B2705	14	100

Table 28C
CEA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NTH
CEA 532-558	540-547	SLPVPRL	313	B08	19	<5
	539-547	QSLPVPRL	314	A0201	15	<5
				B1510	15	<5
				B2705	18	10
				B2709	15	
	542-549	PVSPRLQL	315	B08	18	<5
	541-549	LPVSPRLQL	316	B0702	26	180
				B08	16	<1.0
				B5101	19	130
	540-549	SLPVPRLQL	317	A0201	23	21
				A26	18	N.A.
				A3	18	<5
	541-548	LPVSPRLQ	318	B5101	17	N.A.
	548-556	QLSNGNRIL	319	A0201	24	4
				A26	16	N.A.
				A3	19	<1.0
				B08	17	<1.0
				B1510	15	
	547-556	LQLSNGNRIL	320	A0201	16	3
	535-543	WVNGQSLPV	321	A0201	18	28
				A3	15	<1.0
	533-541	LWWVNGQSL	322	A0201	15	<5

Table 28D
CEA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
CEA 532-558 (continued)	532-541	YLWWVNGQSL	323	A0201	25	816
				A26	18	N.A.
	538-546	GQSLPVSPR	324	B2705	17	100

Table 29A
HER2/NEU: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
Her-2 25-52	30-37	DMKRLRLPA	325	B08	19	8
	28-37	GTDMLRLRLPA	326	A1	23	6
	42-49	HLDMLRHL	327	B08	17	<5
	41-49	THLDMLRHL	328	A0201	17	<5
	40-49	ETHLDMLRHL	329	B1510	24	N.A.
	36-43	PASPETHL	330	A26	29	N.A.
				B5101	17	N.A.
				A0201	15	<5
	35-43	LPASPETHL	331	B5101	20	130
				B5102	N.A.	100
	34-43	RLPASPETHL	332	A0201	20	21
				A0201	15	<5
	38-46	SPETHLDML	333	B0702	20	24
				B08	18	<5
				B5101	18	110
	37-46	ASPETHLDML	334	A0201	18	<5
				A1	29	25
	42-50	HLDMLRHL	335	A26	20	N.A.
				A3	17	4
	41-50	THLDMLRHL	336	A1	18	<1.0

Table 29B
HER2/NEU: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
Her-2 705-732	719-726	ELRKVKVL	337	B08	24	16
				A0201	16	1
	718-726	TELKVKVL	338	B08	22	<5
				B5101	16	<5
	717-726	ETELRKVKVL	339	A1	18	2
				A26	28	6
	715-723	LKETELRKV	340	A0201	17	<5
				B5101	15	<5
	714-723	ILKETELRKV	341	A0201	29	8
				A0201	15	<5
	712-720	MRILKETEL	342	B08	22	<5
				B2705	27	2000
				B2709	21	N.A.
	711-720	QMRILKETEL	343	A0201	20	2
				B0702	13	40
	717-725	ETELRKVKV	344	A1	18	5
				A26	18	N.A.
	716-725	KETELRKVKV	345	A0201	16	19
	706-714	MPNQAQMRI	346	B0702	16	8
				B5101	22	629
	705-714	AMPNQAQMRI	347	A0201	18	8
	706-715	MPNQAQMRL	348	B0702	20	80

Table 29C
HER2/NEU: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
Her-2 954-982	966-973	RPRFREL V	349	B08	20	24
	965-973	CRPRFREL V	350	B5101	18	N.A.
				B2709	18	
				A26	25	N.A.
	968-976	RFREL VSEF	351	A24	N.A.	32
				A3	15	<5
				B08	16	<5
				B2705	19	
	967-976	PRFREL VSEF	352	A26	18	N.A.
				A26	21	N.A.
				A24	N.A.	6
	964-972	ECRPRFREL	353	B0702	15	40
				B8	27	640
				B1510	16	<5

Table 30
NY-ESO-1: Preferred Epitopes Revealed by Housckkeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
NY-ESO-1 51-77	67-75	GAASGLNGC	354	A0201	15	<5
	52-60	RASGPGGGA	355	B0702	15	<5
	64-72	PHGGAASGL	356	B1510	21	N.A.
	63-72	GPHGGAASGL	357	B0702	22	80
	60-69	APRGPHGGAA	358	B0702	23	60

Table 31A
PRAME: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
PRAME 103-135	112-119	VRPRRWKL	359	B08	19	
	111-119	EVRPRRWKL	360	A26	27	N.A.
				A24	N.A.	5
				A3	19	N.A.
				B0702	15	(B7) 300.00
				B08	26	160
	113-121	RPRRWKLQV	361	B0702	21	(B7) 40.00
	114-122	PRRWKLQVL	362	B5101	19	110
				B08	26	<5
				B2705	23	200
	113-122	RPRRWKLQVL	363	B0702	24	(B7) 800.00
				B8	N.A.	160
				B5101	N.A.	61
				B5102	N.A.	61
				A24	N.A.	10
PRAME 161-187	116-124	RWKLQVLDL	364	B08	22	<5
	115-124	RRWKLQVLDL	365	B2705	17	3
	174-182	PVEVLVDLF	366	A0201	16	<5
				A26	25	N.A.

Table 31B
PRAME: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
PRAME 185-215	199-206	VKRKKNVL	367	B08	27	8
	198-206	KVKRKKNVL	368	A0201	16	<1.0
				A26	20	N.A.
				A3	22	<1.0
				B08	30	40
	197-206	EKVKRKKNVL	369	B2705	16	
				A26	15	N.A.
				B08	20	6
				B08	20	<5
	200-208	KRKKNVRL	372	A0201	15	<1.0
				A26	15	N.A.
				B0702	15	<1.0
				B08	21	<1.0
	199-208	VKRKKNVRL	373	B2705	28	
				B2709	25	
				A0201	16	<1.0
				B0702	16	4
PRAME 185-215	189-196	DELFSYLI	374	B5101	15	N.A.
				A0201	22	3
	205-213	VLRLOCKKL	375	A26	17	N.A.
				B08	25	8
	204-213	NVLRLOCKKL	376	A0201	17	7
				A26	19	N.A.

Table 31C

PRAME: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
PRAME 185-215 (continued)	194-202	YLIEKVKRK	377	A0201	20	<1.0
				A26	18	N.A.
				A3	25	68
				B08	20	<1.0
				B2705	17	
PRAME 71-98	74-81	QAWPFTCL	378	B5101	17	n.a.
	73-81	VQAWPFTCL	379	A0201	14	7
				A24	n.a.	5
				B0702	16	6
	72-81	MVQAWPFTCL	380	A26	22	n.a.
				A24	n.a.	7
				B0702	13	30
	81-88	LPLGVLMK	381	B5101	18	n.a.
	80-88	CLPLGVLMK	382	A0201	17	<1.0
				A3	27	120
				A1	12	10
	79-88	TCLPLGVLMK	383	A3	19	3
				A0201	18	7
				A26	21	n.a.
	84-92	GVLKMGQHL	384	B08	21	4
	81-89	LPLGVLMKG	385	B5101	20	2
	80-89	CLPLGVLMKG	386	A0201	16	<1.0
	76-85	WPFTCLPLGV	387	B0702	18	4

Table 31D
PRAME: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
PRAME 39-65	51-59	ELFPPLFMA	388	A0201	19	18
				A26	23	N.A.
	49-57	PRELFPPLF	389	B2705	22	
				B2709	19	
	48-57	LPRELFPPLF	390	B0702	19	4
				B2705	16	
	50-58	RELFPPLFM	391	B2705	15	
	49-58	PRELFPPLFM	392	A1	16	<1.0

Table 32
PSA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
PSA 232-258	239-246	RPSLYTKV	393	B5101	21	N.A.
	238-246	ERPSLYTKV	394	B2705	15	60
	236-243	LPERPSLY	395	B5101	18	N.A.
	235-243	ALPERPSLY	396	A1	19	<1.0
				A26	22	N.A.
				A3	26	6
				B08	16	<1.0
				B2705	11	15
				B2709	19	N.A.
	241-249	SLYTKVVHY	397	A0201	20	<1.0
				A1	19	<1.0
				A26	25	N.A.
				A3	26	60
				B08	20	<1.0
				B2705	13	75
	240-249	PSLYTKVVHY	398	A1	20	<1.0
	239-247	RPSLYTKVV	399	A26	16	N.A.
				B0702	21	4
				B5101	23	110

Table 33A

PSMA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
PSMA 202-228	211-218	GNKVKNAQ	400	B08	22	<5
	202-209	IARYGKVF	401	B08	18	<5
	217-225	AQLAGAKGV	402	A0201	16	26
	207-215	KVFRGNKVK	403	A3	32	15
	211-219	GNKVKNAQL	404	B8	33	80
PSMA 255-282	269-277	TPGYPANEY	405	B2705	17	20
	268-277	LTPGYPANEY	406	A1	16	<5
	271-279	GYPANEYAY	407	A1	21	1
	270-279	PGYPANEYAY	408	A26	24	N.A.
	266-274	DPLTPGYPA	409	A1	15	<5
PSMA 483-509	492-500	SLYESWTKK	410	A1	19	<5
	491-500	KSLYESWTKK	411	B0702	21	3
	486-494	EGFEGKSLY	412	B5101	17	20
	485-494	DEGFEGKSLY	413	A0201	17	<5
	498-506	TKKSPSPEF	414	A3	27	150
				B2705	18	150
				A3	16	<5

Table 33B
PSMA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Blinding Prediction		
				HLA type	SYFPEITHI	NIH
PSMA 483-509 (continued)	497-506	WTKKSPSEF	415	A26	24	N.A.
	492-501	SLYESWTKKS	416	A0201	16	<5
				A3	16	<5
				B08	17	<5
	725-732	WGEVKRQI	417	B5101	17	N.A.
	724-732	AWGEVKRQI	418	B5101	15	6
	723-732	KAWGEVKRQI	419	A0201	16	<1.0
	723-730	KAWGEVKR	420	B5101	15	N.A.
	722-730	SKAWGEVKR	421	B2705	15	<5
	731-739	QIYVAAFTV	422	A0201	21	177
				A3	21	<1.0
				B5101	15	5
PSMA 721-749	733-741	YVAAFTVQA	423	A0201	17	6
	725-733	WGEVKRQIY	424	A3	20	<1.0
				A1	26	11
	727-735	EVKRQIYVA	425	A26	22	N.A.
				A3	18	<1.0
	738-746	TVQAAAETL	426	A26	18	N.A.
				A3	19	<1.0
	737-746	FIVQAAAETL	427	A0201	17	<1.0
				A26	19	N.A.

Table 33C

PSMA: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
PSMA 721-749 (continued)	729-737	KRQIYVAAF	428	A26	16	N.A.
				B2705	24	3000
				B2709	21	N.A.
	721-729	PSKAWGEVK	429	A3	20	<1.0
PSMA 95-122	723-731	KAWGEVKRQ	430	B5101	16	<1.0
	100-108	WKEFGDSV	431	A0201	16	<5
	99-108	QWKEFGDSV	432	A0201	17	<5
	102-111	EFGLDSVELA	433	A26	16	N.A.

Table 34A
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 117-143	126-134	ELRQKESKL	434	A0201	20	<5
				A26	26	N.A.
				A3	17	<5
				B0702	13	(B7) 40.00
				B8	34	320
SCP-1 281-308	125-134	AELRQKESKL	435	A0201	16	<5
	133-141	KIQENRKII	436	A0201	20	61
	298-305	QLEEKTKL	437	B08	28	2
	297-305	NQLEEKTKL	438	A0201	16	33
				B2705	19	200
	288-296	LLEESRDKV	439	A0201	25	15
				B5101	15	3
	287-296	FLLEESRDKV	440	A0201	27	2378
	291-299	ESRDKNVQL	441	A26	21	N.A.
				B08	29	240
SCP-1 471-498	290-299	EESRDKNVQL	442	A26	19	N.A.
	475-483	EKEVHDLEY	443	A1	31	11
				A26	17	N.A.
	474-483	REKEVHDLEY	444	A1	21	<1.0
	480-488	DLEYSYCHY	445	A1	26	45
				A26	30	N.A.
	477-485	EVHDLEYSY	446	A3	16	<5
				A1	15	1

Table 34B

SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 471-498 (continued)	477-485	EVHDLQSY		A26	29	N.A.
	477-486	EVHDLQSY		A3	19	<1.0
	502-509	KLSSKREL	447	A26	22	N.A.
	508-515	ELKNTQYF	448	B08	26	4
			449	B08	24	<1.0
	507-515	RELKNTQYF	450	B2705	18	45
	496-503	KRGQRPKL	451	B4403	N.A.	120
				B08	18	<1.0
				B0702	22	120
	494-503	LPKRGQRPKL	452	B8	N.A.	16
SCP-1 493-520				B5101	N.A.	130
				B3501	N.A.	60
	509-517	LKNTQYFTL	453	A0201	15	<5
				A0201	18	<1.0
	508-517	ELKNTQYFTL	454	A26	27	N.A.
				A3	16	<1.0
	506-514	KRELKNTQY	455	A1	26	2
	502-510	KLSSKRELK	456	B2705	26	3000
	498-506	GORPKLSSK	457	A3	25	60
				A3	22	4
	497-506	RGQRPKLSSK	458	B2705	18	200
	500-508	RPKLSSKRE	459	A3	22	<1.0
				B08	18	<1.0

Table 34C
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 570-596	573-580	LEYVREEL	460	B08	19	<5
	572-580	ELEYVREEL	461	A0201	17	<1.0
				A26	23	N.A.
				A24	N.A.	9
				B08	20	N.A.
	571-580	N ELEYVREEL	462	A0201	16	4
	579-587	ELKQKRDEV	463	A0201	19	<1.0
				A26	18	N.A.
				B08	29	48
	575-583	YVREELKQK	464	A26	17	N.A.
SCP-1 618-645	632-640	QLNVYEIKV	465	A3	27	2
				A0201	24	70
	630-638	SKQLNVYEI	466	A0201	17	<5
	628-636	AESKQLNVY	467	A1	19	<5
				A26	16	N.A.
	627-636	TAESKQLNVY	468	A1	26	45
				A26	15	N.A.

Table 34D
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 633-660	638-645	IKVVKLEL	469	B08	21	<1.0
	637-645	EIKVVKLEL	470	A0201	17	<1.0
				A26	26	N.A.
				B08	28	8
				B1510	15	N.A.
	636-645	YEIKVVKLEL	471	A0201	17	2
	642-650	KLELEESA	472	A0201	20	1
				A3	16	<1.0
				A0201	18	<1.0
	635-643	VYEIKVNKL	473	A24	N.A.	396
				B08	22	<1.0
				A0201	24	56
				A26	25	N.A.
				A24	N.A.	6
				A3	15	<5
				B0702	11	(B7) 20
SCP-1 640-668	646-654	ELESAKQKF	475	B08	N.A.	6
	642-650	KLELEESA	476	A26	27	N.A.
				A0201	20	1
	646-654	ELESAKQKF	477	A3	16	<1.0
				A26	27	N.A.

Table 34E
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 768-796	771-778	KEKLRREA	478	B08	21	<5
				A0201	18	<5
				A26	18	N.A.
				A24	N.A.	5
	777-785	EAKENTATL	479	B0702	13	12
				B08	28	48
				B5101	20	121
	776-785	RBAKENTATL	480	A0201	16	<5
	773-782	KLKREAKENT	481	A3	17	<5
	112-119	EAEKKKW	482	B5101	17	N.A.
SCP-1 92-125				A0201	23	32
				A26	22	N.A.
	101-109	GLSRVYSKL	483	A24	N.A.	6
				A3	17	3
				B08	17	<1.0
	100-109	EGLSRVYSKL	484	A26	21	N.A.
				A24	N.A.	9
	108-116	KLYKEAEKI	485	A0201	22	57
				A3	20	9
				B5101	18	5
	98-106	NSEGLSRVY	486	A1	31	68
	97-106	ENSEGLSRVY	487	A26	18	N.A.
	102-110	LSRVYSKLY	488	A1	22	<1.0

Table 34F
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 92-125 (continued)	101-110	GLSRVYSKLY	489	A1	18	<1.0
				A26	18	N.A.
				A3	19	18
	96-105	LENSEGLSRV	490	A0201	17	5
SCP-1 931-958	108-117	KLYKEAEKIK	491	A3	27	150
	949-956	REDRWAVI	492	B5101	15	N.A.
				B2705	18	600
	948-956	MRDRWAVI	493	B2709	18	N.A.
				B5101	15	1
				A0201	21	6
	947-956	KMRDRWAVI	494	B08	N.A.	15
	947-955	KMRDRWAV	495	A0201	22	411
	934-942	TTPGSTLKF	496	A26	25	N.A.
	933-942	LTPGSTLKF	497	A26	23	N.A.
SCP-1 232-259	937-945	GSTLKFGAI	498	B08	19	1
	945-953	IRKMRDRW	499	B08	19	<5
	236-243	RLEMHFKL	500	B08	16	<5
				A0201	18	<5
	235-243	SRLEMHFKL	501	B2705	25	2000
				B2709	22	
	242-250	KLKEDYEKI	502	A0201	22	4

Table 34G
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 232-259 (continued)				A26	16	N.A.
				A3	15	3
				B08	24	<5
				B5101	14	2
				A1	15	<5
	249-257	KIQHLEQY	503	A26	23	N.A.
				A3	17	<5
	248-257	EKIQHLEQY	504	A1	15	<5
				A26	21	N.A.
	233-242	ENSRLEMHF	505	A26	19	N.A.
SCP-1 310-340	236-245	RLEMHFKLKE	506	A1	19	<5
				A3	17	<5
	324-331	LEDIKVSL	507	B08	20	<1.0
				A0201	21	<1.0
				A26	25	N.A.
	323-331	ELEDIKVSL	508	A24	N.A.	10
				A3	17	<1.0
				B08	19	<1.0
				B1510	16	N.A.
	322-331	KELEDIKVSL	509	A0201	19	22
SCP-1 310-340	320-327	LTKLEDI	500	B08	18	<5
	319-327	HLTKLEDI	511	A0201	21	<1.0
	330-338	SLQRSVSTQ	512	A0201	18	<1.0

Table 34H

SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 310-340 (continued)	321-329	TKELEDIKV	513	A1	16	<1.0
	320-329	LTKLEDIKV	514	A0201	19	<1.0
	326-335	DIKVSQRSV	515	A26	18	N.A.
	281-288	KMKDLTFL	516	B08	20	3
SCP-1 272-305	280-288	NKMKDLTFL	517	A0201	15	1
	279-288	ENKMKDLTFL	518	A26	19	N.A.
	288-296	LLEESRDKV	519	A0201	25	15
				B5101	15	3
	287-296	FLLEESRDKV	520	A0201	27	2378
	291-299	ESRDKVNQL	521	A26	21	N.A.
				B08	29	240
	290-299	EESRDKVNQL	522	A26	19	N.A.
	277-285	EKENKMKDL	523	A26	19	N.A.
				B08	23	<1.0
SCP-1 211-239	276-285	TEKENKMKDL	524	A26	15	N.A.
	279-287	ENKMKDLTF	525	A26	18	N.A.
				B08	28	4
	218-225	IEKMITAF	526	B08	17	<5
	217-225	NIEKMITAF	527	A26	26	N.A.
	216-225	SNIEKMITAF	528	A26	19	N.A.
	223-230	TAFEELRV	529	B5101	23	N.A.
	222-230	ITAFEELRV	530	A0201	18	2
	221-230	MITAFEELRV	531	A0201	18	16

Table 34f
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 211-239 (continued)	220-228	KMITAFEEL	532	A0201	23	50
				A26	15	N.A.
				A24	N.A.	16
	219-228	EKMITAFEEL	533	A26	19	N.A.
	227-235	ELRVQAENS	534	A3	16	<1.0
				B08	15	<1.0
	213-222	DLNSNIEKMI	535	A0201	17	<1.0
				A26	16	N.A.
	837-844	WTSAKNTL	536	B08	20	4
				A0201	18	2
SCP-1 836-863	846-854	TPLPKAYTV	537	B0702	17	4
				B08	16	2
				B5101	25	220
	845-854	STPLPKAYTV	538	A0201	19	<5
	844-852	LSTPLPKAY	539	A1	23	8
				A1	16	<1.0
	843-852	TLSTPLPKAY	540	A26	19	N.A.
				A3	18	2
	842-850	NLTSTPLPK	541	A3	16	3
	841-850	KNTLSTPLPK	542	A3	18	<1.0

Table 34J
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYRPEITHI	NIH
SCP-1 819-845	828-835	ISKDKRDY	543	B08	21	3
				A26	21	N.A.
	826-835	HGISKDKRDY	544	A1	15	<5
	832-840	KRDYLVWTS	545	B2705	16	600
	829-838	SKDKRDYLVW	546	A1	18	<5
SCP-1 260-288	279-286	ENKMKDLT	547	B08	22	8
				A0201	17	3
	260-268	EINDKEKQV	548	A26	19	N.A.
				B08	17	<5
	274-282	QITEKENKM	549	A0201	17	3
				A26	22	N.A.
				B08	16	<5
	269-277	SLLLIQITE	550	A0201	16	<1.0
				A3	18	<1.0
	453-460	FEKIAEEL	551	B08	21	<1.0
SCP-1 437-464	452-460	QFEKIAEEL	552	B2705	15	
	451-460	KQFEKIAEEL	553	A0201	16	56
	449-456	DNKQFEKI	554	B08	16	2
	448-456	YDNKQFEKI	555	B5101	16	N.A.
	447-456	LYDNKQFEKI	556	B5101	16	1
				A1	15	<1.0

Table 34K

SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 437-464 (continued)	440-447	LGEKETLL	557	B5101	16	N.A.
	439-447	VLGEKETLL	558	A0201	24	149
				A26	19	N.A.
				B08	29	12
	438-447	KVLGEKETLL	559	A0201	19	24
				A26	20	N.A.
				A24	N.A.	12
				A3	18	<1.0
				B0702	14	20
				A0201	22	3
SCP-1 383-412	390-398	LLRTEQQRL	560	A26	18	N.A.
				B08	22	1.6
				B2705	15	30
	389-398	ELLRTEQQRL	561	A0201	19	6
				A26	24	N.A.
				A3	15	<1.0
	393-401	TEQQRLNY	562	A1	15	<5
				A26	16	N.A.
	392-401	RTEQQRLNY	563	A1	31	113
				A26	26	N.A.
	402-410	EDQLILTM	564	A26	18	N.A.
	397-406	RLENYEDQLI	565	A0201	17	<1.0
				A3	15	<1.0

Table 34K
SCP-1: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SCP-1 366-394	368-375	KARAAHSF	566	B08	16	<1.0
	376-384	VVTEFETIV	567	A0201	19	161
				A3	16	<1.0
	375-384	FVVTEFETIV	568	A0201	17	106
	377-385	VTEFETIVC	569	A1	18	2
SCP-1 331-357	376-385	VVTEFETIVC	570	A3	16	<5
	344-352	DLQIATNTI	571	A0201	22	<5
				A3	15	<1.0
				B5101	17	11
	347-355	IATNTICQL	572	A0201	19	1
				B08	16	<1.0
				B5101	20	79
SCP-1 331-357	346-355	QIATNTICQL	573	A0201	24	7
				A26	24	N.A.

Table 35
SSX-4: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
SSX4 45-76	57-65	VMTKLGFKV	574	A0201	21	495
	53-61	LNVEVMTKL	575	A0201	17	7
				A0201	23	172
				A26	21	N.A.
	52-61	KLNYEVMTKL	576	A24	N.A.	18
SSX4 98-124				A3	14	4
				B7	N.A.	4
	66-74	TLPPFMRSK	577	A26	16	N.A.
				A3	25	14
	110-118	KIMPKKPAE	578	A0201	15	<5
SSX4 98-124				A26	15	N.A.
				A3	16	<5
				A0201	15	8
	103-112	SLQRIFPKIM	579	A26	16	N.A.
				A3	15	<5

Table 36
Tyrosinase: Preferred Epitopes Revealed by Housekeeping Proteasome Digestion

Substrate	Epitope	Sequence	Seq. ID No.	Binding Prediction		
				HLA type	SYFPEITHI	NIH
Tyr 445-474	463-471	YKSYLEQA	580	A0201	18	<5
				A26	17	N.A.
	459-467	SFQDYKSY	581	A1	18	<5
				A26	22	N.A.
	458-467	DSFQDYKSY	582	A1	19	<5
Tyr 490-518				A26	24	N.A.
	507-514	IPEEKQPL	583	B08	28	5
				B5101	18	N.A.
				A0201	22	88
	506-514	QLPEEKQPL	584	A26	20	N.A.
				A24	N.A.	9
				B08	18	<5
	505-514	KQLPEEKQPL	585	A0201	15	28
				A24	N.A.	17
				A0201	15	<5
Tyr 490-518	507-515	LPEEKQPLL	586	B0702	21	24
				B08	28	5
				B5101	21	157
	506-515	QLPEEKQPLL	587	A0201	23	88
				A26	20	N.A.
	497-505	SLLCRHKRK	588	A24	N.A.	7
				A3	25	15

Example 15
Evaluating Likelihood of Epitope Cross-reactivity on Non-target Tissues.

As noted above PSA is a member of the kallikrein family of proteases, which is itself a subset of the serine protease family. While the members of this family sharing the greatest degree of sequence identity with PSA also share similar expression profiles, it remains possible that individual epitope sequences might be shared with proteins having distinctly different expression profiles. A first step in evaluating the likelihood of undesirable cross-reactivity is the identification of shared sequences. One way to accomplish this is to conduct a BLAST search of an epitope sequence against the SWISSPROT or Entrez non-redundant peptide sequence databases using the "Search for short nearly exact matches" option; hypertext transfer protocol accessible on the world wide web (<http://www.ncbi.nlm.nih.gov/blast/index.html>) at "ncbi.nlm.nih.gov/blast/index.html". Thus searching SEQ ID NO. 214, WVLTAAHCL, against SWISSPROT (limited to entries for homo sapiens) one finds four exact matches, including PSA. The other three are from kallikrein 1 (tissue kallikrein), and elastase 2A and 2B. While these nine amino acid segments are identical, the flanking sequences are quite distinct, particularly on the C-terminal side, suggesting that processing may proceed differently and that thus the same epitope may not be liberated from these other proteins. (Please note that kallikrein naming is confused. Thus the kallikrein 1 [accession number P06870] is a different protein than the one [accession number AAD13817] mentioned in the paragraph on PSA above in the section on tumor-associated antigens).

It is possible to test this possibility in several ways. Synthetic peptides containing the epitope sequence embedded in the context of each of these proteins can be subjected to *in vitro* proteasomal digestion and analysis as described above. Alternatively, cells expressing these other proteins, whether by natural or recombinant expression, can be used as targets in a cytotoxicity (or similar) assay using CD8⁺ T cells that recognize the epitope, in order to determine if the epitope is processed and presented.

Example 16
Epitope Clusters.

Known and predicted epitopes are generally not evenly distributed across the sequences of protein antigens. As referred to above, we have defined segments of sequence containing a higher than average density of (known or predicted) epitopes as epitope clusters. Among the uses of epitope clusters is the incorporation of their sequence into substrate peptides used in proteasomal digestion analysis as described herein. Epitope clusters can also be useful as vaccine components. A fuller discussion of the definition and uses of epitope clusters is found in U.S. Patent Application No. 09/561,571 entitled EPITOPE CLUSTERS, previously incorporated by reference.

The following tables (37-60) present 9-mer epitopes predicted for HLA-A2 binding using both the SYFPEITHI and NIH algorithms and the epitope density of regions of overlapping

epitopes, and of epitopes in the whole protein, and the ratio of these two densities. (The ratio must exceed one for there to be a cluster by the above definition; requiring higher values of this ratio reflect preferred embodiments). Individual 9-mers are ranked by score and identified by the position of their first amino in the complete protein sequence. Each potential cluster from a protein is numbered. The range of amino acid positions within the complete sequence that the cluster covers is indicated as are the rankings of the individual predicted epitopes it is made up of.

Table 37
BIMAS-NIH/Parker algorithm Results for gp100

Rank	Start	Score	Rank	Start	Score
1	619	1493	21	416	19
2	602	413	22	25	18
3	162	226	23	566	17
4	18	118	24	603	15
5	178	118	25	384	14
6	273	117	26	13	14
7	601	81	27	290	12
8	243	63	28	637	10
9	606	60	29	639	9
10	373	50	30	485	9
11	544	36	31	453	8
12	291	29	32	102	8
13	592	29	33	399	8
14	268	29	34	456	7
15	47	27	35	113	7
16	585	26	36	622	7
17	576	21	37	69	7
18	465	21	38	604	6
19	570	20	39	350	6
20	9	19	40	583	5

Table 38
SYFPEITHI (Rammensee algorithm) Results for gp100

Rank	Start	Score	Rank	Start	Score	Rank	Start	Score
1	606	30	37	291	20	73	60	18
2	162	29	38	269	20	74	17	18
3	456	28	39	2	20	75	613	17
4	18	28	40	610	19	76	599	17
5	602	27	41	594	19	77	572	17
6	598	27	42	591	19	78	557	17
7	601	26	43	583	19	79	556	17
8	597	26	44	570	19	80	512	17
9	13	26	45	488	19	81	406	17
10	585	25	46	446	19	82	324	17
11	449	25	47	322	19	83	290	17
12	4	25	48	267	19	84	101	17
13	603	24	49	250	19	85	95	17
14	576	24	50	205	19	86	635	16
15	453	24	51	180	19	87	588	16
16	178	24	52	169	19	88	584	16
17	171	24	53	88	19	89	577	16
18	11	24	54	47	19	90	559	16
19	619	23	55	10	19	91	539	16
20	280	23	56	648	18	92	494	16
21	268	23	57	605	18	93	482	16
22	592	22	58	604	18	94	468	16
23	544	22	59	595	18	95	442	16
24	465	22	60	571	18	96	413	16
25	399	22	61	569	18	97	408	16
26	373	22	62	450	18	98	402	16
27	273	22	63	409	18	99	286	16
28	243	22	64	400	18	100	234	16
29	566	21	65	371	18	101	217	16
30	563	21	66	343	18	102	211	16
31	485	21	67	298	18	103	176	16
32	384	21	68	209	18	104	107	16
33	350	21	69	102	18	105	96	16
34	9	21	70	97	18	106	80	16
35	463	20	71	76	18	107	16	16
36	397	20	72	69	18	108	14	16
						109	7	16

Table 39

Prediction of clusters for gp100

Total AAs: 661

Total 9-mers: 653

SYFPEITHI 16: 109 9-mers

NIH 5: 40 9-mers

			Epitopes/AA			
	Cluster #	AAs	Epitopes (by Rank)	Cluster	Whole Pr	Ratio
SYFPEITHI	1	2 to 26	39, 12, 109, 34, 55, 11, 9, 108, 107, 74, 4	0.440	0.165	2.668
	2	69-115	72, 71, 106, 53, 85, 105, 70, 84, 69, 104	0.213	0.165	1.290
	3	95-115	85, 105, 70, 84, 69	0.238	0.165	1.444
	4	162-188	2, 52, 17, 103, 16, 51	0.222	0.165	1.348
	5	205-225	50, 68, 102, 101	0.190	0.165	1.155
	6	243-258	28, 49	0.125	0.165	0.758
	7	267-306	48, 21, 38, 27, 20, 99, 83, 37, 67	0.225	0.165	1.364
	8	322-332	47, 82	0.182	0.165	1.103
	9	343-358	66, 33	0.125	0.165	0.758
	10	371-381	65, 26	0.182	0.165	1.103
	11	397-421	36, 25, 64, 98, 81, 97, 63, 96	0.320	0.165	1.941
	12	442-476	95, 46, 11, 62, 15, 3, 35, 24, 94	0.257	0.165	1.559
	13	482-502	93, 31, 45, 93	0.190	0.165	1.155
	14	539-552	91, 23	0.143	0.165	0.866
	15	556-627	79, 78, 90, 30, 29, 61, 44, 60, 77, 14, 89, 43, 88, 10, 87, 42, 22, 41, 59, 8, 6, 76, 7, 5, 13, 58, 57, 1, 40, 75, 19	0.431	0.165	2.611
NIH	1	9 to 33	20, 26, 4, 22	0.160	0.061	2.644
	2	268-281	14, 6	0.143	0.061	2.361
	3	290-299	27, 12	0.200	0.061	3.305
	4*	102-121	32, 35	0.100	0.061	1.653
	5*	373-392	10, 25	0.100	0.061	1.653
	6	453-473	31, 34, 18	0.143	0.061	2.361
	7	566-600	23, 19, 17, 40, 16, 13	0.171	0.061	2.833
	8	601-614	7, 2, 24, 38, 9	0.357	0.061	5.902
	9	619-630	1, 36	0.17	0.061	2.754
	10	637-647	28, 29	0.18	0.061	3.005

*Nearby but not overlapping epitopes

Table 40
BIMAS-NIH/Parker algorithm Results for PSMA

Rank	Start	Score
1	663	1360
2	711	1055
3	4	485
4	27	400
5	26	375
6	668	261
7	707	251
8	469	193
9	731	177
10	35	67
11	33	64
12	554	59
13	427	50
14	115	47
15	20	40
16	217	26
17	583	24
18	415	19
19	193	14
20	240	12
21	627	11
22	260	10
23	130	10
24	741	9
25	3	9
26	733	8
27	726	7
28	286	6
29	174	5
30	700	5

Table 41
SYFPEITHI (Rammensee algorithm) Results for PSMA

Rank	Start	Score	Rank	Start	Score	Rank	Start	Score
1	469	27	31	26	20	61	305	17
2	27	27	32	3	20	62	304	17
3	741	26	33	583	19	63	286	17
4	711	26	34	579	19	64	282	17
5	354	25	35	554	19	65	169	17
6	4	25	36	550	19	66	142	17
7	663	24	37	547	19	67	122	17
8	130	24	38	390	19	68	738	16
9	57	24	39	219	19	69	634	16
10	707	23	40	193	19	70	631	16
11	260	23	41	700	18	71	515	16
12	20	23	42	472	18	72	456	16
13	603	22	43	364	18	73	440	16
14	218	22	44	317	18	74	385	16
15	109	22	45	253	18	75	373	16
16	731	21	46	91	18	76	365	16
17	668	21	47	61	18	77	361	16
18	660	21	48	13	18	78	289	16
19	507	21	49	733	17	79	278	16
20	454	21	50	673	17	80	258	16
21	427	21	51	671	17	81	247	16
22	358	21	52	642	17	82	217	16
23	284	21	53	571	17	83	107	16
24	115	21	54	492	17	84	100	16
25	33	21	55	442	17	85	75	16
26	606	20	56	441	17	86	37	16
27	568	20	57	397	17	87	30	16
28	473	20	58	391	17	88	21	16
29	461	20	59	357	17			
30	200	20	60	344	17			

Table 42

Prediction of clusters for prostate-specific membrane antigen (PSMA)

Total AAs: 750

Total 9-mers: 742

SYFPEITHI 16: 88 9-mers

NIH 5: 30 9-mers

	Cluster #	Aas	Epitopes (by rank)	Epitopes/AA		
				Cluster	Whole Pr	Ratio
SYFPEITHI	1	3 to 12	32, 6	0.200	0.117	1.705
	2	13-45	13, 12, 88, 31, 2, 87, 25, 86	0.242	0.117	2.066
	3	57-69	9, 47	0.154	0.117	1.311
	4	100-138	84, 83, 15, 24, 67, 8	0.154	0.117	1.311
	5	193-208	40, 30	0.111	0.117	0.947
	6	217-227	82, 14, 39	0.273	0.117	2.324
	7	247-268	81, 45, 80, 11	0.182	0.117	1.550
	8	278-297	79, 64, 23, 63, 78	0.250	0.117	2.131
	9	354-381	5, 59, 22, 77, 43, 76, 75	0.250	0.117	2.131
	10	385-405	74, 38, 58, 57	0.190	0.117	1.623
	11	440-450	73, 56, 55	0.273	0.117	2.324
	12	454-481	20, 72, 29, 1, 42, 28	0.214	0.117	1.826
	13	507-523	17, 71	0.118	0.117	1.003
	14	547-562	37, 36, 35	0.188	0.117	1.598
	15	568-591	27, 53, 34, 33	0.167	0.117	1.420
	16	603-614	13, 26	0.167	0.117	1.420
	17	631-650	70, 69, 52	0.150	0.117	1.278
	18	660-681	18, 7, 17, 51, 50	0.227	0.117	1.937
	19	700-719	41, 10, 4	0.150	0.117	1.278
	20	731-749	16, 49, 68, 3	0.211	0.117	1.794
NIH	1	3 to 12	25, 3	0.200	0.040	5.000
	2	20-43	15, 5, 4, 11, 10	0.208	0.040	5.208
	3*	415-435	18, 13	0.095	0.040	2.381
	4	663-676	1, 6	0.143	0.040	3.571
	5	700-715	30, 7, 3	0.188	0.040	4.688
	6	726-749	27, 9, 26, 24	0.167	0.040	4.167

*Nearby but not overlapping epitopes

Table 43
BIMAS-NIH/Parker algorithm Results for PSA

Rank	Start	Score
1	7	607
2	170	243
3	52	124
4	53	112
5	195	101
6	165	23
7	72	18
8	245	18
9	2	16
10	59	16
11	122	15
12	125	15
13	191	13
14	9	8
15	14	6
16	175	5
17	130	5

Table 44
SYFPEITHI (Rammensee algorithm) Results for PSA

Rank	Start	Score
1	72	26
2	170	22
3	53	22
4	7	22
5	234	21
6	166	21
7	140	21
8	66	21
9	241	20
10	175	20
11	12	20
12	41	19
13	20	19
14	14	19
15	130	18
16	124	18
17	121	18
18	47	18
19	17	18
20	218	17
21	133	17
22	125	17
23	122	17
24	118	17
25	110	17
26	67	17
27	52	17
28	21	17
29	16	17
30	2	17
31	184	16
32	179	16
33	158	16
34	79	16
35	73	16
36	4	16

Table 45

Prediction of clusters for prostate specific antigen (PSA)

Total AAs: 261

Total 9-mers: 253

SYFPEITHI 16: 36 9-mers

NIH 5: 17 9-mers

	Cluster #	AAs	Epitopes (by rank)	Epitopes/AA		
				Cluster	Whole Pr	Ratio
SYFPEITHI	1	2 to 29	30, 36, 4, 11, 14, 29, 19, 13, 28	0.321	0.138	2.330
	2	41-61	12, 18, 27, 3	0.190	0.138	1.381
	3	66-87	8, 26, 1, 35, 34	0.227	0.138	1.648
	4	110-148	25, 24, 17, 23, 16, 22, 15, 21, 7	0.184	0.138	1.332
	5	158-192	33, 6, 2, 10, 32, 31	0.171	0.138	1.243
	6	234-249	5, 9	0.125	0.138	0.906
	7*	118-133	24, 17, 23, 16, 22	0.313	0.138	2.266
	8*	118-138	24, 17, 23, 16, 22, 15	0.286	0.138	2.071
NIH	1	2-22	9, 1, 14, 15	0.190	0.065	2.924
	2	52-67	3, 4, 10	0.188	0.065	2.879
	3	122-138	11, 12, 17	0.176	0.065	2.709
	4	165-183	6, 2, 16	0.158	0.065	2.424
	5	191-203	13, 5	0.154	0.065	2.362
	6**	52-80	3, 4, 10, 7	0.138	0.065	2.118

*These clusters are internal to the less preferred cluster #4.

**Includes a nearby but not overlapping epitope.

Table 46
BIMAS-NIH/Parker algorithm Results for PSCA

Rank	Start	Score
1	43	53
2	5	84
3	7	79
4	109	36
5	105	25
6	108	24
7	14	21
8	20	18
9	115	17
10	42	15
11	36	15
12	99	9
13	58	8

20

Table 47
SYFPEITHI (Rammensee algorithm) Results for PSCA

25

Rank	Start	Score	Rank	Start	Score
1	108	30	17	54	19
2	14	30	18	12	19
3	105	29	19	4	19
4	5	28	20	1	19
5	115	26	21	112	18
6	99	26	22	101	18
7	7	26	23	98	18
8	109	24	24	51	18
9	53	23	25	43	18
10	107	21	26	106	17
11	20	21	27	104	17
12	8	21	28	83	17
13	13	20	29	63	17
14	102	19	30	50	17
15	60	19	31	3	17
16	57	19	32	9	16
			33	92	16

Table 48**Prediction of clusters for prostate stem cell antigen (PSCA)**

Total AAs: 123

Total 9-mers: 115

SYFPEITHI 16: 33;

SYFPEITHI 20: 13

NIH 5: 13

	Cluster #	AAs	Epitopes (by rank)	Epitopes/AA		
				Cluster	Whole Pr.	Ratio
SYFPEITHI >16	1	1 to 28	20, 31, 19, 4, 7, 12, 33, 18, 13, 2, 11	0.393	0.268	1.464
	2	43-71	25, 30, 24, 9, 17, 16, 15, 29	0.276	0.268	1.028
	3	92-123	32, 23, 6, 27, 14, 22, 3, 26, 10, 1, 8, 21, 5	0.406	0.268	1.514
SYFPEITHI >20	1	5 to 28	4, 7, 12, 13, 2, 11	0.250	0.106	2.365
	2	99-123	6, 3, 10, 1, 8, 5	0.240	0.106	2.271
NIH	1	5 to 28	2, 3, 7, 8	0.167	0.106	1.577
	2	36-51	11, 10, 1	0.188	0.106	1.774
	3	99-123	12, 5, 6, 4, 9	0.200	0.106	1.892
	4*	105-116	5, 6, 4	0.250	0.106	2.365

*This cluster is internal to the less preferred cluster #3.

In tables 49-60 epitope prediction and cluster analysis data for each algorithm are presented together in a single table.

Table 49

Prediction of clusters for MAGE-1 (NIH algorithm)

Total AAs: 309
Total 9-mers: 301
NIH 5:19 9-mers

Cluster #	AAs	Epitope Rank	Start Position	NIH Score	Cluster	Epitopes/AA Whole Pr.	Ratio
1	18-32	16	18	9	0.133	0.063	2.112
		19	24	7			
2	101-113	14	101	11	0.154	0.063	2.442
		7	105	44			
3	146-159	9	146	32	0.143	0.063	2.263
		3	151	169			
4	169-202	10	169	32	0.176	0.063	2.796
		13	174	16			
		18	181	8			
		17	187	8			
		6	188	74			
		5	194	110			
5	264-277	2	264	190	0.143	0.063	2.263
		12	269	20			
6	278-290	1	278	743	0.154	0.063	2.437
		11	282	28			

Table 50

Prediction of clusters for MAGE-1 (SYFPEITHI algorithm)

Total AAs: 309

Total 9-mers: 301

SYFPEITHI 16: 46 9-mers

Cluster #	Aas	Epitope Rank	Start Position	SYFPEITHI Score	Epitopes/AA		
					Cluster	Whole	Ratio
1	7-49	22	7	19	0.233	0.153	1.522
		9	15	22			
		27	18	18			
		16	20	20			
		28	22	18			
		29	24	18			
		33	31	17			
		30	35	18			
		2	38	26			
		17	41	20			
2	89-132	10	89	22	0.273	0.153	1.783
		18	92	20			
		7	93	23			
		23	96	19			
		43	98	16			
		4	101	25			
		8	105	23			
		34	107	17			
		35	108	17			
		36	113	17			
		37	118	17			
		19	124	20			
3	167-203	44	167	16	0.270	0.153	1.766
		20	169	20			
		12	174	21			
		24	181	19			
		6	187	24			
		31	188	18			
		25	191	19			
		38	192	17			
		1	194	27			
		13	195	21			
4	230-246	14	230	21	0.118	0.153	0.769
		39	238	17			
5	264-297	15	264	21	0.235	0.153	1.538
		32	269	18			
		40	270	17			
		26	271	19			
		46	275	16			
		3	278	26			
		21	282	20			
		41	289	17			

Table 51

Prediction of clusters for MAGE-2 (NIH algorithm)

Total AAs: 314

Total 9-mers: 308

NIH \geq 5: 20 9-mers

Cluster #	AAs	Epitope Rank	Start Position	NIH Score	Cluster	Epitope/AA Whole Pr.	Ratio
1	101-120	18	101	5.373	0.150	0.065	2.310
		16	108	6.756			
		1	112	2800.697			
2	153-167	8	153	31.883	0.200	0.065	3.080
		4	158	168.552			
		7	159	32.138			
3	169-211	14	169	8.535	0.209	0.065	3.223
		19	174	5.346			
		6	176	49.993			
		11	181	15.701			
		15	188	7.536			
		12	195	12.809			
		5	200	88.783			
		10	201	16.725			
4	271-284	3	271	398.324	0.143	0.065	2.200
		9	276	19.658			

Table 52

Prediction of clusters for MAGE-2 (SYFPEITHI algorithm)

Total AAs: 314

Total 9-mers: 308

SYFPEITHI 16: 52 9-mers

Cluster #	AAs	Enitone Rank	Start Position	SYFPEITHI Score	Cluster	Enitones/AA Whole Pr.	Ratio
1	15-32	13	15	21	0.278	0.169	1.645
		29	18	18			
		43	20	16			
		30	22	18			
		21	24	19			
2	37-56	31	37	18	0.250	0.169	1.481
		16	40	20			
		44	44	16			
		14	45	21			
		22	48	19			
3	96-133	36	96	17	0.211	0.169	1.247
		46	101	16			
		6	108	25			
		47	109	16			
		2	112	27			
		37	120	17			
		38	125	17			
		17	131	20			
4	153-216	12	153	22	0.344	0.169	2.036
		39	158	17			
		7	159	25			
		23	161	19			
		24	162	19			
		48	164	16			
		49	167	16			
		32	170	18			
		50	171	16			
		4	174	26			
		9	176	24			
		51	177	16			
		15	181	21			
		25	188	19			
		18	194	20			
		33	195	18			
		19	198	20			
		3	200	27			
		1	201	28			
		40	202	17			
		10	203	23			
		52	208	16			
5	237-254	26	237	19	0.167	0.169	0.987
		27	245	19			
		34	246	18			
6	271-299	8	271	25	0.241	0.169	1.430
		35	276	18			
		41	277	17			
		11	278	23			
		28	283	19			
		20	285	20			
		42	291	17			

Table 53

Prediction of clusters for MAGE-3 (NIH algorithm)

Total AAs: 314

Total 9-mers: 308

NIH 5: 22 9-mers

Cluster #	AAs	Epitope Rank	Start Position	NIH Score	Cluster	Epitopes/AA Whole Pr.	Ratio
1	101-120	15	101	11.002	0.200	0.071	2.800
		21	105	6.488			
		8	108	49.134			
		2	112	339.313			
2	153-167	18	153	7.776	0.200	0.071	2.800
		6	158	51.77			
		22	159	5.599			
3	174-209	17	174	8.832	0.194	0.071	2.722
		7	176	49.993			
		13	181	15.701			
		19	188	7.536			
		14	195	12.809			
		5	200	88.783			
		12	201	16.725			
4	237-251	16	237	10.868	0.200	0.071	2.800
		4	238	148.896			
		20	243	6.88			
5	271-284	1	271	2655.495	0.143	0.071	2.000
		11	276	19.658			

Table 54

Prediction of clusters for MAGE-3 (SYFPEITHI algorithm)

Total AAs: 314

Total 9-mers: 308

SYFPEITHI 16: 47 9-mers

Cluster #	AAs	Epitope Rank	Start Position	SYFPEITHI Score	Cluster	Epitones/AA Whole Pr.	Ratio
1	15-32	12	15	21	0.278	0.153	1.820
		26	18	18			
		37	20	16			
		27	22	18			
		18	24	19			
2	38-56	38	38	16	0.263	0.153	1.725
		15	40	20			
		39	44	16			
		13	45	21			
		19	48	19			
3	101-142	28	101	18	0.190	0.153	1.248
		40	105	16			
		1	108	31			
		6	112	25			
		31	120	17			
		32	125	17			
		16	131	20			
		41	134	16			
4	153-216	20	153	19	0.313	0.153	2.048
		29	156	18			
		33	158	17			
		21	159	19			
		34	161	17			
		42	164	16			
		43	167	16			
		10	174	22			
		8	176	23			
		14	181	21			
		22	188	19			
		44	193	16			
		11	194	22			
		23	195	19			
		45	197	16			
		17	198	20			
		3	200	27			
		2	201	28			
		35	202	17			
		46	208	16			
5	220-230	5	220	26	0.182	0.153	1.191
		47	222	16			
6	237-246	7	237	25	0.200	0.153	1.311
		9	238	23			
7	271-293	4	271	27	0.217	0.153	1.425
		30	276	18			
		24	278	19			
		36	283	17			
		25	285	19			

Table 55

Prediction of clusters for PRAME (NIH algorithm)

Total AAs: 509

Total 9-mers: 501

NIH 5: 40 9-mers

Cluster #	AAs	Epitope Rank	Start Position	NIH Score	Cluster	Epitopes/AA Whole Pr.	Ratio
1	33-47	20	33	18	0.133	0.080	1.670
		17	39	21			
2	71-81	9	71	50	0.2	0.07984	2.505
		32	73	7			
3	99-108	23	100	15	0.2	0.07984	2.505
		24	99	13			
4	126-135	38	126	5	0.2	0.07984	2.505
		35	127	6			
5	224-246	5	224	124	0.130	0.080	1.634
		8	230	63			
		39	238	5			
6	290-303	18	290	18	0.214	0.080	2.684
		14	292	23			
		7	295	66			
7	305-324	28	305	10	0.200	0.080	2.505
		30	308	8			
		25	312	13			
		36	316	6			
8	394-409	2	394	182	0.188	0.080	2.348
		12	397	42			
		31	401	7			
9	422-443	10	422	49	0.227	0.080	2.847
		3	425	182			
		34	431	7			
		29	432	9			
		4	435	160			
10	459-487	15	459	21	0.172	0.080	2.159
		11	462	45			
		22	466	15			
		40	472	5			
		37	479	6			

Table 56

Prediction of clusters for PRAME (SYFPEITHI algorithm)

Total AAs: 509

Total 9-mers: 501

SYFPEITHI 17: 80 9-mers

Cluster #	AAs	Epitope Rank	Start Position	SYFPEITHI Score	Cluster	Epitopes/AA Whole Pr.	Ratio
1	18-59	65	18	17	0.238	0.160	1.491
		50	21	18			
		66	26	17			
		35	33	20			
		22	34	22			
		51	37	18			
		5	39	27			
		23	40	22			
		13	44	24			
		46	51	19			
2	78-115	36	78	20	0.263	0.160	1.648
		67	80	17			
		52	84	18			
		24	86	22			
		53	91	18			
		25	93	22			
		9	99	25			
		8	100	26			
		54	103	18			
		55	107	18			
3	191-202	56	191	18	0.167	0.160	1.044
		38	194	20			
4	205-215	26	205	22	0.182	0.160	1.139
		27	207	22			
5	222-238	47	222	19	0.235	0.160	1.474
		14	224	24			
		69	227	17			
		57	230	18			
6	241-273	70	241	17	0.212	0.160	1.328
		15	248	24			
		71	255	17			
		30	258	21			
		39	259	20			
		58	261	18			
		40	265	20			
7	290-342	72	290	17	0.208	0.160	1.300
		48	293	19			
		31	298	21			
		73	301	17			
		18	305	23			
		6	308	27			
		10	312	25			
		19	316	23			
		28	319	22			

Prediction of clusters for PRAME (SYFPEITHI algorithm)

Total AAs: 509

Total 9-mers: 501

SYFPEITHI 17: 80 9-mers

Cluster #	AAs	Epitope Rank	Start Position	SYFPEITHI Score	Cluster	Epitopes/AA Whole Pr.	Ratio
		41	326	20			
		74	334	17			
8	343-363	59	343	18	0.238	0.160	1.491
		60	348	18			
		75	351	17			
		20	353	23			
		76	355	17			
9	364-447	49	364	19	0.250	0.160	1.566
		32	371	21			
		11	372	25			
		61	375	18			
		77	382	17			
		21	390	23			
		78	391	17			
		1	394	30			
		42	397	20			
		62	403	18			
		33	410	21			
		43	418	20			
		34	419	21			
		7	422	27			
		2	425	29			
		79	426	17			
		63	428	18			
		64	431	18			
		12	432	25			
		16	435	24			
		80	439	17			
10	455-474	29	455	22	0.200	0.160	1.253
		17	459	24			
		4	462	28			
		3	466	29			

Table 57
Predication of clusters for CEA (NIH algorithm)

Total AAs:702

Total 9-mers: 694

NIH 5: 30 9-mers

Cluster #	AA	Peptides Rank	Start Position	Score	Cluster	Peptides/AAs Whole Pr.	Ratio
1	17-32	5	17	79.041	0.188	0.043	4.388
		7	18	46.873			
		20	24	12.668			
2	113-129	2	113	167.991	0.118	0.043	2.753
		15	121	21.362			
3	172-187	25	172	9.165	0.125	0.043	2.925
		14	179	27.995			
4	278-291	30	278	5.818	0.143	0.043	3.343
		17	283	19.301			
5	350-365	9	350	43.075	0.125	0.043	2.925
		12	357	27.995			
6	528-543	8	528	43.075	0.125	0.043	2.925
		13	535	27.995			
7	631-645	23	631	9.563	0.200	0.043	4.680
		19	634	13.381			
		24	637	9.245			
8	691-702	1	691	196.407	0.167	0.043	3.900
		27	694	7.769			

Table 58
Predication of clusters for CEA (SYFPEITHI algorithm)

Total AAs: 702

Total 9-mers: 694

SYFPEITHI 16: 81 9-mers

Cluster #	AA	Peptides Rank	Start Position	Score	Cluster	Peptides/AAs Whole Pr.	Ratio
1	5-36	67	5	16	0.250	0.117	2.140
		23	12	19			
		24	16	19			
		9	17	22			
		25	18	19			
		32	19	18			
		68	23	16			
		33	28	18			
2	37-62	41	37	17	0.269	0.117	2.305
		20	44	20			
		26	45	19			
		42	46	17			
		27	50	19			
		43	53	17			
		44	54	17			
3	99-115	14	99	21	0.235	0.117	2.014
		5	100	23			
		45	104	17			
		34	107	18			
4	116-129	69	116	16	0.143	0.117	1.223
		21	121	20			
5	172-187	46	172	17	0.125	0.117	1.070
		70	179	16			
6	192-202	3	192	24	0.182	0.117	1.557
		47	194	17			
7	226-241	48	226	17	0.188	0.117	1.605
		49	229	17			
		15	233	21			
8	307-318	11	307	22	0.250	0.117	2.140
		71	308	16			
		51	310	17			
9	319-349	52	319	17	0.129	0.117	1.105
		53	327	17			
		72	335	16			
		35	341	18			
10	370-388	12	370	22	0.211	0.117	1.802
		54	372	17			
		74	375	16			
		6	380	23			
11	403-419	56	403	17	0.235	0.117	2.014
		57	404	17			
		58	407	17			
		28	411	19			

12	427-442	59	427	17	0.188	0.117	1.605
		75	432	16			
		76	434	16			
13	450-462	77	450	16	0.154	0.117	1.317
		13	454	22			
14	488-505	36	488	18	0.167	0.117	1.427
		18	492	21			
		60	497	17			
15	548-558	4	548	24	0.182	0.117	1.557
		61	550	17			
16	565-577	62	565	17	0.154	0.117	1.317
		19	569	21			
17	579-597	78	579	16	0.143	0.117	1.223
		79	582	16			
		7	589	23			
18	605-618	2	605	25	0.143	0.117	1.223
		38	610	18			
19	631-669	29	631	19	0.154	0.117	1.317
		63	637	17			
		80	644	16			
		64	652	17			
		39	660	18			
		81	661	16			
20	675-702	22	675	20	0.286	0.117	2.446
		30	683	19			
		31	687	19			
		40	688	18			
		65	690	17			
		1	691	31			
		66	692	17			
		8	694	23			

Table 59
Predication of clusters for SCP-1 (NIH algorithm)

Total AAs: 976
 Total 9-mers: 968
 NIH 5: 37 9-mers

Cluster #	AA	Peptides Rank	Start Position	Score	Cluster	Peptides/AAs Whole Pr.	Ratio
1	101-116	15	101	40.589	0.125	0.038	3.270
		13	108	57.255			
2*	281-305	14	281	44.944	0.12	0.038	3.139
		24	288	15.203			
		17	297	32.857			
3	431-447	8	431	80.217	0.073	0.038	1.914
		26	438	11.861			
		4	439	148.896			
4	557-579	11	557	64.335	0.174	0.038	4.550
		19	560	24.937			
		6	564	87.586			
		18	571	32.765			
5	635-650	10	635	69.552	0.125	0.038	3.270
		34	642	6.542			
6	755-767	36	755	5.599	0.154	0.038	4.025
		35	759	5.928			
7	838-854	2	838	284.517	0.118	0.038	3.078
		28	846	11.426			

Table 60
Predication of clusters for SCP-1

Total AAs: 976

Total 9-mers: 968

Rammensee 16: 118 9-mers

Cluster #	AA	Peptides Rank	Start Position	Score	Cluster	Peptides/AAs Whole Pr.	Ratio
1	8-28	99	8	16	0.143	0.121	1.182
		77	15	17			
		100	20	16			
2	63-80	78	63	17	0.222	0.121	1.838
		50	66	19			
		102	69	16			
		60	72	18			
3	94-123	79	94	17	0.133	0.121	1.103
		12	101	23			
		17	108	22			
		103	115	16			
4	126-158	35	126	20	0.182	0.121	1.504
		36	133	20			
		51	139	19			
		80	140	17			
		61	143	18			
		37	150	20			
5	161-189	38	161	20	0.207	0.121	1.711
		52	165	19			
		81	171	17			
		82	177	17			
		62	178	18			
		39	181	20			
6	213-230	40	213	20	0.167	0.121	1.379
		13	220	23			
		28	222	21			
7	235-250	63	235	18	0.125	0.121	1.034
		18	242	22			
8	260-296	83	260	17	0.243	0.121	2.012
		105	262	16			
		84	267	17			
		106	269	16			
		41	270	20			
		64	271	18			
		85	274	17			
		19	281	22			
		3	288	25			
9	312-338	108	312	16	0.148	0.121	1.225
		29	319	21			
		30	323	21			
		65	330	18			
10	339-355	66	339	18	0.235	0.121	1.946
		31	340	21			
		42	344	20			
		53	347	19			

11	376-447	54	376	19	0.194	0.121	1.608
		43	382	20			
		44	386	20			
		20	390	22			
		55	397	19			
		6	404	24			
		86	407	17			
		45	411	20			
		67	417	18			
		21	425	22			
		46	431	20			
		68	432	18			
		32	438	21			
		7	439	24			
12	455-488	33	455	21	0.235	0.121	1.946
		47	459	20			
		56	462	19			
		87	463	17			
		88	466	17			
		14	470	23			
		109	473	16			
		34	480	21			
13	515-530	57	515	19	0.125	0.121	1.034
		22	522	22			
14	557-590	8	557	24	0.147	0.121	1.216
		23	564	22			
		9	571	24			
		90	575	17			
		58	582	19			
15	610-625	69	610	18	0.125	0.121	1.034
		91	617	17			
16	633-668	92	633	17	0.222		
		10	635	24			
		70	638	18			
		93	640	17			
		48	642	20			
		49	645	20			
		111	652	16			
		112	660	16			
17	674-685	71	674	18	0.167	0.121	1.379
		11	677	24			
18	687-702	1	687	26	0.125	0.121	1.034
		94	694	17			
19	744-767	113	744	16	0.250	0.121	2.068
		95	745	17			
		4	745	25			
		24	752	22			
		2	755	26			
		72	759	18			
20	812-827	97	812	17	0.125	0.121	1.034
		115	819	16			
21	838-857	116	838	16	0.150	0.121	1.241
		25	846	22			
		74	849	18			

22	896-913	117	896	16	0.222	0.121	1.838
		98	899	17			
		26	902	22			
		76	905	18			

1 MLLAVLYCLL WSFQTSAGHF PRACVSSKNL MEKECCPPWS GDRSPCGQLS
 GRGSCQNILL
 5 61 SNAPLGPOFP FTGVDDRESW PSVFYNRTCQ CSGNFMGFNC GNCKFGFWGP
 NCTERRLLVR
 121 RNIFDLSAPE KDKFFAYLTL AKHTISSDYV IPIGTYQMK NGSTPMFNDI
 NIYDLFVWMH
 181 YYVSM DALLG GSEIWRDIDF AHEAPFLPW HRLFLLRWEQ EIQKLTGDEN
 10 FTIPYDWRD
 241 AEKCDICTDE YMGGQHPTNP NLLSPASFFS SWQIVCSRLE EYN SHQSLCN
 GTPEGPLRRN
 301 PGNHDKSRTP RLPSSADVEF CLSLTQYESG SMDKAANFSF RNTLEGFASP
 LTGIADASQS
 15 361 SMHNALHIYM NGTMSQVQGS ANDPIFLLHH AFVDSIFEQW LRRHRPLQEV
 YPEANAPIGH
 421 NRESYMVPFI PLYRNGDFFI SSKDLGYDYS YLQSDPDPSF QDYIKSYLEQ
 ASRIWSWLLG
 481 AAMVGAVLTA LLAGLVSLLC RHKRKQLPEE KQPLLMEKED YHSLYQSHL
 20 TYROSINASE PROTEIN

25

30 1 MNGDDAFARR PTVGAQIPEK IQKAFDDIAK YFSKEEWEKM KASEKIFYVY
 MKRKYEAMTK
 61 LGFKATLPPF MCNKRAEDFQ GNDLDNDPNR GNQVERPQMT FGR LQGISPK
 IMPKKPAEEG
 121 NDSEEVPEAS GPQNDGKELC PPGKPTTSEK IHERSGPKRG EHAWTHRLRE
 35 RKQLVIYEEI
 181 SDPEEDDE
 SSX-2 PROTEIN

40

1 MWNLLHETDS AVATARRPRW LCAGALVLAG GFFLLGFLFG WFIKSSNEAT
 NITPKHNMKA
 61 FLDELKAENI KKFLYNFTQI PHLAGTEQNF QLAKQIQSQW KEFGLDSEVL
 5 AHYDVLLSYP
 121 NKTHPNYISI INEDGNEIFN TSLFEPPLPG YENVSDIVPP FSAFSPQGMP
 EGDLYVNYA
 181 RTEDFFKLER DMKINCSGKI VIARYGKVR GNKVKNAQLA GAKGVILYSD
 PADYFAPGVK
 10 241 SYPDGWNLP GGVQRGNILN LNGAGDPLTP GYPANEYAYR RGIAEAVGLP
 SIPVHPIGYY
 301 DAQKLEKMG GSAPPDSSWR GSKVPYNVG PGFTGNFSTQ KVKMHIHSTN
 EVTRIYNVIG
 361 TLRGAVEPDR YVILGGHRDS WVFGGIDPQS GAAVVHEIVR SFGTLKKEGW
 15 RPRRTILFAS
 421 WDAEEFGLLG STEWAEENSR LLQERGVAYI NADSSIEGNY TLRVDCTPLM
 YSLVHNLTK
 481 LKSPDEGFEG KSLYESWTKK SPSPEFSGMP RISKLGSGND FEVFFQRLGI
 ASGRARYTKN
 20 541 WETNKFSGYP LYHSVYETYE LVEKFYDPMF KYHLTVAQVR GGMVFELANS
 IVLPFDCRDY
 601 AVVLRKYADK IYSISMKHPQ EMKTYSVSFD SLFSAVKNFT EIASKFSERL
 QDFDKSNPIV
 661 LRMMNDQLMF LERAFIDPLG LPDRPFYRHV IYAPSSHNKY AGESFPGIYD
 25 ALFDIESKVD
 721 PSKAWGEVKR QIYVAAFTVQ AAAETLSEVA

PSMA PROTEIN

30

1: Homo sapiens tyrosinase
 NM_00037 (oculocutaneous albinism IA) (TYR), PubMed, Protein,
 2 Mrna Taxonomy, OMIM, LinkOut
 LOCUS NM_000372 1964 bp mRNA PRI 31-
 OCT-2000
 DEFINITION Homo sapiens tyrosinase (oculocutaneous albinism IA)
 35 (TYR), mRNA.
 ACCESSION NM_000372
 VERSION NM_000372.1 GI:4507752
 KEYWORDS
 SOURCE human.
 40 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
 Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
 Homo.
 45 REFERENCE 1 (bases 1 to 1964)
 AUTHORS Kwon BS, Haq AK, Pomerantz SH and Halaban R.
 TITLE Isolation and sequence of a cDNA clone for human
 tyrosinase that
 maps at the mouse c-albino locus
 50 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 84 (21), 7473-7477
 (1987)
 MEDLINE 88041128
 PUBMED 2823263
 REMARK Erratum: [published erratum appears in Proc Natl Acad
 55 Sci U S A

- 1988 Sep;85(17):6352]]
 REFERENCE 2 (bases 1 to 1964)
 AUTHORS Barton DE, Kwon BS and Francke U.
 TITLE Human tyrosinase gene, mapped to chromosome 11 (q14---
 5 -q21),
 defines second region of homology with mouse
 chromosome 7
 JOURNAL Genomics 3 (1), 17-24 (1988)
 MEDLINE 89122007
 10 PUBMED 3146546
 REFERENCE 3 (bases 181 to 1964)
 AUTHORS Shibahara,S., Tomita,Y., Tagami,H., Muller,R.M. and
 Cohen,T.
 TITLE Molecular basis for the heterogeneity of human
 15 tyrosinase
 JOURNAL Tohoku J. Exp. Med. 156 (4), 403-414 (1988)
 MEDLINE 89222868
 REFERENCE 4 (bases 1 to 1964)
 20 AUTHORS Bouchard B, Fuller BB, Vijayasaradhi S and Houghton
 AN.
 TITLE Induction of pigmentation in mouse fibroblasts by
 expression of
 human tyrosinase cDNA
 JOURNAL J. Exp. Med. 169 (6), 2029-2042 (1989)
 25 MEDLINE 89279151
 PUBMED 2499655
 REFERENCE 5 (bases 1 to 1964)
 AUTHORS Takeda,A., Tomita,Y., Okinaga,S., Tagami,H. and
 Shibahara,S.
 30 TITLE Functional analysis of the cDNA encoding human
 tyrosinase precursor
 JOURNAL Biochem. Biophys. Res. Commun. 162 (3), 984-990 (1989)
 MEDLINE 89351001
 REFERENCE 6 (bases 1 to 1964)
 35 AUTHORS Kikuchi H, Miura H, Yamamoto H, Takeuchi T, Dei T and
 Watanabe M.
 TITLE Characteristic sequences in the upstream region of the
 human
 tyrosinase gene
 40 JOURNAL Biochim. Biophys. Acta 1009 (3), 283-286 (1989)
 MEDLINE 90089403
 PUBMED 2480811
 REFERENCE 7 (bases 1 to 1964)
 45 AUTHORS Giebel LB, Strunk KM and Spritz RA.
 TITLE Organization and nucleotide sequences of the human
 tyrosinase gene
 and a truncated tyrosinase-related segment
 JOURNAL Genomics 9 (3), 435-445 (1991)
 50 MEDLINE 91236163
 PUBMED 1903356
 REFERENCE 8 (bases 1 to 1964)
 AUTHORS Brichard V, Van Pel A, Wolfel T, Wolfel C, De Plaen E,
 Lethe B,
 Coulie P and Boon T.
 55 TITLE The tyrosinase gene codes for an antigen recognized by
 autologous
 cytolytic T lymphocytes on HLA-A2 melanomas
 JOURNAL J. Exp. Med. 178 (2), 489-495 (1993)

MEDLINE 93340625
 PUBMED 8340755
 COMMENT PROVISIONAL REFSEQ: This record has not yet been
 subject to final
 5 NCBI review. The reference sequence was derived from
M27160.1.
FEATURES Location/Qualifiers
 source 1..1964
 /organism="Homo sapiens"
 10 /db_xref="taxon:9606"
 /chromosome="11"
 /map="11q14-q21"
 /cell_line="MeWo melanoma DNA, and cDNA to
 mRNA"
 15 /tissue_type="placenta"
 gene 1..1964
 /gene="TYR"
 /note="OCA1A"
 20 /db_xref="LocusID:7299"
 /db_xref="MIM:203100"
 CDS 83..1672
 /gene="TYR"
 /EC_number="1.14.18.1"
 25 /note="Tyrosinase"
 /codon_start=1
 /db_xref="LocusID:7299"
 /db_xref="MIM:203100"
 /product="tyrosinase (oculocutaneous albinism
 IA)"
 30 /protein_id="NP_000363.1"
 /db_xref="GI:4507753"
 /translation="MLLAVLYCLLWSFQTSAGHFPRACVSSKNLMEKECCPPWSGDRS
 35 PCGQLSGRGSCQNILLSNAPLGPQFPFTGVDDRESWPSVFYNRTCQCSGNFMGFNCGN
 CKFGFWGPNCTERRLLVRRNIFDLSAPEKDKFFAYLTLAKHTISSDYVIPIGTYGQMK
 NGSTPMFNDINIYDLFVWMHYVVSMDALLGGSEIWRDIDFAHEAPAFLPWHRLFLLRW
 40 EQEIQKLTGDENFTIPYWDWRDAEKCDICTDEYMGQHPNTNPALLSPASFFSSWQIVC
 SRLEEYNHQSLCNGTPEGPLRRNPGNHDKSRTPRLPSSADVEFCLSLTQYESGSMDK
 45 AANFSFRNTLEGFASPLTGIADASQSSMHNALHIYMNGTMSQVQGSANDPIFLLHHAF
 VDSIFEQWLRRHRPLQEVYPEANAPIGHNRESYMPFPIPLYRNGDFFISSKDLGYDYS
 YLQSDPDPSFQDIKSYLEQASRIWSWLLGAAMVGAVLTALLAGLVSLLCRHKRKQLP
 50 EEKQPLLMEKEDYHSLYQSHL"
 sig_peptide 83..136
 mat_peptide 137..1669
 /note="tyrosinase"
 /product="tyrosinase (oculocutaneous albinism
 55 IA)"
 misc_feature 560..1429
 /note="tyrosinase; Region: Common central
 domain of

tyrosinase"

	BASE COUNT	520 a	462 c	445 g	537 t
	ORIGIN				
	1	atcactgtag	tagtagctgg	aaagagaaat	ctgtgactcc aattagccag
5		ttcctgcaga			
	61	cottgtgagg	actagaggaa	gaatgctcct	ggctgttttg tactgcctgc
		tgtggagttt			
	121	ccagacctcc	gctggccatt	tccctagagc	ctgtgtctcc totaagaacc
		tgatggagaa			
10		181	ggaatgctgt	ccaccgtgga	gcggggacag gagtccctgt ggccagcttt
		caggcagagg			
	241	ttcctgtcag	aatatccttc	tgtccaatgc	accacttggg cctcaatttc
		ccttcacagg			
	301	ggtggatgac	cgggagtcgt	ggccttcctg	cttttataat aggacctgcc
15		agtgtctctgg			
	361	caacttcacg	ggattcaact	gtggaaactg	caagtttggc ttttggggac
		caaactgcac			
	421	agagagacga	ctcttggtga	gaagaaacat	cttcgatttg agtgccccag
		agaaggacaa			
20		481	atTTTTTg	tacctactt	tagcaaagca taccatcagc tcagactatg
		tcacccccat			
	541	agggacctat	ggccaaatga	aaaatggatc	aacacccatg ttttaacgaca
		tcaatattta			
	601	tgacctcttt	gtctggatgc	attattatgt	gtcaatggat gcaactgcttg
25		ggggatctga			
	661	aatctggaga	gacattgatt	ttgccccatga	agcaccagct tttctgcctt
		ggcatagact			
	721	cttcttggtg	cgggtgggaa	aagaaatcca	gaagctgaca ggagatgaaa
		acttcactat			
30		781	tccatattgg	gactggcg	atgcagaaaa gtgtgacatt tgcacagatg
		agtacatggg			
	841	aggtcagcac	cccacaaatc	ctaacttact	cagcccagca tcattctctt
		cctcttgcca			
	901	gattgtctgt	agccgattgg	aggagtacaa	cagccatcag tctttatgca
35		atggaacgcc			
	961	cgagggacct	ttacggcgta	atcctggaaa	ccatgacaaa tccagaaccc
		caaggctccc			
	1021	ctcttcagct	gatgtagaat	tttgccctgag	tttgacccaa tatgaatctg
		gttccatgga			
40		1081	taaagctgcc	aatttcagct	ttagaaatac actggaagga tttgctagtc
		cacttactgg			
	1141	gatagcggat	gcctctcaaa	gcagcatgca	caatgccttg cacatctata
		tgaatggaac			
	1201	aatgtccag	gtacagggat	ctgccaacga	tcctatcttc cttcttcacc
45		atgcatttgt			
	1261	tgacagtatt	tttgagcagt	ggctccgaag	gcaccgtcct cttcaagaag
		tttatccaga			
	1321	agccaatgca	cccattggac	ataaccggga	atcctacatg gttcctttta
		taccactgta			
50		1381	cagaaatgg	gatttcttta	tttcatccaa agatctgggc tatgactata
		gctatctaca			
	1441	agattcagac	ccagactctt	ttcaagacta	cattaagtcc tatttggaac
		aagcagagtcg			
	1501	gatctggtea	tggctccttg	gggcggcgat	ggtagggggc gtcctcactg
55		ccctgctggc			
	1561	agggcttgtg	agcttgctgt	gtcgtcacia	gagaaagcag cttcctgaag
		aaaagcagcc			

```

1621 actcctcatg gagaaagagg attaccacag cttgtatcag agccatttat
aaaaggctta
1681 ggcaatagag tagggccaaa aagcctgacc tcaacttaac tcaaagtaat
gtccagggttc
5 1741 ccagagaata tctgctggta tttttctgta aagaccattt gcaaaattgt
aacctaatac
1801 aaagtgtagc cttcttccaa ctcaggtaga acacacctgt ctttgtcttg
ctgtttttcac
1861 tcagcccttt taacattttc ccctaagccc atatgtctaa ggaaaggatg
10 ctatttggtgta
1921 atgaggaact gttatttgta tgtgaattaa agtgccttta tttt

```

	1:	Homo sapiens synovial sarcoma, NM_003147 X breakpoint 2 (SSX2), mRNA	PubMed, Protein, Related Sequences, Taxonomy, OMIM, LinkOut
15	LOCUS	NM_003147 766 bp mRNA	PRI 14-
	MAR-2001		
	DEFINITION	Homo sapiens synovial sarcoma, X breakpoint 2 (SSX2), mRNA.	
20	ACCESSION	NM_003147	
	VERSION	NM_003147.1 GI:10337582	
	KEYWORDS	.	
	SOURCE	human.	
	ORGANISM	<u>Homo sapiens</u>	
25		Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;	
		Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.	
	REFERENCE	1 (bases 1 to 766)	
30	AUTHORS	Shipley JM, Clark J, Crew AJ, Birdsall S, Rocques PJ, Gill S,	
		Chelly J, Monaco AP, Abe S, Gusterson BA and et al.	
	TITLE	The t(X;18)(p11.2;q11.2) translocation found in human synovial	
35		sarcomas involves two distinct loci on the X chromosome	
	JOURNAL	Oncogene 9 (5), 1447-1453 (1994)	
	MEDLINE	94203675	
	PUBMED	8152806	
	REFERENCE	2 (bases 1 to 766)	
40	AUTHORS	Crew,A.J., Clark,J., Fisher,C., Gill,S., Grimer,R., Chand,A.,	
		Shipley,J., Gusterson,B.A. and Cooper,C.S.	
	TITLE	Fusion of SYT to two genes, SSX1 and SSX2, encoding proteins with	
45		homology to the Kruppel-associated box in human synovial sarcoma	
	JOURNAL	EMBO J. 14 (10), 2333-2340 (1995)	
	MEDLINE	95292974	
	REFERENCE	3 (bases 1 to 766)	
50	AUTHORS	Tureci O, Sahin U, Schobert I, Koslowski M, Scmitt H, Schild HJ,	
		Stenner F, Seitz G, Rammensee HG and Pfreundschuh M.	
	TITLE	The SSX-2 gene, which is involved in the t(X;18) translocation of	

synovial sarcomas, codes for the human tumor antigen

HOM-MEL-40
 JOURNAL Cancer Res. 56 (20), 4766-4772 (1996)
 MEDLINE 96438636
 PUBMED 8840996

5 COMMENT PROVISIONAL REFSEQ: This record has not yet been
 subject to final
 NCBI review. The reference sequence was derived from

X86175.1.

10 FEATURES Location/Qualifiers
 source 1..766
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /chromosome="X"
 /map="Xp11.23-p11.22"
 /dev_stage="adult"
 15 gene 1..766
 /gene="SSX2"
 /note="HD21; HOM-MEL-40; SSX"
 20 /db_xref="LocusID:6757"
 /db_xref="MIM:300192"
 misc feature 20..61
 /note="Kruppel associated box homology"
 CDS 92..658
 25 /gene="SSX2"
 /codon_start=1
 /db_xref="LocusID:6757"
 /db_xref="MIM:300192"
 /product="synovial sarcoma, X breakpoint 2"
 30 /protein_id="NP_003138.1"
 /db_xref="GI:10337583"

/translation="MNGDDAFARRPTVGAQIPEKIQKAFDDIAKYFSKEEWEKMKASE

35 KIFYVYMKRKYEAMTKLGFKATLPPFCMCNKRAEDFQGNLDNDPNRGNQVERPQMTFG

RLQGISPKIMPKKPAEEGNDSEEVPRASGPQNDGKELCPPGKPTTSEKIHERSGPKRG

EHAWTHRLRERKQLVIYBEISDPEEDDE"

40 misc feature 161..337
 /note="KRAB; Region: krueppel associated box"

BASE COUNT 229 a 181 c 200 g 156 t

ORIGIN

1 ctctcttttcg attcttccat actcagagta cgcacggtct gattttctct
 ttggattctt

45 61 ccaaaatcag agtcagactg ctcccgggtgc catgaacgga gacgacgcct
 ttgcaaggag

121 acccacggtt ggtgctcaaa taccagagaa gatccaaaag gccttcgatg
 atattgccaa

181 atacttctct aaggaagagt gggaaaagat gaaagcctcg gagaaaatct

50 tctatgtgta

241 tatgaagaga aagtatgagg ctatgactaa actaggtttc aaggccaccc
 tcccaccttt

301 catgtgtaat aaacggggccg aagacttcca ggggaatgat ttggataatg
 accctaaccg

361 tgggaaatcag gttgaacgtc ctcatgatgac ttccggcagg ctccagggaa

55 tctccccgaa

421 gatcatgccc aagaagccag cagaggaagg aaatgattcg gaggaagtgc
 cagaagcatc

481 tggcccacaa aatgatggga aagagctgtg cccccggga aaaccaacta
 cctctgagaa
 541 gattcacgag agatctggac ccaaaagggg ggaacatgcc tggaccacaca
 gactgcgtga
 5 601 gagaaaacag ctggtgattt atgaagagat cagcgaccct gaggaagatg
 acgagtaact
 661 cccctcaggg atacgacaca tgcccatgat gagaagcaga acgtggtgac
 ctttcacgaa
 721 catgggcatg gctgcggacc cctcgtcacc aggtgcatag caagtg
 10

1: **Homo sapiens folate hydrolase** PubMed, Protein, Related
NM_004 (prostate-specific membrane antigen) Sequences, Taxonomy, OMIM,
 476 1 (FOLH1), mRNA LinkOut
 LOCUS NM_004476 2653 bp mRNA PRI 01-
 NOV-2000
 15 DEFINITION Homo sapiens folate hydrolase (prostate-specific
 membrane antigen)
 1 (FOLH1), mRNA.
 ACCESSION NM_004476
 VERSION NM_004476.1 GI:4758397
 20 KEYWORDS .
 SOURCE human.
 ORGANISM Homo sapiens
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
 Euteleostomi;
 25 Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
 Homo.
 REFERENCE 1 (bases 1 to 2653)
 AUTHORS Israeli,R.S., Powell,C.T., Fair,W.R. and Heston,W.D.
 TITLE Molecular cloning of a complementary DNA encoding a
 30 prostate-specific membrane antigen
 JOURNAL Cancer Res. 53 (2), 227-230 (1993)
 MEDLINE 93113576
 REFERENCE 2 (bases 1 to 2653)
 35 AUTHORS Rinker-Schaeffer CW, Hawkins AL, Su SL, Israeli RS,
 Griffin CA,
 Isaacs JT and Heston WD.
 TITLE Localization and physical mapping of the prostate-
 specific membrane
 antigen (PSM) gene to human chromosome 11
 40 JOURNAL Genomics 30 (1), 105-108 (1995)
 MEDLINE 96129312
 PUBMED 8595888
 REFERENCE 3 (bases 1 to 2653)
 45 AUTHORS O'Keefe DS, Su SL, Bacich DJ, Horiguchi Y, Luo Y,
 Powell CT,
 Zandvliet D, Russell PJ, Molloy PL, Nowak NJ, Shows
 TB, Mullins C,
 Vonder Haar RA, Fair WR and Heston WD.
 TITLE Mapping, genomic organization and promoter analysis of
 50 the human
 prostate-specific membrane antigen gene
 JOURNAL Biochim. Biophys. Acta 1443 (1-2), 113-127 (1998)
 MEDLINE 99057588
 PUBMED 9838072
 55 REFERENCE 4 (bases 1 to 2653)

AUTHORS Maraj BH, Leek JP, Karayi M, Ali M, Lench NJ and
 Markham AF.
 TITLE Detailed genetic mapping around a putative prostate-
 specific
 5 membrane antigen locus on human chromosome 11p11.2
 JOURNAL Cytogenet. Cell Genet. 81 (1), 3-9 (1998)
 MEDLINE 98358137
 PUBMED 9691167
 COMMENT PROVISIONAL REFSEQ: This record has not yet been
 10 subject to final
 NCBI review. The reference sequence was derived from
 M99487.1.
 FEATURES Location/Qualifiers
 source 1..2653
 15 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /chromosome="11"
 /map="11p11.2"
 /sex="male"
 20 /cell_line="LNCaP-ATCC"
 /cell_type="prostate"
 /tissue_type="prostatic carcinoma metastatic
 lymph node"
 /tissue_lib="LNCaP cDNA of Ron Israeli"
 25 gene 1..2653
 /gene="FOLH1"
 /note="FOLH; PSM; PSMA"
 /db_xref="LocusID:2346"
 /db_xref="MIM:600934"
 30 CDS 262..2514
 /gene="FOLH1"
 /note="folate hydrolase 1 (prostate-specific
 membrane
 antigen)"
 35 /codon_start=1
 /db_xref="LocusID:2346"
 /db_xref="MIM:600934"
 /evidence=experimental
 /product="folate hydrolase (prostate-specific
 40 membrane
 antigen) 1"
 /protein_id="NP_004467.1"
 /db_xref="GI:4758398"
 45 /translation="MWNLLHETDSAVATARRPRWLCAGALVLAGGFLLGFLFGWFIK
 SSNEATNITPKHNMKAFLDELKAENIKKFLYNFTQIPHLAGTEQNFQLAKQIQSQWKE
 FGLDSVELAHYDVLLSYPNKTHPNYISIIINEDGNEIFNTSLFEPFPPGYENVSDIVPP
 50 FSAFSPQGMPEGDLVYVNYARTEDFFKLERDMKINCSGKIVIARYGKVFRGNKVKNAQ
 LAGAKGVILYSDPADYFAPGVKSYPDGWNLPGGGVQRGNIILNLNGAGDPLTPGYANE
 YAYRRGIAEAVGLPSIPVHPIGYYDAQKLEKMGGSAAPPDSSWRGSLKVPYNVPGPFT
 55 GNFSTQKVKMHIHSTNEVTRIYNVIGTLRGAVEPDRYVILGGHRDSWVFGGIDPQSGA

AVVHEIVRSFGTLKKEGWRPRRTILFASWDAEEFGLLGSTEWAEENSRLQLQERGVAII
 NADSSIEGNYTLRVDCPTPLMYSLVHNLTKELKSPDEGFEGKSLYESWTKKSPSPEFSG
 5 MPRISKLGSGNDFEVFFQRLGIASGRARYTKNWETNKFSGYPLYHSVYETYELVEKFY
 DPMFKYHLTVAQVRGGMVFELANSIVLPFDCRDYAVVLRKYADKIYSISMKHPQEMKT
 10 YSVSFDLSLFAVKNFTEIASKFSERLQDFDKSNPIVLRMMNDQLMFLERAFIDPLGLP
 DRPFYRHVIYAPSSHNKYAGESFPGIYDALFDIESKVDPSKAWGEVVKRQIYVAAFTVQ
 AAAETLSEVA"
 15 misc feature 778..1029
 /note="PA; Region: PA domain"
 BASE COUNT 782 a 524 c 640 g 707 t
 ORIGIN
 1 ctcaaaagg ggcggatttc cttctcctgg aggcagatgt tgcctctctc
 tctcgtctcg
 20 61 attggttcag tgcactctag aaacactgct gtggtggaga aactggaccc
 caggctctgga
 121 gcgaattcca gcctgcaggg ctgataagcg aggcattagt gagattgaga
 gagactttac
 181 ccgcgcgtgg tgggttgagg gcgcgcagta gagcagcagc acaggcgagg
 25 gtcccgagg
 241 gccggctctg ctgcgcgcga gatgtggaat ctccctcacg aaaccgactc
 ggctgtggcc
 301 accgcgcgcc gccgcgcgtg gctgtgcgct ggggcgcgtg tgctggcggg
 tggcttcttt
 361 ctccctcggt tccctcttcg gtggtttata aaatcctcca atgaagctac
 taacattact
 421 ccaaagcata atatgaaagc atttttggat gaattgaaag ctgagaacat
 caagaagttc
 481 ttatataatt ttacacagat accacattta gcaggaacag aacaaaactt
 35 tcagcttgca
 541 aagcaaattc aatcccagtg gaaagaattt ggcttgatt ctggtgagct
 agcacattat
 601 gatgtcctgt tgtcctaccc aaataagact catcccaact acatctcaat
 aattaatgaa
 40 661 gatggaaatg agattttcaa cacatcatta tttgaaccac ctccctcagg
 atatgaaaat
 721 gtttcggata ttgtaccacc tttcagtgtt ttctctcctc aaggaatgcc
 agagggcgat
 781 ctagtgtatg ttaactatgc acgaactgaa gacttcttta aattggaacg
 45 ggacatgaaa
 841 atcaattgct ctgggaaaat tgtaattgcc agatatggga aagttttcag
 aggaaataag
 901 gttaaaaatg ccagctggc aggggccaaa ggagtcattc tctactccga
 ccctgctgac
 50 961 tactttgtc ctggggtgaa gtcctatcca gatggttgga atcttctcgg
 aggtggtgtc
 1021 cagcgtggaa atatcctaaa tctgaatggt gcaggagacc ctctcacacc
 aggttaccga
 1081 gcaaatgaat atgcttatag gcgtggaatt gcagaggctg ttggtcttcc
 55 aagtattcct
 1141 gttcatccaa ttggatacta tgatgcacag aagctcctag aaaaaatggg
 tggtcagca

1201 ccaccagata gcagctggag aggaagtctc aaagtgccct acaatgttgg
 acctggcttt
 1261 actggaaact tttctacaca aaaagtcaag atgcacatcc actctacaa
 tgaagtgaca
 5 1321 agaatttaca atgtgatagg tactctcaga ggagcagtgg aaccagacag
 atatgtcatt
 1381 ctgggaggtc accgggactc atgggtgttt ggtggtattg accctcagag
 tggagcagct
 1441 gttgttcatg aaattgtgag gagctttgga aacttgaaaa aggaaggggtg
 10 gagacctaga
 1501 agaacaattt tgtttgcaag ctgggatgca gaagaatttg gtcttcttgg
 ttctactgag
 1561 tgggcagagg agaattcaag actccttcaa gagcgtggcg tggcttatat
 taatgctgac
 1621 tcacttatag aaggaaaacta cactctgaga gttgattgta caccgctgat
 15 gtacagcttg
 1681 gtacacaacc taacaaaaga gctgaaaagc cctgatgaag gctttgaagg
 caaatctctt
 1741 tatgaaagtt ggactaaaaa aagtccttcc ccagagttca gtggcatgcc
 20 caggataagc
 1801 aaattgggat ctggaaatga ttttgagggtg ttcttccaac gacttggaaat
 tgcttcaggc
 1861 agagcacggt atactaaaaa ttgggaaaca aacaaattca gcggctatcc
 actgtatcac
 1921 agtgtctatg aaacatatga gttgggtggaa aagttttatg atccaatggt
 25 taaatatcac
 1981 ctcactgtgg cccagggttcg aggagggatg gtgtttgagc tagccaatcc
 catagtgtc
 2041 ccttttgatt gtcgagatta tgctgtagtt ttaagaaagt atgctgacaa
 30 aatctacagt
 2101 atttctatga aacatccaca ggaaatgaag acatacagtg tatcatttga
 ttcacttttt
 2161 tctgcagtaa agaattttac agaaattgct tccaagttca gtgagagact
 ccaggacttt
 2221 gacaaaagca acccaatagt attaagaatg atgaatgac aactcatggt
 35 tctggaaaaga
 2281 gcattttattg atccattagg gttaccagac aggccttttt ataggcatgt
 catctatgct
 2341 ccaagcagcc acaacaagta tgcaggggag tcattcccag gaatttatga
 40 tgctctgttt
 2401 gatattgaaa gcaaagtgga cccttccaag gcctggggag aagtgaagag
 acagatttat
 2461 gttgcagcct tcacagtga ggcagctga gagactttga gtgaagtagc
 ctaagaggat
 2521 tcttttagaga atccgtattg aatttgtgtg gtatgtcact cagaaagaat
 45 cgtaatgggt
 2581 atattgataa attttaaaat tggatatatt gaaataaagt tgaatattat
 atataaaaaa
 2641 aaaaaaaaaa aaa

U20093. Human melanocyte-...[gi:1142634]
 LOCUS HSPMEL17S2 2817 bp DNA PRI 20-
 NOV-1996
 DEFINITION Human melanocyte-specific (pmel 17) gene, exons 2-5,
 and complete
 cds.
 ACCESSION U20093
 VERSION U20093.1 GI:1142634
 KEYWORDS .
 SEGMENT 2 of 2
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
 Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
 Homo.
 REFERENCE 1 (bases 1 to 2817)
 AUTHORS Kim,K.K., Youn,B.S., Heng,H.H., Shi,X.M., Tsui,L.C.,
 Lee,Z.H.,
 Pickard,R.T. and Kwon,B.S.
 TITLE Genomic organization and FISH mapping of human Pmel
 17, the
 putative silver locus
 JOURNAL Pigment Cell Res. 9 (1), 42-48 (1996)
 MEDLINE 96314705
 REFERENCE 2 (bases 1 to 2817)
 AUTHORS Kwon,B.S.
 TITLE Direct Submission
 JOURNAL Submitted (05-JAN-1995) Indiana University School of
 Medicine,
 Microbiology and Immunology, 635 Barnhill Drive,
 Indianapolis, IN
 46202, USA
 FEATURES Location/Qualifiers
 source 1..2817
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /chromosome="12"
 /map="12pter-q21"
 /clone_lib="lambda FIX II from Stratagene"
 gene join(U19491.1:1111..3109,1..2722)
 /gene="pmel 17"
 CDS
 join(U19491.1:1111..1186,67..459,853..1014,1266..2396,
 2499..2722)
 /gene="pmel 17"
 /codon_start=1
 /product="Pmel 17"
 /protein_id="AAB19181.1"
 /db_xref="GI:1142636"
 /translation="MDLVLRCLLHLAVIGALLAVGATKVPNRQDWLGVSRLRTKAWNROLYPEW
 TEAQRLLDCWRGGQVSLKVSNDGPTLIGANASFSIALNFPGSQKVLDPGQVIWVNNTIINGSQVWGG
 QPVYPQETDDACIFPDGGPCPSGSWSQKRSFVYVWKTWGQYQVVLGGPVSGLSIGTGRAMLGTHM
 EVTVYHRRGSRSYVPLAHSSSAFTITDQVPFVSQVSLRALDGGNKHFLRNQPLTFALQLHDPSTGY
 LAEADLSYTWDFGDSSGTLISRAPVVTHYLEPGPVTAQVVLQAAIPLTSCGSSPVPVPGTTDGHRT
 AEAPNTTAGQVPTTEVVGTTPGQAPTAEPSTTSVQVPTTEVISTAPVQMPTAESTGMTPEKVPVS
 EVMGTTLAEMSTPEATGMTPAEVSIVVLSGTTAAQVTTTEWVETTARELPIPEPEGPDASSIMSTE"

SITGSLGPLLDGTATLRLVKRQVPLDCVLYRYGSFSVTLDIVQGIESAEILQAVPSGEGDAFELTV
 SCQGGLPKEACMEISSPGCQPPAQRQCQVLPSPACQLVLHQILKGGSGTYCLNVSLADTNSLAVV
 STQLIMPGQEAGLGQVPLIVGILLVLMVVLASLIYRRRLMKQDFSVPLPHSSSHWLRLPRIFCS
 CPIGENSPLLSGQV"

5	exon	67..459				
		/gene="pmel 17"				
		/number=2				
	exon	853..1014				
10		/gene="pmel 17"				
		/number=3				
	exon	1266..2396				
		/gene="pmel 17"				
		/number=4				
	exon	2499..>2722				
15		/gene="pmel 17"				
		/number=5				
	BASE COUNT	627 a	772 c	739 g	679 t	
	ORIGIN					
	1	gtgctaaaaa	gatgccttct	tcatttggct	gtgatagggtg	ctttgtggct
20	gtgggggcta	61 caaaagtacc	cagaaaccag	gactggcttg	gtgtctcaag	gcaactcaga
	accaaagcct	121 ggaacaggca	gctgtatcca	gagtggacag	aagcccagag	acttgactgc
	tggagaggtg	25 181 gtcaagtgtc	cctcaagggtc	agtaatgatg	ggcctacaact	gattggtgca
	aatgcctcct	241 tctctattgc	cttgaacttc	cctggaagcc	aaaaggtatt	gccagatggg
	caggttatct	301 ggggtcaacaa	taccatcatc	aatgggagcc	aggtgtgggg	aggacagcca
30	gtgtatcccc	361 aggaaactga	cgatgcctgc	atcttccctg	atggtggacc	ttgcccattc
	ggctcttggg	421 ctccagaagag	aagctttggt	tatgtctgga	agacctgggg	tgagggaactc
	ccttctcagc	35 481 ctatcatcca	cacttgtggt	tacttctttc	tacctgatca	cctttctttt
	ggccgcccct	541 tccaccttaa	cttctgtgat	tttctcta	cttcattttc	ctcttagatc
	ttttctcttt	601 cttagcacct	agcccccttc	aagctctatc	ataattcttt	ctggcaactc
40	ttggcctcaa	661 ttgtagtcct	accccatgga	atgcctcatt	aggaccccctt	ccctgtcccc
	ccatatcaca	721 gccttccaaa	caccctcaga	agtaatcata	cttcctgacc	tcccatctcc
	agtgcctgtt	45 781 cgaagcctgt	ccctcagtc	cctttgacca	gtaatctctt	cttccttgc
	tttcattcca	841 aaaatgcttc	aggccaatac	tggcaagttc	tagggggccc	agtgtctggg
	ctgagcattg	901 ggacaggcag	ggcaatgctg	ggcacacaca	ccatggaagt	gactgtctac
50	catcgccggg	961 gatcccgagg	ctatgtgcct	cttgctcatt	ccagctcagc	cttcaccatt
	actggtaagg	1021 gttcagggaag	ggcaaggcca	gttgtagggc	aaagagaagg	cagggaggct
	tggatggact	55 1081 gcaaaggaga	aaggtgaaat	gctgtgcaaa	cttaaagtag	aagggccagg
	aagacctagg	1141 cagagaaatg	tgaggcttag	tgccagtga	gggccagcca	gtcagcttgg
	agttggaggg					

```

1201 tgtggctgtg aaaggagaag ctgtggctca ggcctgggtc tcaccttttc
tggtccaat
1261 cccagaccag gtgcctttct ccgtgagcgt gtcccagttg cgggccttgg
atggagggaa
5 1321 caagcacttc ctgagaaatc agcctctgac ctttgccctc cagctccatg
accccagtgg
1381 ctatctggct gaagctgacc tctcctacac ctgggacttt ggagacagta
gtggaaccct
1441 gatctctcgg gcacctgtgg tcactcatac ttacctggag cctggcccag
10 tcaactgcca
1501 ggtggtcctg caggctgcca ttctctcac ctctgtggc tcctccccag
ttccaggcac
1561 cacagatggg cacaggccaa ctgcagaggc ccctaacacc acagctggcc
aagtgcctac
15 1621 tacagaagtt gtgggtacta cacctgggtc ggcgccaaact gcagagccct
ctggaaccac
1681 atctgtgcag gtgccaacca ctgaagtcac aagcactgca cctgtgcaga
tgccaactgc
1741 agagagcaca ggtatgacac ctgagaaggt gccagtttca gaggtcatgg
20 gtaccacact
1801 ggcagagatg tcaactccag aggtacagg tatgacacct gcagaggtat
caattgtggt
1861 gctttctgga accacagctg cacaggtaac aactacagag tgggtggaga
ccacagctag
25 1921 agagctacct atccctgagc ctgaagggtc agatgccagc tcaatcatgt
ctacggaaag
1981 tattacaggt tccctgggcc ccctgctgga tggtagagcc accttaaggc
tggtagagag
2041 acaagtcccc ctggattgtg ttctgtatcg atatggttcc ttttcctgca
30 ccctggacat
2101 tgtccagggt attgaaagtg ccgagatcct gcaggctgtg ccgtccggtg
agggggatgc
2161 atttgagctg actgtgtcct gccaaaggcg gctgccccaa gaagcctgca
tggagatctc
35 2221 atcgccaggg tgccagcccc ctgccagcg gctgtgccag cctgtgctac
ccagcccagc
2281 ctgccagctg gttctgcacc agatactgaa gggtaggctc gggacatact
gcctcaatgt
2341 gtctctggct gataccaaca gcctggcagt ggtcagcacc cagcttatca
40 tgccctggtag
2401 gtccctggac agagactaag tgaggaggga agtggataga ggggacagct
ggcaagcagc
2461 agacatgagt gaagcagtgc ctgggattct tctcacaggt caagaagcag
gccttgggca
45 2521 ggttccgctg atcgtgggca tcttgctggt gttgatggct gtggtccttg
catctctgat
2581 atataggcgc agacttatga agcaagactt ctccgtaccc cagttgccac
atagcagcag
2641 tcaactggctg cgtctacccc gcattctctg ctcttgctcc attggtgaga
50 atagccccct
2701 cctcagtggg cagcaggtct gagtactctc atatgatgct gtgattttcc
tgaggttgac
2761 agaaacacct atatttcccc cagtcttccc tgggagacta ctattaactg
aaataaaa
55 //

```

NM_001648. Homo sapiens kall...[gi:4502172]

LOCUS NM_001648 1466 bp mRNA PRI
 31-OCT-2000
 DEFINITION Homo sapiens kallikrein 3, (prostate specific antigen)
 (KLK3),
 5 mRNA.
 ACCESSION NM_001648
 VERSION NM_001648.1 GI:4502172
 KEYWORDS .
 SOURCE human.
 10 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata;
 Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini;
 Hominidae; Homo.
 15 REFERENCE 1 (bases 1 to 1466)
 AUTHORS Lundwall, A. and Lilja, H.
 TITLE Molecular cloning of human prostate specific antigen
 cDNA
 JOURNAL FEBS Lett. 214 (2), 317-322 (1987)
 20 MEDLINE 87190978
 REFERENCE 2 (bases 1 to 1466)
 AUTHORS Sutherland GR, Baker E, Hyland VJ, Callen DF, Close JA,
 Tregear GW,
 Evans BA and Richards RI.
 25 TITLE Human prostate-specific antigen (APS) is a member of
 the glandular
 kallikrein gene family at 19q13
 JOURNAL Cytogenet. Cell Genet. 48 (4), 205-207 (1988)
 MEDLINE 89250658
 30 PUBMED 2470553
 REFERENCE 3 (bases 1 to 1466)
 AUTHORS Riegman PH, Klaassen P, van der Korput JA, Romijn JC
 and Trapman J.
 TITLE Molecular cloning and characterization of novel
 35 prostate antigen
 cDNA's
 JOURNAL Biochem. Biophys. Res. Commun. 155 (1), 181-188 (1988)
 MEDLINE 88326297
 PUBMED 2458104
 40 REFERENCE 4 (bases 1 to 1466)
 AUTHORS Schulz P, Stucka R, Feldmann H, Combriato G, Klobeck HG
 and Fittler F.
 TITLE Sequence of a cDNA clone encompassing the complete
 mature human
 45 prostate specific antigen (PSA) and an unspliced
 leader sequence
 JOURNAL Nucleic Acids Res. 16 (13), 6226 (1988)
 MEDLINE 88289366
 PUBMED 2456523
 50 REFERENCE 5 (bases 1 to 1466)
 AUTHORS Riegman PH, Vlietstra RJ, van der Korput JA, Romijn JC
 and Trapman J.
 TITLE Characterization of the prostate-specific antigen gene:
 a novel
 55 human kallikrein-like gene
 JOURNAL Biochem. Biophys. Res. Commun. 159 (1), 95-102 (1989)
 MEDLINE 89165891
 PUBMED 2466464

REFERENCE 6 (bases 1 to 1466)
AUTHORS Henttu P and Vihko P.
TITLE cDNA coding for the entire human prostate specific
antigen shows high homologies to the human tissue kallikrein
genes
JOURNAL Biochem. Biophys. Res. Commun. 160 (2), 903-910 (1989)
MEDLINE 89246551
PUBMED 2470373
REFERENCE 7 (bases 1 to 1466)
AUTHORS Klobeck HG, Combriato G, Schulz P, Arbusow V and
Fittler F.
TITLE Genomic sequence of human prostate specific antigen
(PSA)
JOURNAL Nucleic Acids Res. 17 (10), 3981 (1989)
MEDLINE 89282407
PUBMED 2471958
REFERENCE 8 (bases 1 to 1466)
AUTHORS Lundwall A.
TITLE Characterization of the gene for prostate-specific
antigen, a human glandular kallikrein
JOURNAL Biochem. Biophys. Res. Commun. 161 (3), 1151-1159
(1989)
MEDLINE 89302090
PUBMED 2472789
COMMENT PROVISIONAL REFSEQ: This record has not yet been
subject to final NCBI review. The reference sequence was derived
from X05332.1.
FEATURES Location/Qualifiers
source 1..1466
/organism="Homo sapiens"
/db_xref="taxon:9606"
/chromosome="19"
/map="19q13"
/clone="lambda HPSA-1"
/tissue_type="prostate"
/clone_lib="(lambda)gt11"
gene 1..1466
/gene="KLK3"
/note="PSA; APS"
/db_xref="LocusID:354"
/db_xref="MIM:176820"
sig_peptide 44..94
CDS 44..829
/gene="KLK3"
/EC_number="3.4.21.77"
/note="antigen, prostate specific"
/codon_start=1
/db_xref="LocusID:354"
/db_xref="MIM:176820"
/product="kallikrein 3, (prostate specific
antigen)"
/protein_id="NP_001639.1"
/db_xref="GI:4502173"

```

/translation="MWVPVVFLLSVTWIGAAPLILSRIVGGWECEKHSQPWQVLVAS
RGRAVCGGVLVHPQWVLTAAHCIRNKSIVLLGRHSLFHPEDTGQVFQVSHSFPHPYDMSLLKNRF
LRPGDDSSHDMLLRLSEPAELTDAVKVMDLPTQEPALGTTTCYASGWGSIEPEEFLLTPKKLQCVDL
5 HVISNDVCAQVHPQKVTKFMLCAGRWTGGKSTCSGDSGGPLVCNGVLQGITSWGSEPCALPERPSL
YTKVVHYRKWIKDTIVANP"
      variation      91
                        /allele="C"
                        /allele="T"
10      variation      91
                        /db_xref="dbSNP:1135765"
                        /allele="C"
                        /allele="T"
                        /db_xref="dbSNP:11573"
15      misc_feature    95..115
                        /note="propeptide (AA 1-7)"
      variation      97
                        /allele="G"
                        /allele="A"
20      misc_feature    113..802
                        /db_xref="dbSNP:1135766"
                        /note="Tryp_SPc; Region: Trypsin-like serine
protease"
      misc_feature    116..802
25      mat_peptide     116..826
                        /note="trypsin; Region: Trypsin"
                        /product="kallikrein 3, (prostate specific
antigen)"
      variation      280
30      variation      280
                        /allele="C"
                        /allele="T"
                        /db_xref="dbSNP:1058072"
      variation      280
35      variation      280
                        /allele="C"
                        /allele="T"
                        /db_xref="dbSNP:12946"
      variation      526
                        /allele="A"
                        /allele="G"
40      variation      643
                        /db_xref="dbSNP:1803130"
                        /allele="A"
                        /allele="G"
                        /db_xref="dbSNP:1803129"
45      variation      788
                        /note="WARNING: map location ambiguous"
                        /allele="T"
                        /allele="C"
                        /db_xref="dbSNP:1802720"
50      variation      844
                        /allele="T"
                        /allele="C"
                        /db_xref="dbSNP:1058205"
      variation      894
55      variation      894
                        /allele="A"
                        /allele="G"
                        /db_xref="dbSNP:1058274"
      variation      906

```

		/allele="T"
		/allele="A"
		/db_xref="dbSNP:1803132"
5	variation	918
		/allele="T"
		/allele="C"
		/db_xref="dbSNP:1802722"
	variation	953
10		/allele="A"
		/allele="G"
		/db_xref="dbSNP:1802721"
	variation	1017
		/allele="T"
		/allele="C"
15		/db_xref="dbSNP:1803138"
	variation	1074
		/allele="C"
		/allele="T"
		/db_xref="dbSNP:12101"
20	variation	1107
		/allele="A"
		/allele="G"
		/db_xref="dbSNP:1803131"
	variation	1107
25		/allele="G"
		/allele="A"
		/db_xref="dbSNP:6998"
	variation	1211
		/allele="C"
30		/allele="T"
		/db_xref="dbSNP:1803136"
	variation	1229
		/allele="A"
		/allele="G"
35		/db_xref="dbSNP:1803135"
	variation	1249
		/allele="T"
		/allele="C"
		/db_xref="dbSNP:1803134"
40	variation	1273
		/allele="A"
		/allele="G"
		/db_xref="dbSNP:12040"
	variation	1326
45		/allele="T"
		/allele="C"
		/db_xref="dbSNP:1803133"
	variation	1344
		/allele="A"
50		/allele="G"
		/db_xref="dbSNP:12863"
	variation	1363
		/allele="A"
		/allele="G"
55		/db_xref="dbSNP:1803137"
	misc_feature	1445..1450
		/note="put.polyA signal"
	polyA_site	1466

```

                                /note="polyA site"
BASE COUNT      338 a      382 c      422 g      324 t
ORIGIN
5      1 agccccaagc ttaccacctg caccceggaga gctgtgtgtc accatgtggg
      61 cttcctcacc ctgtccgtga cgtggattgg tgctgcaccc ctcacacctgt
      121 gggaggctgg gagtgcgaga agcattccca accctggcag gtgcttgtgg
10     181 cagggcagtc tgcggcgggtg ttctgggtgca cccccagtgg gtcctcacag
      241 catcaggaac aaaagcgtga tcttgtctggg tcggcacagc ctgtttcatc
      301 aggccaggta tttcaggtea gccacagctt cccacacccg ctctacgata
15     361 gaagaatcga ttcctcaggc caggtgatga ctccagccac gacctcatgc
      421 gtcagagcct gccgagctca cggatgctgt gaaggctcatg gacctgccc
      481 agcactgggg accacctgct acgcctcagg ctggggcagc attgaaccag
20     541 gaccccaaag aaacttcagt gtgtggacct ccatgttatt tccaatgacg
      601 agttcacctc cagaaggatga ccaagtcat gctgtgtgct ggacgctgga
25     661 aagcacctgc tcgggtgatt ctgggggccc acttgtctgt aatggtgtgc
      721 cacgtcatgg ggcagtgaac catgtgccct gcccgaaagg ccttcctctgt
      781 ggtgcattac cggaagtgga tcaaggacac catcgtggcc aacccttgag
30     841 aacccttatc tgtagtaaac ttggaacctt ggaaatgacc aggccaagac
      901 ccagttctac tgacctttgt ccttaggtgt gaggtccagg gttgctagga
35     961 gcagacacag gtgtagacca gagtgtttct taaatggtgt aattttgtcc
      1021 ctggggaata ctggccatgc ctggagacat atcactcaat ttctctgagg
      1081 gatgggggtgt ctgtgttatt tgtggggtac agagatgaaa gaggggtggg
40     1141 agagagtgga gagtgcacat tgctggacac tgtccatgaa gcaactgagca
      1201 gcacaacgca ccagacactc acagcaagga tggagctgaa aacataaccc
45     1261 ggaggcactg ggaagcctag agaaggctgt gagccaagga gggagggctc
      1321 tgggatgggg atgaagtaag gagagggact ggacccctg gaagctgatt
      1381 ggaggtgtat tgaagtcctc cagacaaccc tcagatttga tgatttccta
50     1441 cagaaataaa gagctgttat actgtg
      //

55     U87459. Human autoimmunog...[gi:1890098]
      LOCUS      HSU87459      752 bp      mRNA      PRI
      22-DEC-1999

```


DEFINITION Human autoimmunogenic cancer/testis antigen NY-ESO-1 mRNA, complete cds.
 ACCESSION U87459
 VERSION U87459.1 GI:1890098
 5 KEYWORDS .
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata;
 10 Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini;
 Hominidae; Homo.
 REFERENCE 1 (bases 1 to 752)
 AUTHORS Chen,Y.T., Scanlan,M.J., Sahin,U., Tureci,O.,
 Gure,A.O., Tsang,S.,
 15 Williamson,B., Stockert,E., Pfreundschuh,M. and
 Old,L.J.
 TITLE A testicular antigen aberrantly expressed in human
 cancers detected
 by autologous antibody screening
 20 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 94 (5), 1914-1918 (1997)
 MEDLINE 97203161
 PUBMED 9050879
 REFERENCE 2 (bases 1 to 752)
 AUTHORS Chen,Y.-T.
 25 TITLE Direct Submission
 JOURNAL Submitted (28-JAN-1997) Ludwig Institute for Cancer
 Research, New
 York Branch, 1275 York Avenue, New York, NY 10021,
 30 USA
 FEATURES Location/Qualifiers
 source 1..752
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /clone_lib="esophageal squamous cell
 35 carcinoma expression
 cDNA library"
 CDS 54..596
 /codon_start=1
 /product="autoimmunogenic cancer/testis
 40 antigen NY-ESO-1"
 /protein_id="AAB49693.1"
 /db_xref="GI:1890099"
 /translation="MQAEGRGTTGGSTGDADGPGGPGIPDGPGGNAGGPGEAG
 45 ATGGRGPRGAGAARASGPGGGAPRGPHGGAASGLNGCCRCGARGPESRLLEF
 YLAMPFATPMEAEIARRSLAQDAPPLFVPGVLLKEFTVSGNILTIRLTAADH
 RQLQLSISSCLQLSLLMWITQCFLPVFLAQPPSGQRR"
 BASE COUNT 126 a 230 c 256 g 140 t
 ORIGIN
 50 1 atcctcgtgg gccctgacct tctctctgag agccgggcag aggctccgga
 gccatgcagg
 61 ccgaaggccg gggcacaggg ggttcgacgg gcgatgctga tggcccagga
 ggccctggca
 121 ttctgatgg cccagggggc aatgctggcg gcccaggaga ggcgggtgcc
 55 acgggcggca
 181 gaggtccccg gggcgcaggg gcagcaaggg cctcggggcc gggaggaggc
 gccccgcggg

```

241 gtccgcatgg cggcgcggtc tcagggtga atggatgctg cagatgcggg
gccagggggc
301 cggagagccg cctgcttgag ttctacctcg ccatgccttt cgcgacaccc
atggaagcag
5 361 agctggcccc caggagcctg gcccaggatg ccccacccgt tcccggtcca
gggggtgcttc
421 tgaaggagtt cactgtgtcc ggcaacatac tgactatccg actgactgct
gcagaccacc
481 gccaaactgca gctctccatc agctcctgtc tccagcagct ttccctgttg
10 atgtggatca
541 cgcagtgctt tctgcccgtg tttttggctc agcctccctc agggcagagg
cgctaagccc
601 agcctggcgc cccttccctag gtcatgcctc ctcccctagg gaatgggtccc
agcacgagtg
15 661 gccagttcat tgtggggggc tgattgtttg togtggagg aggacggctt
acatgtttgt
721 ttctgtagaa aataaaactg agctacgaaa aa
//

20 CAA11116. LAGE-1a protein [...[gi:3255959]
LOCUS CAA11116 180 aa PRI
23-JUN-1998
DEFINITION LAGE-1a protein [Homo sapiens].
ACCESSION CAA11116
25 PID g3255959
VERSION CAA11116.1 GI:3255959
DBSOURCE embl locus HOS223093, accession AJ223093.1
KEYWORDS .
SOURCE human.
30 ORGANISM Homo sapiens
Eukaryota; Metazoa; Chordata; Craniata;
Vertebrata; Euteleostomi;
Mammalia; Eutheria; Primates; Catarrhini;
Hominidae; Homo.
35 REFERENCE 1 (residues 1 to 180)
AUTHORS Lethe,B., Lucas,S., Michaux,L., De Smet,C.,
Godelaine,D.,
Serrano,A., De Plaen,E. and Boon,T.
40 TITLE LAGE-1, a new gene with tumor specificity
JOURNAL Int. J. Cancer 76 (6), 903-908 (1998)
MEDLINE 98289662
REFERENCE 2 (residues 1 to 180)
AUTHORS Lethe,B.G.
45 TITLE Direct Submission
JOURNAL Submitted (08-JAN-1998) Lethe B.G., Brussels Branch,
Ludwig
Institute for Cancer Research, 74 Avenue
Hippocrate, B - 1200 -
Bruxelles, BELGIUM
50 COMMENT Related sequences: AJ223040, AJ223041 and AJ003149.
FEATURES Location/Qualifiers
source 1..180
/organism="Homo sapiens"
/isolate="individual LB33"
55 /db_xref="taxon:9606"
/chromosome="X"
/map="q28"
/clone="c2RB"

```

```

                    /sex="Female"
                    /cell_line="LB33-MELA"
                    /cell_type="melanoma"
                    /clone_lib="B6"
5      Protein      1..180
                    /product="LAGE-1a protein"
      CDS           1..180
                    /gene="LAGE-1"
                    /db_xref="SPTREMBL:O75637"
10      /coded_by="join(AJ223093.1:1231..1499,
AJ223093.1:2115..2249,AJ223093.1:2479..2617)"
ORIGIN
15      1 mqaegrgtgg stgdadgpgg pgipdgpggn aggpgeagat gggrgprgaga
arasgprgga
      61 prgphggaas aqdgrocpca rrpdsrllel hitmpfsspm eaelvrrils
rdaaplprpg
      121 avlkdfvtvg nllfirltaa dhrqlqlsis sclqqslslm witqcflpvf
laqapsggrr
20      181
//

CAA11117. LAGE-1b protein [...[gi:3255960]
LOCUS      CAA11117      210 aa      PRI
25 23-JUN-1998
DEFINITION LAGE-1b protein [Homo sapiens].
ACCESSION CAA11117
PID      g3255960
VERSION  CAA11117.1 GI:3255960
30 DBSOURCE  embl locus HOS223093, accession AJ223093.1
KEYWORDS
SOURCE    human.
ORGANISM  Homo sapiens
          Eukaryota; Metazoa; Chordata; Craniata;
35 Vertebrata; Euteleostomi;
          Mammalia; Eutheria; Primates; Catarrhini;
Hominidae; Homo.
REFERENCE  1 (residues 1 to 210)
AUTHORS   Lethe,B., Lucas,S., Michaux,L., De Smet,C.,
40 Godelaine,D.,
          Serrano,A., De Plaen,E. and Boon,T.
TITLE     LAGE-1, a new gene with tumor specificity
JOURNAL   Int. J. Cancer 76 (6), 903-908 (1998)
MEDLINE   98289662
45 REFERENCE 2 (residues 1 to 210)
AUTHORS   Lethe,B.G.
TITLE     Direct Submission
JOURNAL   Submitted (08-JAN-1998) Lethe B.G., Brussels Branch,
50 Ludwig
          Institute for Cancer Research, 74 Avenue
Hippocrate, B - 1200 -
          Bruxelles, BELGIUM
COMMENT   Related sequences: AJ223040, AJ223041 and AJ003149.
FEATURES  Location/Qualifiers
55      source      1..210
                    /organism="Homo sapiens"
                    /isolate="individual LB33"
                    /db_xref="taxon:9606"

```

```

                    /chromosome="X"
                    /map="q28"
                    /clone="c2RB"
                    /sex="Female"
5                    /cell_line="LB33-MELA"
                    /cell_type="melanoma"
                    /clone_lib="B6"
                Protein 1..210
                    /product="LAGE-1b protein"
10                CDS 1..210
                    /gene="LAGE-1"
                    /db_xref="SPTREMBL:O75638"
                    /coded_by="join(AJ223093.1:1231..1499,
15                    AJ223093.1:2115..2478)"
                    /note="alternate splicing at exon 2 - intron
                2 boundary"
                ORIGIN
                    1 mqaegrgtgg stgdadgpgg pgipdgpagn aggpgeagat ggrgprgaga
                arasgprgga
20                    61 prgphggaas aqdgrcpcga rrpdsrllel hitmpfsspm eaelvrrils
                rdaaplprpg
                    121 avlkdfvtsg nllfmsvwdq dregagrmrv vgwglgsasp egqkardlrt
                pkhkvsegrp
                    181 gtpgpppppeg aqgdgcrgva fnvmfsaphi
25                //

30                M77481. Human antigen (MA...[gi:416114]
                LOCUS HUMMAG1A 2420 bp DNA PRI
                15-NOV-1993
                DEFINITION Human antigen (MAGE-1) gene, complete cds.
                ACCESSION M77481
35                VERSION M77481.1 GI:416114
                KEYWORDS antigen.
                SOURCE Homo sapiens (individual_isolate patient MZ2)
                melanoma abdominal
                                metastasis of melanoma DNA.
40                ORGANISM Homo sapiens
                                Eukaryota; Metazoa; Chordata; Craniata;
                Vertebrata; Euteleostomi;
                                Mammalia; Eutheria; Primates; Catarrhini;
                Hominidae; Homo.
45                REFERENCE 1 (bases 785 to 1286)
                AUTHORS van der Bruggen,P., Traversari,C., Chomez,P.,
                Lurquin,C., De
                                Plaen,E., Van den Eynde,B., Knuth,A. and Boon,T.
                TITLE A gene encoding an antigen recognized by cytolytic T
50                lymphocytes on
                                a human melanoma
                JOURNAL Science 254, 1643-1647 (1991)
                MEDLINE 92086861
                REFERENCE 2 (bases 1 to 2420)
55                AUTHORS van der Bruggen P.
                TITLE Direct Submission
                JOURNAL Submitted (05-FEB-1992) Pierre van der Bruggen, Ludwig
                Institute

```

for Cancer Research, Brussels Branch, Avenue
Hippocrate, 74, UCL
7459, Brussels, B-1200, Belgium

COMMENT On Nov 15, 1993 this sequence version replaced
gi:187294.

5 FEATURES Location/Qualifiers

source 1..2420
/organism="Homo sapiens"
/db_xref="taxon:9606"

10 /map="X"
/sex="female"
/cell_line="MZ2-MEL.43"

intron 1..412
/partial

15 /gene="MAGE-1"
/number=1

exon 413..485
/gene="MAGE-1"
/number=2

20 mRNA join(413..485,561..2111)
/partial
/gene="MAGE-1"

intron 486..560
/gene="MAGE-1"
/number=2

25 exon 561..2111
gene="MAGE-1"
/number=3

gene 626..1555
/gene="MAGE-1"

30 CDS 626..1555
/gene="MAGE-1"
/codon_start=1
/product="MAGE-1"
/protein_id="AAA03229.1"
/db_xref="GI:416115"

35 /translation="MSLEQRSIHCKPBEALEAQOEALGLVCVQATSSSSPL
VLGTL EEVPTAGSTDPQPQSGASAFPTTINFTRQRPSEGSSSREEEGPST
40 SCILES LFRAVITKKVADLVGFLL LKYRAREPVTKAEMLESVIKNYKHC FPE
IFGKASESLQLVFGIDVKEADPTGHSYVLVTCLGLSYDGLLDGNDQIMPKTGF
LIIVLVM IAMEGGHAPEEBIWEELSVMEVYDGREHSAYGEPRKLLTQDLVQE
KYLEYRQVPDSDPARYEF LWGPRLAETSYVKVLEYVIKVSARVRFFFP SLR
EAALREEEEGV"

45 polyA_signal 2095..2100
/gene="MAGE-1"

intron 2112..2420
/partial
gene="MAGE-1"
/number=3

50 BASE COUNT 562 a 582 c 677 g 599 t
ORIGIN
1 ggatccaggc cctgccagga aaaatataag ggccctgcgt gagaacagag
ggggtcaccc
55 61 actgcatgag agtggggatg tcacagagtc cagcccaccc tcttggttagc
actgagaagc
121 cagggctgtg cttgcggtct gcaccctgag ggcccgtgga ttcctcttcc
tgagagctcca

	181	ggaaccaggc	agtgaggcct	tggtctgaga	cagtatcctc	aggtcacaga
		gcagaggatg				
	241	cacaggggtgt	gccagcagtg	aatgtttgcc	ctgaatgcac	accaagggcc
5		ccacctgcca				
	301	caggacacat	aggactccac	agagtctggc	ctcacctccc	tactgtcagt
		cctgtagaat				
	361	cgacctctgc	tggcoggetg	taccctgagt	accctctcac	ttcctccttc
		aggttttcag				
10	421	gggacaggcc	aaccagagg	acaggattcc	ctggaggcca	cagaggagca
		ccaaggagaa				
	481	gatctgtaag	taggcctttg	ttagagtctc	caaggttcag	ttctcagctg
		aggcctctca				
	541	cacactccct	ctctccccag	gcctgtgggt	cttcattgcc	cagctcctgc
		ccacactcct				
15	601	gcctgctgcc	ctgacgagag	tcattcatgtc	tcttgagcag	aggagtctgc
		actgcaagcc				
	661	tgaggaagcc	cttgaggccc	aacaagaggc	cctgggcctg	gtgtgtgtgc
		aggctgccac				
20	721	ctcctcctcc	tctcctctgg	tcctgggcac	cctggaggag	gtgcccactg
		ctgggtcaac				
	781	agatcctccc	cagagtccctc	agggagccctc	cgcctttccc	actaccatca
		acttcactcg				
	841	acagaggcaa	cccagtgagg	gttccagcag	ccgtgaagag	gagggggcaa
		gcacctcttg				
25	901	tatcctggag	tccttggttc	gagcagtaat	cactaagaag	gtggctgatt
		tggttggttt				
	961	tctgctcctc	aaatatcgag	ccaggagacc	agtcacaaaag	gcagaaatgc
		tggagagtgt				
30	1021	catcaaaaaat	tacaagcact	gttttcctga	gatcttcggc	aaagcctctg
		agtccttgca				
	1081	gctggctcttt	ggcattgacg	tgaaggaagc	agaccccacc	ggccactcct
		atgtccttgt				
	1141	cacctgccta	ggtctctcct	atgatggcct	gctgggtgat	aatcagatca
		tgcccaagac				
35	1201	aggcttcctg	ataattgtcc	tggtcatgat	tgcaatggag	ggcggccatg
		ctcctgagga				
	1261	ggaaatctgg	gaggagctga	gtgtgatgga	ggtgtatgat	gggaggggagc
		acagtgccta				
40	1321	tggggagccc	aggaagctgc	tcaccaaga	tttgggtgcag	gaaaagtacc
		tggagtaaccg				
	1381	gcaggtgccg	gacagtgatc	ccgcacgcta	tgagttcctg	tgggggtccaa
		gggccctcgc				
	1441	tgaaaccagc	tatgtgaaag	tccttgagta	tgtgatcaag	gtcagtgcaa
		gagttcgctt				
45	1501	tttcttccca	tcctgcgtg	aagcagcttt	gagagaggag	gaagagggag
		tctgagcatg				
	1561	agttgcagcc	aaggccagtg	ggagggggac	tgggccagtg	caccttccag
		ggccgcgtcc				
50	1621	agcagcttcc	cctgcctcgt	gtgacatgag	gccattctt	cactctgaag
		agagcgggtca				
	1681	gtgttctcag	tagtaggttt	ctgttctatt	gggtgacttg	gagatttatc
		tttgttctct				
	1741	tttgaattg	ttcaaatgtt	tttttttaag	ggatgggtga	atgaacttca
		gcattccaagt				
55	1801	ttatgaatga	cagcagtcac	acagttctgt	gtatatagtt	taagggttaag
		agtcttgtgt				
	1861	tttattcaga	ttgggaaatc	cattctatct	tgtgaattgg	gataataaca
		gcagtgggaat				

```

1921 aagtacttag aaatgtgaaa aatgagcagt aaaatagatg agataaagaa
ctaaagaaat
1981 taagagatag tcaattcttg cttataacct cagtctattc tgtaaaattt
ttaaagatat
5 2041 atgcataacct ggatttcctt ggcttctttg agaatgtaag agaaattaaa
tctgaataaa
2101 gaattcttcc tgttcaactgg ctcttttctt ctccatgcac tgagcatctg
cttttttgaa
2161 ggccctgggt tagtagtgga gatgctaagg taagccagac tcatacccac
10 ccatagggtc
2221 gtagagtcta ggagctgcag tcacgtaatc gaggtggcaa gatgtcctct
aaagatgtag
2281 ggaaaagtga gagaggggtg aggggtgtggg gctccgggtg agagtgggtg
agtgtcaatg
15 2341 ccctgagctg gggcattttg ggctttggga aactgcagtt ctttctgggg
gagctgattg
2401 taatgatctt ggttgatcc
//

20 L18920. Human MAGE-2 gene...[gi:436180]
LOCUS HUMMAGE2X 4559 bp DNA PRI
20-APR-1994
DEFINITION Human MAGE-2 gene exons 1-4, complete cds.
ACCESSION L18920
25 VERSION L18920.1 GI:436180
KEYWORDS
SOURCE Homo sapiens (human).
ORGANISM Homo sapiens
Eukaryota; Metazoa; Chordata; Craniata;
30 Vertebrata; Euteleostomi;
Mammalia; Eutheria; Primates; Catarrhini;
Hominidae; Homo.
REFERENCE 1 (bases 1 to 4559)
AUTHORS De Smet,C., Lurquin,C., van der Bruggen,P., De
35 Plaen,E.,
Brasseur,F. and Boon,T.
TITLE Sequence and expression pattern of the human MAGE2 gene
JOURNAL Immunogenetics 39 (2), 121-129 (1994)
MEDLINE 94102805
40 FEATURES Location/Qualifiers
source 1..4559
/organism="Homo sapiens"
/isolate="MZ-2"
/db_xref="taxon:9606"
45 /map="X"
/haplotype="A1 A29 B37 B44 C6"
/sex="female"
/cell_line="MZ2-MEL.43"
/cell_type="melanoma cell line"
50 gene 561..4496
/gene="MAGE-2"
exon 561..754
/gene="MAGE-2"
/number=1
55 intron 755..1850
/gene="MAGE-2"
/number=1
exon 1851..1969

```

```

                    /gene="MAGE-2"
                    /number=2
intron            1970..2788
                    /gene="MAGE-2"
5                /number=2
exon              2789..2854
                    /gene="MAGE-2"
                    /number=3
10              intron      2855..2934
                    /gene="MAGE-2"
                    /number=3
exon              2935..4496
                    /gene="MAGE-2"
                    /number=4
15              CDS         3000..3944
                    /gene="MAGE-2"
                    /codon_start=1
                    /protein_id="AAA17729.1"
                    /db_xref="GI:436181"
20
    /translation="MPLEQRSQHCKPEEGLEARGEALGLVGAQAPATEEQQTASSSSTLVEVT
LGEVPAADSPSPPHSPQGASSFSTTINYTLWRQSDGSSNQEEEGPRMFPDLE
SEFQAAISRKMVELVHFLLLKYRAREPVTKAEMLESVLRNCQDFFPVIFSKASEYLQLVFGIE
VVEVVPISHLYILVTCLGLSYDGLLDNQVMPKTGLLIIVLAIIEGDCAPEEKIWEELSML
25 EVFEGREDSVFAHPRKLLMQDLVQBNYLEYRQVPGSDPACYEFLWGPRLIETSYVKVLHHTL
KIGGEPHISYPPLHERALREGE"
polyA_signal      4479..4484
                    /gene="MAGE-2"
BASE COUNT      1049 a   1280 c   1283 g   947 t
30 ORIGIN
      1 attccttcat caaacagcca ggagtgagga agaggaccct cctgagtgag
gactgaggat
      61 ccaccctcac cacatagtgg gaccacagaa tccagctcag cccctcttgt
cagccctggt
35      121 acacactggc aatgatctca ccccgagcac acccctcccc ccaatgccac
ttcgggccga
      181 ctcagagtca gagacttggg ctgagggggag cagacacaat cggcagagga
tggcggtcca
      241 ggctcagtct ggcattccaag tcaggacctt gagggatgac caaaggcccc
40      tccccacccc
      301 aactcccccg accccaccag gatctacagc ctcaggatcc ccgtcccaat
ccctaccctt
      361 acaccaacac catcttcatg cttaccccca cccccccatc cagatcccca
tccgggcaga
45      421 atccggttcc acccttgccg tgaaccaggg gaagtcacgg gcccggtgtg
gacgccactg
      481 acttgacatc tggaggtcag aggacagcga gattctcgcc ctgagcaacg
gcctgacgtc
      541 ggcggaggga agcaggcgca ggctccgtga ggaggcaagg taagacgccg
50      agggaggact
      601 gaggcggggc tcaccccaga cagagggccc ccaataatcc agcgctgcct
ctgctgccgg
      661 gcctggacca ccctgcaggg gaagacttct caggctcagt cgccaccacc
tcaccccgcc
55      721 accccccgcc gctttaaccg cagggaactc tggcgtaaga gctttgtgtg
accagggcag
      781 ggctgggttag aagtgtcag ggcccagact cagccaggaa tcaaggtcag
gacccaaga

```


	841	ggggactgag	ggcaaccac	cccctaccct	cactaccaat	cccatcccc
		aacaccaacc				
	901	ccacccccat	ccctcaaaca	ccaacccac	ccccaaacc	cattcccatc
		tcctcccca				
5	961	ccaccatcct	ggcagaatcc	ggctttgccc	ctgcaatcaa	cccacggaag
		ctccgggaat				
	1021	ggcgccaag	cacgcggatc	ctgacgttca	catgtacggc	taagggaggg
		aaggggttg				
	1081	gtctcgtgag	tatggccttt	gggatgcaga	ggaagggccc	aggcctcctg
10		gaagacagt				
	1141	gagtccttag	gggaccagc	atgccaggac	agggggccca	ctgtaccctt
		gtctcaaact				
	1201	gagccacctt	ttcattcagc	cgagggaatc	ctagggatgc	agaccactt
		cagcaggggg				
15	1261	ttggggccca	gcctgcgagg	agtcaagggg	aggaagaaga	gggaggactg
		aggggacctt				
	1321	ggagtccaga	tcagtggcaa	ccttgggctg	ggggatcctg	ggcacagtgg
		ccgaatgtgc				
	1381	cccgtgctca	ttgcaccttc	agggtgacag	agagttgagg	gctgtggtct
20		gagggctggg				
	1441	acttcaggtc	agcagagggg	ggaatcccag	gatctgccgg	accaaggtg
		tgcccccttc				
	1501	atgaggactg	gggatacccc	cggcccagaa	agaagggatg	ccacagagtc
		tgggaagtccc				
25	1561	ttgttcttag	ctctggggga	acctgatcag	ggatggccct	aagtgacaat
		ctcatttgta				
	1621	ccacaggcag	gaggttgggg	aaccctcagg	gagataaggt	gttgggtgtaa
		agaggagctg				
	1681	tctgctcatt	tcagggggtt	gggggttgag	aaagggcagt	ccctggcagg
30		agtaaagatg				
	1741	agtaaccac	aggaggccat	cataacgttc	accctagaac	caaaggggtc
		agccctggac				
	1801	aacgcacgtg	ggggtaacag	gatgtggccc	ctcctcactt	gtctttccag
		atctcagggg				
35	1861	gttgatgacc	ttgttttcag	aagggtgactc	aggtcaacac	aggggcccc
		tctggtcgac				
	1921	agatgcagtg	gttctaggat	ctgccaagca	tccagggtga	gagcctgagg
		taggattgag				
	1981	ggtacccctg	ggccagaatg	cagcaagggg	gccccataga	aatctgccct
40		gcccctgcgg				
	2041	ttacttcaga	gaccctgggc	agggtgtgca	gctgaagtcc	ctccattatc
		ctgggatctt				
	2101	tgatgtcagg	gaaggggagg	ccttgggtctg	aaggggctgg	agtcaggtca
		gtagagggag				
45	2161	ggtctcaggc	cctgccagga	gtggacgtga	ggaccaagcg	gactcgtcac
		ccaggacacc				
	2221	tggactccaa	tgaatttgga	catctctcgt	tgtccttcgc	gggaggacct
		ggtcacgtat				
	2281	ggccagatgt	gggtcccttc	atatccttct	gtaccatata	agggatgtga
50		gttcttgaca				
	2341	tgagagattc	tcaagccagc	aaaaggggtg	gattaggccc	tacaaggaga
		aaggtgaggg				
	2401	ccctgagtga	gcacagaggg	gaccctccac	ccaagtagag	tggggacctc
		acggagtctg				
55	2461	gccaaacctg	ctgagacttc	tgggaatccg	tggctgtgct	tgaggtctgc
		acactgaagg				
	2521	cccgtgcatt	cctctcccag	gaatcaggag	ctccaggaac	caggcagtga
		ggccttggtc				

	2581	tgagtcagt	tcctcaggtc	acagagcaga	ggggacgcag	acagtgccaa
		cactgaagg				
	2641	ttgcctggaa	tgcacaccaa	gggccccacc	cgcccagaac	aaatgggact
5		ccagagggcc				
	2701	tggcctcacc	ctccctattc	tcagtcctgc	agcctgagca	tgtgctggcc
		ggctgtaccc				
	2761	tgaggtgccc	tcccacttcc	tccttcaggt	tctgaggggg	acaggctgac
		aagtaggacc				
10	2821	cgaggcactg	gaggagcatt	gaaggagaag	atctgtaagt	aagcctttgt
		cagagcctcc				
	2881	aaggttcagt	tcagttctca	cctaaggcct	cacacacgct	ccttctctcc
		ccaggcctgt				
	2941	gggtcttcat	tgcccagctc	ctgcccgcac	tcctgcctgc	tgccctgacc
		agagtcacca				
15	3001	tgctcttga	gcagaggagt	cagcactgca	agcctgaaga	aggccttgag
		gcccgaggag				
	3061	aggccctggg	cctgggtggg	gcgcaggctc	ctgctactga	ggagcagcag
		accgcttctt				
20	3121	cctcttctac	tctagtggaa	gttaccctgg	gggaggtgcc	tgctgccgac
		tcaccgagtc				
	3181	ctccccacag	tcctcagggg	gcctccagct	tctcgactac	catcaactac
		actctttgga				
	3241	gacaatccga	tgagggctcc	agcaaccaag	aagaggaggg	gccagaatg
25		tttcccgacc				
	3301	tgaggtccga	gttccaagca	gcaatcagta	ggaagatggg	tgagttgggt
		cattttctgc				
	3361	tcctcaagta	tcgagccagg	gagccggtca	caaaggcaga	aatgctggag
		agtgtcctca				
30	3421	gaaattgcc	ggacttcttt	cccgtgatct	tcagcaaagc	ctccgagtac
		ttgcagctgg				
	3481	tctttggcat	cgaggtgggt	gaagtgggtc	ccatcagcca	cttgtagatc
		cttgtcacct				
	3541	gcctgggcct	ctcctacgat	ggcctgctgg	gcgacaatca	ggcatgccc
		aagacaggcc				
35	3601	tcctgataat	cgtcctggcc	ataatcgcaa	tagagggcga	ctgtgcccct
		gaggagaaaa				
	3661	tctgggagga	gctgagtatg	ttggaggtgt	ttgaggggag	ggaggacagt
		gtcttcgcac				
40	3721	atcccaggaa	gctgctcatg	caagatctgg	tgaggaaaa	ctacctggag
		taccggcagg				
	3781	tgcccggcag	tgatcctgca	tgctacgagt	tcctgtgggg	tccaagggcc
		ctcattgaaa				
	3841	ccagctatgt	gaaagtcctg	caccatacac	taaagatcgg	tgaggaaacct
45		cacatttcct				
	3901	acccacccct	gcatgaacgg	gctttgagag	agggagaaga	gtgagtctca
		gcacatgttg				
	3961	cagccagggc	cagtgggagg	gggtctgggc	cagtgcacct	tccagggccc
		catccattag				
50	4021	cttccactgc	ctcgtgtgat	atgaggccca	ttcctgcctc	tttgaagaga
		gcagtcagca				
	4081	ttcttagcag	tgagtttctg	ttctgttgga	tgactttgag	atttatcttt
		ctttcctgtt				
	4141	ggaattgttc	aaatgttcct	tttaacaaat	ggttgatga	acttcagcat
55		ccaagtttat				
	4201	gaatgacagt	agtcacacat	agtgtgtttt	atatagttta	ggggtaagag
		tcctgttttt				
	4261	tattcagatt	gggaaatcca	ttccattttg	tgagttgtca	cataataaca
		gcagtgggaat				

```

      4321 atgtatttgc ctatattgtg aacgaattag cagtaaaata catgatacaa
ggaactcaaa
      4381 agatagttaa ttcttgccctt atacctcagt ctattatgta aaattaaaaa
tatgtgtatg
5      4441 tttttgcttc tttgagaatg caaaagaaat taaatctgaa taaattcttc
ctgttcactg
      4501 gctcatttct ttaccattca ctcagcatct gctctgtgga aggccttggg
agtagtggg
//
10
U03735. Human MAGE-3 anti...[gi:468825]
LOCUS      HSU03735      4204 bp      DNA      PRI
07-APR-1994
DEFINITION Human MAGE-3 antigen (MAGE-3) gene, complete cds.
15  ACCESSION  U03735
VERSION     U03735.1  GI:468825
KEYWORDS    .
SOURCE      human.
ORGANISM    Homo sapiens
20          Eukaryota;      Metazoa;      Chordata;      Craniata;
Vertebrata; Euteleostomi;
          Mammalia;      Eutheria;      Primates;      Catarrhini;
Hominidae; Homo.
REFERENCE   1  (bases 1 to 4204)
25  AUTHORS   Gaugler,B., Van den Eynde,B., van der Bruggen,P.,
Romero,P.,
          Gaforio,J.J., De Plaen,E., Lethe,B., Brasseur,F.
and Boon,T.
TITLE       Human gene MAGE-3 codes for an antigen recognized on
30  a melanoma by
          autologous cytolytic T lymphocytes
JOURNAL     J. Exp. Med. 179, 921-930 (1994)
MEDLINE     94157413
REFERENCE   2  (bases 1 to 4204)
35  AUTHORS   Gaugler,B.
TITLE       Direct Submission
JOURNAL     Submitted (25-NOV-1993) Beatrice Gaugler, Ludwig
Institute for
          Cancer Research, 74 Avenue Hippocrate, Brussels
40  1200, Belgium
FEATURES    Location/Qualifiers
          source      1..4204
                        /organism="Homo sapiens"
                        /isolate="patient MZ2"
45          /db_xref="taxon:9606"
                        /chromosome="X"
                        /sex="female"
                        /cell_type="lymphocyte"
                        /tissue_type="blood"
50          /dev_stage="adult"
          exon        434..468
                        /number=1
          exon        2254..2319
                        /number=2
55          exon        2400..3978
                        /number=3
          gene        2465..3409
                        /gene="MAGE-3"

```

CDS 2465..3409
 /gene="MAGE-3"
 /codon_start=1
 /product="MAGE-3 antigen"
 /protein_id="AAA17446.1"
 /db_xref="GI:468826"
 /translation="MPLEQRSQHCKPEEGLEARGEALGLVGAQAPATEEQEAASSSSTLVEVTLGE
 VPAESPDPQSPQGASSLPTTMNYPLWSQSYEDSSNQEEEGPSTFPDLESEFQAALSRKVAELVH
 FLLLKYRAREPVTKAEMLGSVVGNWQYFFPVIFSKASSSLQLVFGIELMEVDPIGHLIYIFATCLGL
 SYDGLLDNQIMPKAGLLIIVLAI IAREGDCAPEEKIWEELSVLEVFEGRSDSILGDPKKLLTQHF
 VQENYLBRYQVPGSDPACYEFLWGPRLVETSYVKVLHMHMVKISGGPHISYPPLHEWVLRGEE"
 polyA_signal 3958..3963
 polyA_site 3979
 15 BASE COUNT 944 a 1144 c 1223 g 893 t
 ORIGIN
 1 acgcaggcag tgatgtcacc cagaccacac cccttcccc aatgccactt
 caggggggtac
 20 61 tcagagtcag agacttggtc tgaggggagc agaagcaatc tgcagaggat
 ggcgggtccag
 121 gctcagccag gcatcaactt caggaccctg agggatgacc gaaggccccc
 cccacccacc
 181 cccaactccc ccgacccccc caggatctac agcctcagga cccccgtccc
 aatccttacc
 25 241 ccttgcccca tcaccatctt catgcttacc tccaccccca tccgatcccc
 atccaggcag
 301 aatccagttc caccctgcc cggaaccag ggtagtaccg ttgccaggat
 gtgacgccac
 361 tgacttgccg attggaggtc agaagaccgc gagattctcg ccctgagcaa
 cgagcgacgg
 421 cctgacgtcg gcggagggaa gccggcccag gctcggtagg gaggcaaggt
 aagacgctga
 481 gggaggactg aggcgggcct cacctcagac agagggcctc aaataatcca
 gtgctgcctc
 541 tgctgccggg cctgggccac cccgcagggg aagacttcca ggctgggtcg
 ccactacctc
 601 accccgcega ccccccgcgc ttagccacg gggaaactctg gggacagagc
 ttaatgtggc
 661 cagggcaggg ctggttagaa gaggtcaggg cccacgctgt ggcaggaatc
 aaggtcagga
 721 ccccgagagg gaactgaggg cagcctaacc accaccctca ccaccattcc
 cgtcccccaa
 781 caccacaacc cacccccac cccattccc atccccacc ccaccctat
 cctggcagaa
 841 tccgggcttt gccctggta tcaagtcacg gaagctccgg gaatggcggc
 caggcacgtg
 901 agtcctgagg ttcacatcta cggctaaggg agggaagggg ttcggtatcg
 cgagtatggc
 961 cgttgggagg cagcgaaagg gcccaggcct cctggaagac agtggagtcc
 tgaggggacc
 1021 cagcatgcca ggacaggggg cccactgtac ccctgtctca aaccgaggca
 ccttttcatt
 1081 cggctacggg aatcctaggg atgcagaccc acttcagcag ggggttgggg
 cccagccctg
 1141 cgaggagtca tggggaggaa gaagagggag gactgagggg accttgaggt
 ccagatcagt
 1201 ggcaaccttg ggctggggga tgctgggcac agtggccaaa tgtgctctgt
 gctcattgcg

	1261	ccttcagggg	gaccagagag	ttgagggctg	tggtctgaag	agtgggactt
		caggtcagca				
	1321	gagggaggaa	tcccaggatc	tgcagggccc	aaggtgtacc	ccaagggggc
5		ccctatgtgg				
	1381	tggacagatg	cagtggctct	aggatctgcc	aagcatccag	gtgaagagac
		tgagggagga				
	1441	ttgagggtag	ccctgggaca	gaatgcggac	tgggggcccc	ataaaaatct
		gccctgctcc				
10	1501	tgctgttacc	tcaagagacc	tgggcagggc	tgctcagctga	ggtccctcca
		ttatcctagg				
	1561	atcactgatg	tcaggggaagg	ggaagccttg	gtctgagggg	gctgcactca
		gggcagtaga				
	1621	gggaggctct	cagaccctac	taggagtggg	ggtgaggacc	aagcagtctc
		ctcaccaggg				
15	1681	gtacatggac	ttcaataaat	ttggacatct	ctcggtgtcc	tttccggggg
		gacctgggaa				
	1741	tgtatggcca	gatgtgggtc	ccctcatgtt	tttctgtacc	atatcaggta
		tgtgagttct				
20	1801	tgacatgaga	gattctcagg	ccagcagaag	ggagggatta	ggccctataa
		ggagaaaggt				
	1861	gagggccctg	agtgagcaca	gaggggatcc	tccaccccag	tagagtgggg
		acctcacaga				
	1921	gtctggccaa	ccctcctgac	agttctggga	atccgtggct	gcgtttgctg
		tctgcacatt				
25	1981	gggggcccgt	ggattcctct	cccaggaatc	aggagctcca	ggaacaaggc
		agtgaggact				
	2041	tggtctgagg	cagtgtcctc	aggtcacaga	gtagaggggg	ctcagatagt
		gccaacgggtg				
30	2101	aaggtttgcc	ttggattcaa	accaagggcc	ccacctgccc	cagaacacat
		ggactccaga				
	2161	gcgcctggcc	tcacctcaa	tactttcagt	cctgcagcct	cagcatgcgc
		tggccggatg				
	2221	tacctgagg	tgccctotca	cttcctcctt	caggtttctga	ggggacaggc
		tgacctggag				
35	2281	gaccagaggc	ccccggagga	gactgaagg	agaagatctg	taagtaagcc
		tttggttagag				
	2341	cctccaaggt	tccattcagt	actcagctga	ggtctctcac	atgctccctc
		tctccccagg				
40	2401	ccagtgggtc	tccattgccc	agctcctgcc	cacactcccg	cctggtgccc
		tgaccagagt				
	2461	catcatgcct	cttgagcaga	ggagtcagca	ctgcaagcct	gaagaaggcc
		ttgaggcccg				
	2521	aggagaggcc	ctgggcctgg	tgggtgcgca	ggctcctgct	actgaggagc
		aggaggctgc				
45	2581	ctcctcctct	tctactctag	ttgaagtcac	cctggggggag	gtgcctgctg
		ccgagtcacc				
	2641	agatcctccc	cagagtcctc	agggagcctc	cagcctcccc	actaccatga
		actaccctct				
50	2701	ctggagccaa	tcctatgagg	actccagcaa	ccaagaagag	gagggggccaa
		gcaccttccc				
	2761	tgacctggag	tccgagttcc	aagcagcact	cagtaggaag	gtggccgagt
		tggttcattt				
	2821	tctgctcctc	aagtatcgag	ccaggagacc	ggtcacaaag	gcagaaatgc
		tggggagtggt				
55	2881	cgtcggaaat	tggcagtatt	tctttcctgt	gatcttcagc	aaagcttcca
		gttccttgca				
	2941	gctgggtcttt	ggcatcgagc	tgatggaagt	ggaccccatc	ggccacttgt
		acatctttgc				

```

      3001 cacctgcctg ggcctctcct acgatggcct gctgggtgac aatcagatca
tgcccaaggc
      3061 aggcctcctg ataatcgctc tggccataat cgcaagagag ggcgactgtg
ccccctgagga
5      3121 gaaaatctgg gaggagctga gtgtgttaga ggtgtttgag gggaggggaag
acagtatctt
      3181 gggggatccc aagaagctgc tcaccaaca tttcgtgcag gaaaactacc
tgaggtaccg
      3241 gcaggtcccc ggcagtgatc ctgcatgtta tgaattcctg tgggggccaa
10      gggccctcgt
      3301 tgaaaccagc tatgtgaaag tcctgcacca tatggtaaag atcagtggag
gacctcacat
      3361 ttcctaccca cccctgcatg agtggggttt gagagagggg gaagagttag
tctgagcacg
15      3421 agttgcagcc agggccagtg ggaggggggtc tggggccagt cactttccgg
ggccgcatcc
      3481 cttagtttcc actgcctcct gtgacgtgag gccattctt cactctttga
agcgagcagt
      3541 cagcattctt agtagtgggt ttctgttctg ttggatgact ttgagattat
20      tctttgtttc
      3601 ctgttggagt tgttcaaagt ttcttttaa cggatggtt aatgagcgtc
agcatccagg
      3661 tttatgaatg acagtagtca cacatagtgc tgtttatata gtttaggagt
aagagtcttg
25      3721 ttttttactc aaattgggaa atccattcca ttttgtgaat tgtgacataa
taatagcagt
      3781 ggtaaaagta tttgcttaaa attgtgagcg aattagcaat aacatacatg
agataactca
      3841 agaaatcaaa agatagttga ttcttgccct gtacctcaat ctattctgta
30      aaattaaaca
      3901 aatatgcaaa ccaggatttc cttgacttct ttgagaatgc aagcgaaatt
aaatctgaat
      3961 aaataattct tcctcttcac tggctcgttt cttttccgtt cactcagcat
ctgctctgtg
35      4021 ggaggccctg ggtagtagt ggggatgcta aggtaagcca gactcacgcc
taccatagg
      4081 gctgtagagc ctaggacctg cagtcataata attaggtgg tgagaagtcc
tgtaagatgt
      4141 agaggaaatg taagagaggg gtgagggtgt ggcgctccgg gtgagagtag
40      tggagtgtca
      4201 gtgc
//

```

```

45      AF043498. Homo sapiens pros...[gi:2909843]
      LOCUS      AF043498      990 bp      mRNA      PRI      24-
      FEB-1998
      DEFINITION  Homo sapiens prostate stem cell antigen (PSCA)
      mRNA, complete cds.
      ACCESSION  AF043498
50      VERSION    AF043498.1 GI:2909843
      KEYWORDS    .
      SOURCE      human.
      ORGANISM    Homo sapiens
85      Euteleostomi;
      Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
      Homo.
      REFERENCE   1 (bases 1 to 990)

```

AUTHORS Reiter,R.E., Gu,Z., Watabe,T., Thomas,G., Kinga,S.,
 Davis,E., Wahl,M., Nisitani,S., Yamashiro,J., Le Beau,M.M.,
 5 Losa,M. and Witte,O.N.
 TITLE Prostate stem cell antigen: a cell surface marker
 overexpressed in
 prostate cancer
 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 95 (4), 1735-1740 (1998)
 10 MEDLINE 98132661
 REFERENCE 2 (bases 1 to 990)
 AUTHORS Reiter,R.E.
 TITLE Direct Submission
 JOURNAL Submitted (19-JAN-1998) Urology, UCLA, 66-134 CHS
 15 10833 Le Conte
 Ave., Los Angeles, CA 90095, USA
 FEATURES Location/Qualifiers
 source 1..990
 20 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /chromosome="8"
 /map="8q24.2"
 /note="LAPC-4 prostate cancer xenograft"
 gene 1..990
 25 /gene="PSCA"
 CDS 18..389
 /gene="PSCA"
 /note="GPI-anchored cell surface protein"
 /codon_start=1
 30 /product="prostate stem cell antigen"
 /protein_id="AAC39607.1"
 /db_xref="GI:2909844"
 /translation="MKAVLLALLMAGLALQPGTALLCYSCKAQVSNEDCLQVENCTQLGEQCW
 35 TARIRAVGLLTVISKGCSLNCVDDSDYYVVGKKNITCCDTDLNAGAHALQPAAAILALLPA
 LGLLLWGPGQL"
 BASE COUNT 193 a 299 c 285 g 202 t 11 others
 ORIGIN
 1 agggagagggc agtgaccatg aaggctgtgc tgcttgccct gttgatggca
 40 ggcttggccc
 61 tgcagccagg cactgccctg ctgtgtact cctgcaaagc ccagggtgagc
 aacgaggact
 121 gcctgcaggt ggagaactgc acccagctgg gggagcagtg ctggaccgcg
 45 cgcacccgcg
 181 cagttggcct cctgaccgtc atcagcaaag gctgcagctt gaactgcgtg
 gatgactcac
 241 aggactacta cgtgggcaag aagaacatca cgtgctgtga caccgacttg
 tgcaacgcca
 301 gcgggggcca tgccctgcag cgggctgccg ccatccttgc gctgctccct
 50 gcaactcgcc
 361 tgctgctctg gggacccggc cagctatagg ctctgggggg ccccgctgca
 gccacactg
 421 ggtgtggtgc cccaggcctt tgtgccactc ctcacagaac ctggcccagt
 gggagcctgt
 55 481 cctgggttct gaggcacatc ctaacgcaag tttgaccatg tatgtttgca
 ccccttttcc
 541 ccnaaccctg accttcccat gggccttttc caggattccn accnggcaga
 tcagtttttag

```

        601 tganacanat ccgcntgcag atggcccctc caaccntttt tgttgntggt
tccatggccc
        661 agcattttcc acccttaacc ctgtgttcag gcacttnttc cccaggaag
ccttccctgc
5       721 ccacccatt tatgaattga gccaggtttg gtccgtggtg tccccgcac
ccagcagggg
        781 acaggcaatc aggagggcc agtaaaggct gagatgaagt ggactgagta
gaactggagg
10      841 acaagagttg acgtgagttc ctgggagttt ccagagatgg ggcctggagg
cctggaggaa
        901 ggggccaggc ctcacatttg tgggntccc gaatggcagc ctgagcacag
cgtaggccct
        961 taataaacac ctgttgata agccaaaaaa
//
15
P06870. GLANDULAR KALLIKR...[gi:125170]
LOCUS      KLK1_HUMAN      262 aa                      PRI      20-
AUG-2001
DEFINITION      GLANDULAR KALLIKREIN 1 PRECURSOR (TISSUE
20 KALLIKREIN)
(KIDNEY/PANCREAS/SALIVARY GLAND KALLIKREIN).
ACCESSION      P06870
PID            gi125170
VERSION        P06870 GI:125170
25 DBSOURCE      swissprot: locus KLK1_HUMAN, accession P06870;
class: standard.
extra accessions: Q9UMJ1, created: Jan 1, 1988.
sequence updated: Jan 1, 1988.
annotation updated: Aug 20, 2001.
30 xrefs: gi: gi: 186652, gi: gi: 186653, gi: gi: 186649,
gi: gi:
186651, gi: gi: 186645, gi: gi: 186646, gi: gi:
186647, gi: gi:
186648, gi: gi: 34026, gi: gi: 34027, gi: gi: 186643,
35 gi: gi:
386843, gi: gi: 67558
xrefs (non-sequence databases): HSSP P00757, MEROPS
S01.160,
GlycoSuiteDB P06870, MIM 147910, InterPro IPR001314,
40 InterPro
IPR001254, Pfam PF00089, PRINTS PR00722, PROSITE
PS50240, PROSITE
PS00134, PROSITE PS00135
KEYWORDS      Hydrolase; Serine protease; Glycoprotein; Multigene
45 family;
Zymogen; Signal.
SOURCE      human.
ORGANISM      Homo sapiens
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
50 Euteleostomi;
Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
Homo.
REFERENCE      1 (residues 1 to 262)
AUTHORS      Fukushima,D., Kitamura,N. and Nakanishi,S.
55 TITLE      Nucleotide sequence of cloned cDNA for human
pancreatic kallikrein
JOURNAL      Biochemistry 24, 8037-8043 (1985)
REMARK      SEQUENCE FROM N.A.

```


TISSUE=Pancreas
 REFERENCE 2 (residues 1 to 262)
 AUTHORS Evans,B.A., Yun,Z.X., Close,J.A., Tregear,G.W.,
 Kitamura,N.,
 5 Nakanishi,S., Callen,D.F., Baker,E., Hyland,V.J.,
 Sutherland,G.R.
 and Richards,R.I.
 TITLE Structure and chromosomal localization of the human
 renal
 10 kallikrein gene
 JOURNAL Biochemistry. 27 (9), 3124-3129 (1988)
 MEDLINE 88269498
 PUBMED 2898948
 REMARK SEQUENCE FROM N.A.
 15 TISSUE=Kidney
 REFERENCE 3 (residues 1 to 262)
 AUTHORS Angermann,A., Bergmann,C. and Appelhans,H.
 TITLE Cloning and expression of human salivary-gland
 kallikrein in
 20 Escherichia coli
 JOURNAL The Biochemical journal. 262 (3), 787-793 (1989)
 MEDLINE 90073574
 PUBMED 2686621
 REMARK SEQUENCE FROM N.A.
 25 TISSUE=Salivary gland
 REFERENCE 4 (residues 1 to 262)
 AUTHORS Baker,A.R. and Shine,J.
 TITLE Human kidney kallikrein: cDNA cloning and sequence
 analysis
 30 JOURNAL DNA (Mary Ann Liebert, Inc.) 4 (6), 445-450 (1985)
 MEDLINE 86135264
 PUBMED 3853975
 REMARK SEQUENCE OF 17-262 FROM N.A.
 TISSUE=Kidney
 35 REFERENCE 5 (residues 1 to 262)
 AUTHORS Lu,H.S., Lin,F.K., Chao,L. and Chao,J.
 TITLE Human urinary kallikrein. Complete amino acid sequence
 and sites of
 40 glycosylation
 JOURNAL International journal of peptide and protein research.
 33 (4),
 237-249 (1989)
 MEDLINE 89326688
 PUBMED 2666327
 45 REMARK SEQUENCE OF 25-262.
 TISSUE=Urine
 REFERENCE 6 (residues 1 to 262)
 AUTHORS Kellermann,J., Lottspeich,F., Geiger,R. and
 Deutzmann,R.
 50 TITLE Human urinary kallikrein--amino acid sequence and
 carbohydrate
 attachment sites
 JOURNAL Protein sequences & data analysis. 1 (3), 177-182
 (1988)
 55 MEDLINE 88203586
 PUBMED 3163150
 REMARK SEQUENCE OF 25-262, AND CARBOHYDRATE-LINKAGE SITES.
 TISSUE=Urine

REFERENCE 7 (residues 1 to 262)
 AUTHORS Lottspeich,F., Geiger,R., Henschen,A. and Kutzbach,C.
 TITLE N-Terminal amino acid sequence of human urinary
 kallikrein homology
 5 with other serine proteases
 JOURNAL Hoppe-Seyler's Zeitschrift fur physiologische Chemie.
 360 (12),
 1947-1950 (1979)
 MEDLINE 80114126
 10 PUBMED 393608
 REMARK SEQUENCE OF 25-55.
 TISSUE=Urine
 REFERENCE 8 (residues 1 to 262)
 AUTHORS Takahashi,S., Irie,A., Katayama,Y., Ito,K. and
 15 Miyake,Y.
 TITLE N-terminal amino acid sequence of human urinary
 prokallikrein
 JOURNAL Journal of biochemistry. 99 (3), 989-992 (1986)
 MEDLINE 86223893
 20 PUBMED 3635530
 REMARK SEQUENCE OF 28-47.
 TISSUE=Urine
 [FUNCTION] GLANDULAR KALLIKREINS CLEAVE MET-LYS AND
 25 ARG-SER BONDS
 IN KININOGEN TO RELEASE LYS-BRADYKININ.
 [CATALYTIC ACTIVITY] PREFERENTIAL CLEAVAGE OF ARG-|-
 XAA BONDS IN
 SMALL MOLECULE SUBSTRATES. HIGHLY SELECTIVE ACTION TO
 30 RELEASE
 KALLIDIN (LYSYL-BRADYKININ) FROM KININOGEN INVOLVES
 HYDROLYSIS OF
 MET-|-XAA OR LEU-|-XAA.
 [SIMILARITY] BELONGS TO PEPTIDASE FAMILY S1; ALSO
 35 KNOWN AS THE
 TRYPSIN FAMILY. KALLIKREIN SUBFAMILY.
 FEATURES Location/Qualifiers
 source 1..262
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 40 Protein 1..262
 /product="GLANDULAR KALLIKREIN 1 PRECURSOR"
 /EC_number="3.4.21.35"
 Region 1..18
 45 /region_name="Signal"
 /note="PROBABLE."
 Region 19..24
 /region_name="Propeptide"
 /note="ACTIVATION PEPTIDE (PROBABLE)."
 50 Region 25..262
 /region_name="Mature chain"
 /note="GLANDULAR KALLIKREIN 1."
 Bond bond(31,174)
 /bond_type="disulfide"
 55 Bond bond(50,66)
 /bond_type="disulfide"
 /note="BY SIMILARITY."

```

Site      65
           /site_type="active"
           /note="CHARGE RELAY SYSTEM."
5  Site      93
           /site_type="glycosylation"
           /note="O-LINKED."
Site      102
           /site_type="glycosylation"
           /note="N-LINKED (GLCNAC...)."
10 Site      104
           /site_type="glycosylation"
           /note="O-LINKED."
Site      108
           /site_type="glycosylation"
           /note="N-LINKED (GLCNAC...)."
15 Site      120
           /site_type="active"
           /note="CHARGE RELAY SYSTEM."
Region    145
           /region_name="Variant"
           /note="Q -> E. /FTId=VAR_006625."
Bond      bond(153,220)
           /bond_type="disulfide"
           /note="BY SIMILARITY."
25 Site      165
           /site_type="glycosylation"
           /note="N-LINKED (GLCNAC...) (PARTIAL)."
Site      167
           /site_type="glycosylation"
           /note="O-LINKED."
30 Bond      bond(185,199)
           /bond_type="disulfide"
           /note="BY SIMILARITY."
Region    186
           /region_name="Variant"
           /note="E -> K. /FTId=VAR_006626."
35 Bond      bond(210,235)
           /bond_type="disulfide"
           /note="BY SIMILARITY."
40 Site      214
           /site_type="active"
           /note="CHARGE RELAY SYSTEM."

ORIGIN
45 1 mwflvlclal slggtgaapp iqsrivggwe ceqhsqpwwa alyhfstfqc
   ggilvhrqvv
   61 ltaahcisdn yqlwlgrhnl fddentaqfv hvsesfphpg fnmsllenht
   rqadedyshd
   121 lmlrltepa dtitdavkvv elptqepevg stclasgwgs iepenfsfpd
   dlqcvdlkil
50 181 pndecekahv qkvtdfmlcv ghleggkdtc vgdsggplmc dgvlqgvtsw
   gyvpcgtpnk
   241 psvavrvlsy vkwiedtiaa ns
   //

55 P08217. ELASTASE 2A PRECU...[gi:119255]
   LOCUS      EL2A_HUMAN      269 aa      PRI      20-
   AUG-2001

```

DEFINITION ELASTASE 2A PRECURSOR.
 ACCESSION P08217
 PID g119255
 VERSION P08217 GI:119255
 DBSOURCE swissprot: locus EL2A_HUMAN, accession P08217;
 class: standard.
 created: Aug 1, 1988.
 sequence updated: Aug 1, 1988.
 annotation updated: Aug 20, 2001.
 xrefs: gi: gi: 182022, gi: gi: 182023, gi: gi: 182057,
 gi: gi: 182058, gi: gi: 88298, gi: gi: 88299
 xrefs (non-sequence databases): MEROPS S01.155,
 InterPro IPR001314,
 InterPro IPR001254, Pfam PF00089, PRINTS PR00722,
 PROSITE PS50240,
 PROSITE PS00134, PROSITE PS00135
 KEYWORDS Hydrolase; Serine protease; Pancreas; Zymogen; Signal.
 SOURCE human.
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
 Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
 Homo.
 REFERENCE 1 (residues 1 to 269)
 AUTHORS Kawashima,I., Tani,T., Shimoda,K. and Takiguchi,Y.
 TITLE Characterization of pancreatic elastase II cDNAs: two
 elastase II mRNAs are expressed in human pancreas
 JOURNAL DNA 6 (2), 163-172 (1987)
 MEDLINE 87217962
 REMARK SEQUENCE FROM N.A.
 REFERENCE 2 (residues 1 to 269)
 AUTHORS Fletcher,T.S., Shen,W.F. and Largman,C.
 TITLE Primary structure of human pancreatic elastase 2
 determined by
 sequence analysis of the cloned mRNA
 JOURNAL Biochemistry 26 (23), 7256-7261 (1987)
 MEDLINE 88107669
 REMARK SEQUENCE FROM N.A.
 [FUNCTION] ACTS UPON ELASTIN.
 [CATALYTIC ACTIVITY] PREFERENTIAL CLEAVAGE: LEU-|-XAA,
 MET-|-XAA
 AND PHE-|-XAA. HYDROLYSES ELASTIN.
 [SUBCELLULAR LOCATION] SECRETED.
 [TISSUE SPECIFICITY] PANCREAS.
 [SIMILARITY] BELONGS TO PEPTIDASE FAMILY S1; ALSO
 KNOWN AS THE TRYPSIN FAMILY. ELASTASE SUBFAMILY.
 FEATURES Location/Qualifiers
 source 1..269
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 Region 1..16
 /region_name="Signal"
 Protein 1..269
 /product="ELASTASE 2A PRECURSOR"
 /EC_number="3.4.21.71"

```

      Region      17..28
                   /region_name="Propeptide"
                   /note="ACTIVATION PEPTIDE."
5      Region      29..269
                   /region_name="Mature chain"
                   /note="ELASTASE 2A."
      Bond        bond(58,74)
                   /bond_type="disulfide"
                   /note="BY SIMILARITY."
10     Site        73
                   /site_type="active"
                   /note="CHARGE RELAY SYSTEM (BY SIMILARITY)."
```

```

      Site        121
                   /site_type="active"
                   /note="CHARGE RELAY SYSTEM (BY SIMILARITY)."
```

```

15     Bond        bond(155,222)
                   /bond_type="disulfide"
                   /note="BY SIMILARITY."
      Bond        bond(186,202)
20     Bond        /bond_type="disulfide"
                   /note="BY SIMILARITY."
      Bond        bond(212,243)
                   /bond_type="disulfide"
                   /note="BY SIMILARITY."
25     Site        216
                   /site_type="active"
                   /note="CHARGE RELAY SYSTEM (BY SIMILARITY)."
```

```

ORIGIN
30     1 mirtlllstl vagalscgrp typppyvtrvv ggeearpnsw pwqvsllqyss
      ngkwyhtcgg
      61 slianswvlt aahcissrt yrvglgrhnl yvaesgslav svskivvhkd
      wnsnqiskgn
      121 diallklanp vsltldkiqla clppagtilp nnypcyvtgw grlqtngavp
      dvlqqgrllv
35     181 vdyatcssa wwgssvktsm icaggdgvis scngdsggpl ncqasdgrwq
      vhgivsfgr
      241 lgcnyyhkps vftrvsnyid winsviann
      //
```

```

40     NP_056933. pancreatic elasta...[gi:7705648]
      LOCUS      NP_056933      269 aa      PRI      02-
      NOV-2000
      DEFINITION pancreatic elastase IIB [Homo sapiens].
      ACCESSION  NP_056933
45     PID      g7705648
      VERSION    NP_056933.1 GI:7705648
      DBSOURCE   REFSEQ: accession NM_015849.1
      KEYWORDS   .
      SOURCE     human.
50     ORGANISM Homo sapiens
                   Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
      Euteleostomi;
                   Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
      Homo.
55     REFERENCE 1 (residues 1 to 269)
      AUTHORS    Kawashima,I., Tani,T., Shimoda,K. and Takiguchi,Y.
      TITLE      Characterization of pancreatic elastase II cDNAs: two
      elastase II
```

mRNAs are expressed in human pancreas

JOURNAL DNA 6 (2), 163-172 (1987)

MEDLINE 87217962

COMMENT PROVISIONAL REFSEQ: This record has not yet been
 5 subject to final
 NCBI review. The reference sequence was derived from
 M16653.1.

FEATURES Location/Qualifiers

10 source 1..269
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /chromosome="12"
 /map="12"

15 Protein 1..269
 /product="pancreatic elastase IIB"
 sig_peptide 1..16
 /note="pancreatic elastase IIB signal
 peptide"
 Region 28..262
 20 /region_name="Trypsin-like serine protease"
 /db_xref="CDD:Tryp_SPC"
 /note="Tryp_SPC"
 mat_peptide 29..269
 /product="pancreatic elastase IIB mature
 25 peptide"
 Region 31..262
 /region_name="Trypsin"
 /db_xref="CDD:pfam00089"
 /note="trypsin"

30 CDS 1..269
 /gene="LOC51032"
 /db_xref="LocusID:51032"
 /coded_by="NM_015849.1:26..835"

ORIGIN

35 1 mirtlllsl vagalscgvs tyapdmrml ggeearpnsw pwqvslyyss
 ngqwyhtcgg
 61 slianswvlt aahciyssri yrvmlgqhn1 yvaesgslav svskivvhkd
 wnsnqvskgn
 121 diallklanp vsltldkiqla clppagtilp nnypcyvtgw grlqtngalp
 40 ddlkqgrllv
 181 vdyatcsssg wwgstvktnm icaggdgvic tcngdsggpl ncqasdgrwe
 vhgigsaltsv
 241 lgcnyyykps iftrvsnynd winsviann
 //

45

PRAME

50 LOCUS NM_006115 2148 bp mRNA PRI 19-JUN-
 2001
 DEFINITION Homo sapiens preferentially expressed antigen in
 melanoma (PRAME),
 mRNA.

55 ACCESSION NM_006115
 VERSION NM_006115.1 GI:5174640
 KEYWORDS .
 SOURCE human.
 ORGANISM Homo sapiens

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
 Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
 Homo.

5 REFERENCE 1 (bases 1 to 2148)
 AUTHORS Ikeda,H., Lethe,B., Lehmann,F., Van Baren,N.,
 Baurain,J.-F., De
 Smet,C., Chambost,H., Vitale,M., Moretta,A., Boon,T. and
 Coulie,P.G.

10 TITLE Characterization of an antigen that is recognized on a
 melanoma
 showing partial HLA loss by CTL expressing an NK inhibitor
 receptor

15 JOURNAL Immunity 6 (2), 199-208 (1997)
 MEDLINE 97199265
 REFERENCE 2 (bases 1 to 2148)
 AUTHORS Williams JM, Chen GC, Zhu L and Rest RF.
 TITLE Using the yeast two-hybrid system to identify human
 epithelial cell
 20 proteins that bind gonococcal Opa proteins: intracellular
 gonococci
 bind pyruvate kinase via their Opa proteins and
 require host
 pyruvate for growth

25 JOURNAL Mol. Microbiol. 27 (1), 171-186 (1998)
 MEDLINE 98125741
 PUBMED 9466265
 REFERENCE 3 (bases 1 to 2148)
 30 AUTHORS van Baren,N., Chambost,H., Ferrant,A., Michaux,L., Ikeda,H
 Millard,I., Olive,D., Boon,T. and Coulie,P.G.
 TITLE PRAME, a gene encoding an antigen recognized on a
 human melanoma by
 cytolytic T cells, is expressed in acute leukaemia cells

35 JOURNAL Br. J. Haematol. 102 (5), 1376-1379 (1998)
 MEDLINE 98423996
 PUBMED 9753074
 REFERENCE 4 (bases 1 to 2148)
 40 AUTHORS Dunham I, Shimizu N, Roe BA, Chisoe S, Hunt AR, Collins JI
 Bruskiewich R, Beare DM, Clamp M, Smink LJ, Ainscough
 R, Almeida
 JP, Babbage A, Bagguley C, Bailey J, Barlow K, Bates
 KN, Beasley O,
 Bird CP, Blakey S, Bridgeman AM, Buck D, Burgess J,
 Burrill WD,

45 O'Brien KP and et al.
 TITLE The DNA sequence of human chromosome 22
 JOURNAL Nature 402 (6761), 489-495 (1999)
 MEDLINE 20057165
 PUBMED 10591208

50 REMARK Erratum:[published erratum appears in Nature 2000 Apr
 20;404(6780):904]]
 COMMENT REVIEWED REFSEQ: This record has been curated by NCBI
 staff. The
 reference sequence was derived from U65011.1.
 55 Summary: The protein encoded by this gene has a 509
 amino acid
 antigen, lacking a signal sequence, and recognized on
 a human

melanoma cell line by a T-lymphocyte clone. A significant level of this mRNA is detected in normal testis as well as in many melanomas, non-small cell lung carcinomas, sarcomas, head and neck tumors and renal carcinomas. The encoded protein is expressed predominantly in acute leukemias carrying chromosomal abnormalities such as translocation t(8:21), which fuses the AML1 and ETO genes. Its expression shares several characteristics with the expression patterns of MAGE, BAGE, and GAGE gene families, all of which are expressed in tumors. This protein is expressed in a higher proportion of samples than genes of the MAGE, BAGE, and GAGE families, and of these four groups, only this protein is expressed by acute myeloid leukemias.

COMPLETENESS: complete on the 3' end.

25 FEATURES Location/Qualifiers
 source 1..2148
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /chromosome="22"
 30 /map="22q11.22"
 /haplotype="A24 A28 B13 B44 Cw6 Cw7"
 gene 1..2148
 /gene="PRAME"
 /note="MAPE; OIP4"
 35 /db_xref="LocusID:23532"
 /db_xref="MIM:606021"
 CDS 236..1765
 /gene="PRAME"
 /note="melanoma antigen preferentially
 40 expressed in tumors; Opa-interacting protein OIP4"
 /codon_start=1
 /db_xref="LocusID:23532"
 /db_xref="MIM:606021"
 45 /product="preferentially expressed antigen of
 melanoma"
 /protein_id="NP_006106.1"
 /db_xref="GI:5174641"
 50 /translation="MERRRLWGSIQSRYSMSVWTSPPRLVELAGQSLLKDEALAIAALELLPREL
 FPPLFMAAFDGRHSQTLKAMVQAWPFTCLPLGVLKMGQHLHLETFKAVLDGLDVLAAQEVPRRWK
 LQVLDLRKNSHQDFWTVWSGNRSALYSFPEPEAAQPMTKKRKVDGLSTEAQQPFIPVEVLVDLFLK
 EGACDELFSYLIEKVKRKKNVLRCLCKKLLKIFAMPQDIKMLKMWQLDSIEDLEVTCTWKLPTLA
 KFSPYLGQMINLRLLLLSHIHASSYISPEKEEQVIAQFTSQFLSLQCLQALYVDSLFFLRGRILDQL
 55 LRHVMNPLETLSITNCRLSEGDVMHLSQSPSVLSVLSLSGVMLTDVSPEPLQALLERASATLQD
 LVFDECGITDDQLLALLPSLSHCSQLTTLSTFYGNSISISALQSLLOHLIGLSNLTHVLYVPPLSY
 EDINGTLHLERLAYLHARLRELLCELGRPSMVWLSANPCPHCGDRTFYDPEPILCPCFMPN"
 variation 254


```

/allele="C"
/allele="T"
/db_xref="dbSNP:1129172"
5      variation      698
/allele="C"
/allele="A"
/db_xref="dbSNP:1063107"
      variation      1001
/allele="C"
10     /allele="T"
/db_xref="dbSNP:2229695"
      variation      1498
/allele="T"
/allele="C"
15     /db_xref="dbSNP:2229696"
      variation      1762
/allele="C"
/allele="T"
/db_xref="dbSNP:13604"
20     variation      1952
/allele="A"
/allele="G"
/db_xref="dbSNP:7104"
      variation      1984
/allele="A"
/allele="C"
/db_xref="dbSNP:8405"
25     variation      2043
/allele="T"
/allele="A"
/db_xref="dbSNP:1804685"
30     polyA_signal    2108..2113
      polyA_site      2130
BASE COUNT      534 a      548 c      558 g      508 t
35     ORIGIN
      1 gcttcagggt acagctcccc cgcagccaga agccgggcct gcagccctc
      agcacgcctc
      61 cgggacaccc caccgccttc ccaggcgtga cctgtcaaca gcaacttcgc
      ggtgtggtga
40     121 actctctgag gaaaaacccat tttgattatt actctcagac gtgcgtggca
      acaagtgact
      181 gagacctaga aatccaagcg ttggagggtcc tgaggccagc ctaagtcgct
      tcaaaatgga
      241 acgaaggcgt ttgtgggggt ccattcagag ccgatacatc agcatgagtg
45     tgtggacaag
      301 cccacggaga cttgtggagc tggcagggca gagcctgctg aaggatgagg
      ccctggccat
      361 tgccgccctg gagttgctgc ccaggagct cttcccgcca ctcttcattg
      cagcctttga
50     421 cgggagacac agccagaccc tgaaggcaat ggtgcaggcc tggcccttca
      cctgcctccc
      481 tctgggagtg ctgatgaagg gacaacatct tcacctggag accttcaaag
      ctgtgcttga
      541 tggacttgat gtgctccttg cccaggaggt tcgccccagg aggtggaaac
55     ttcaagtgct
      601 ggatttacgg aagaactctc atcaggactt ctggactgta tggctctggaa
      acagggccag

```

661 tctgtactca tttccagagc cagaagcagc tcagcccatg acaaagaagc
 gaaaagtaga
 721 tggtttgagc acagaggcag agcagccctt cattccagta gaggtgctcg
 tagacctgtt
 5 781 cctcaaggaa ggtgcctgtg atgaattgtt ctctacctc attgagaaag
 tgaagcgaaa
 841 gaaaaatgta ctacgcctgt gctgtaagaa gctgaagatt tttgcaatgc
 ccatgcagga
 901 tatcaagatg atcctgaaaa tgggtgcagct ggactctatt gaagatttgg
 10 aagtgacttg
 961 tacctggaag ctaccacct tggcgaaatt ttctccttac ctgggccaga
 tgattaatct
 1021 gcgtagactc ctctctccc acatccatgc atcttcctac atttccccgg
 agaaggaaga
 15 1081 gcagtatatc gcccagttca cctctcagtt cctcagtcg cagtgcctgc
 aggctctcta
 1141 tgtggactct ttatTTTTcc ttagaggcog cctggatcag ttgtcaggc
 acgtgatgaa
 1201 ccccttgga accctctcaa taactaactg ccggctttcg gaaggggatg
 20 tgatgcatct
 1261 gtcccagagt cccagcgtca gtcagctaag tgtcctgagt ctaagtgggg
 tcatgctgac
 1321 cgatgtaagt cccgagcccc tccaagctct gctggagaga gcctctgcca
 ccctccagga
 25 1381 cctgggtcttt gatgagtgtg ggatcacgga tgatcagctc cttgccctcc
 tgccttcct
 1441 gagccactgc tcccagctta caaccttaag cttctacggg aattccatct
 ccatatctgc
 1501 cttgcagagt ctctgcagc acctcatcgg gctgagcaat ctgaccacg
 30 tgctgtatcc
 1561 tgtccccctg gagagttatg aggacatcca tggtaacctc cacctggaga
 ggcttgcccta
 1621 tctgcatgcc aggctcaggg agttgctgtg tgagttgggg cggcccagca
 tgggtctggct
 35 1681 tagtgccaac cctgtcctc actgtgggga cagaaccttc tatgacctcg
 agcccatcct
 1741 gtgcccctgt ttcatgccta actagctggg tgcacatata aaatgcttca
 ttctgcatac
 1801 ttggacacta aagccaggat gtgcatgcat cttgaagcaa caaagcagcc
 40 acagtttcag
 1861 acaaatgttc agtgtgagtg aggaaaacat gttcagtgag gaaaaaacat
 tcagacaaat
 1921 gttcagtgag gaaaaaaagg ggaagttggg gataggcaga tgttgacttg
 aggagttaat
 45 1981 gtgatctttg gggagataca tcttatagag ttagaaatag aatctgaatt
 tctaaaggga
 2041 gattctggct tgggaagtac atgtaggagt taatccctgt gtagactgtt
 gtaaagaaac
 2101 tgttgaaaat aaagagaagc aatgtgaagc aaaaaaaaaa aaaaaaaa

ED-B domain of Fibronectin

LOCUS HSFIBEDB 2823 bp DNA linear
 PRI 09-AUG-1999

5 DEFINITION Human fibronectin gene ED-B region.
 ACCESSION X07717
 VERSION X07717.1 GI:31406
 KEYWORDS alternate splicing; fibronectin.
 SOURCE human.

10 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
 Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
 Homo.

15 REFERENCE 1 (bases 1 to 2823)
 AUTHORS Paoletta,G., Henchcliffe,C., Sebastio,G. and
 Baralle,F.E.
 TITLE Sequence analysis and in vivo expression show that
 alternative
 20 splicing of ED-B and ED-A regions of the human
 fibronectin gene are
 independent events
 JOURNAL Nucleic Acids Res. 16 (8), 3545-3557 (1988)
 MEDLINE 88233940

25 FEATURES Location/Qualifiers
 source 1..2823
 /organism="Homo sapiens"
 /db_xref="taxon:9606"
 /clone="MA10"

30 exon 1..104
 /number=1
 /product="fibronectin"

CDS join(<2..104,1375..1647,2758..>2823)
 /codon_start=1
 35 /product="fibronectin"
 /protein_id="CAB52437.1"
 /db_xref="GI:5725425"

/translation="CTFDNLSPGLEYNVSVYTVKDDKESVPISDTIPEVPQLTDLSF

VDITDSSIGLRWTPPLNSSTIIGYRITVVAAGEGIPIFEDFVDSSVGYYTGTGLEPGID

YDISVITLINGGESAPTTLTQQTAVPPPTDLRFTNIGPDIMRVTV"

```

5      intron          105..1374
                        /number=1
      exon            1375..1647
                        /note="ED-B exon"
                        /number=2
10      intron          1648..2757
                        /product="fibronectin"
                        /number=2
      exon            2758..2823
                        /number=3
15      /product="fibronectin"
BASE COUNT      824 a    556 c    528 g    915 t
ORIGIN
      1 ctgcactttt gataacctga gtcccgccct ggagtacaat gtcagtgttt
      acactgtcaa
20      61 ggatgacaag gaaagtgtcc ctatctctga taccatcatc ccaggtaata
      gaaaataagc
      121 tgctatcctg agagtgcacat tccaataaga gtggggatta gcatcttaat
      ccccagatgc
      181 ttaaggggtgt caactatatt tgggatttaa ttccgatctc ccagctgcac
25      tttccaaaac
      241 caagaagtca aagcagcgat ttggacaaaa tgcttgctgt taacactgct
      ttactgtctg
      301 tgcttcaactg ggatgctgtg tgttgcagcg agtatgtaat ggagtggcag
      ccatggcttt
30      361 aactctgtat tgtctgctca catggaagta tgactaaaac actgtcacgt
      gtctgtactc
      421 agtactgata ggctcaaagt aatatggtaa atgcatcca tcagtacatt
      tctgcccgat
      481 ttacaaatcc atatcaattt ccaacagctg cctatttcat cttgcagttt
35      caaatccttc
      541 tttttgaaaa ttggatttta aaaaaaagtt aagtaaaagt cacacottca
      gggttgttct
      601 ttcttggtggc cttgaaagac aacattgcaa aggcctgtcc taaggatagg
      cttgtttgtc

```

661 cattgggtta taacataatg aaagcattgg acagatcgtg tcccccttg
gactcttcag
721 tagaatgctt ttactaacgc taattacatg ttttgattat gaatgaacct
aaaatagtgg
5 781 caatggcctt aacctaggcc tgtctttcct cagcctgaat gtgcttttga
atggcacatt
841 tcacaccata cattcataat gcattagcgt tatggccatg atgttgcac
gagttttgta
901 tgggagaaaa aaaatcaatt taccacccat ttattatttt ttccggttgt
10 tcatgcaagc
961 ttatttttcta ctaaaacagt tttggaatta ttaaaagcat tgctgatact
tacttcagat
1021 attatgtcta ggctctaaga atggtttcga catcctaaac agccatatga
tttttaggaa
15 1081 tctgaacagt tcaaattgta ccctttaagg atgttttcaa aatgtaaaaa
atatatatat
1141 atatatatat tccctaaaag aatattcctg tttattcttc tagggaagca
aactgttcat
1201 gatgcttagg aagtcttttc agagaattta aaacagattg catattacca
20 tcattgcttt
1261 aacattccac caattttact actagtaacc tgatatacac tgctttattt
tttctcttt
1321 ttttccctct attttcttt tgcctcccc tccctttgct ttgtaactca
atagaggtgc
25 1381 cccaactcac tgacctaac tttgttgata taaccgatc aagcatcggc
ctgaggtgga
1441 ccccgctaaa ctcttcacc attattgggt accgcatcac agtagttgag
gcaggagaag
1501 gtatccctat ttttgaagat tttgtggact cctcagtagg atactacaca
30 gtcacagggc
1561 tggagccggg cattgactat gatatcagcg ttatcactct cattaatggc
ggcgagagtg
1621 cccctactac actgacacaa caaacgggtg aattttgaaa acttctgcgt
ttgagacata
35 1681 gatgggtgtg catgctgcca ccagttactc cggttaaata tggatgttcc
atgggggaag
1741 tcagcaattg gccaaagatt cagatagggtg gaattggggg gataaggaat
caaatgcac

1801 tgctaaactg attggagaaa aacacatgca atatcttcag tacactctca
tttaaaccac
1861 aagtagatat aaagcctaga gaaatacaga tgtctgctct gttaaata
aaatagcaaa
5 1921 tgttcattca atttgaagac ctagaatttt tcttcttaaa taccaaacac
gaataccaaa
1981 ttgcgtaagt accaattgat aagaatatat caccaaaatg taccatcatg
ctcttccttc
2041 taccctttga taaactctac catgctcctt ctttgtagct aaaaacccat
10 caaaatttag
2101 ggtagagtgg atgggcattg ttttgaggta ggagaaaagt aaacttgga
ccattctagg
2161 ttttgttget gtcactaggt aaagaaacac ctctttaacc acagtctggg
gacaagcatg
15 2221 caacatttta aaggttctct gctgtgcatg ggaaaagaaa catgctgaga
accaatttgc
2281 atgaacatgt tcacttgtaa gtagaattca ctgaatggaa ctgtagctct
agatatctca
2341 catgggggga agtttaggac cctcttgtct ttttgtctgt gtgcatgtat
20 ttctttgtaa
2401 agtactgcta tgtttctctt tgctgtgtgg caacttaagc ctcttcggcc
tgggataaaa
2461 taatctgcag tggattaat aatgtacata aagtcaacat atttgaaagt
agattaaaat
25 2521 ctttttttaa tatatcaatg atggcaaaaa gggttaaaggg ggcctaacag
tactgtgtgt
2581 agtggtttat ttttaacagt agtacactat aacttaaaat agacttagat
tagactgttt
2641 gcatgattat gattctgttt cctttatgca tgaaatattg attttacctt
30 tccagctact
2701 tcgttagctt taatttttaa atacattaac tgagtcttcc ttcttggtog
aaaccagctg
2761 ttcctcctcc cactgacctg cgattcacca acattggtcc agacaccatg
cgtgtcacct
35 2821 ggg
//

CEA
LOCUS NM_004363 2974 bp mRNA linear
PRI 28-NOV-2000
DEFINITION Homo sapiens carcinoembryonic antigen-related cell
5 adhesion
molecule 5 (CEACAM5), mRNA.
ACCESSION NM_004363
VERSION NM_004363.1 GI:11386170
KEYWORDS .
10 SOURCE human.
ORGANISM Homo sapiens
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
Euteleostomi;
Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
15 Homo.
REFERENCE 1 (bases 1 to 2974)
AUTHORS Oikawa S, Nakazato H and Kosaki G.
TITLE Primary structure of human carcinoembryonic antigen
(CEA) deduced
20 from cDNA sequence
JOURNAL Biochem. Biophys. Res. Commun. 142 (2), 511-518 (1987)
MEDLINE 87128144
PUBMED 3814146
REFERENCE 2 (bases 1 to 2974)
25 AUTHORS Zimmermann W, Weber B, Ortlieb B, Rudert F, Schempp W,
Fiebig HH,
Shively JE, von Kleist S and Thompson JA.
TITLE Chromosomal localization of the carcinoembryonic
antigen gene
30 family and differential expression in various tumors
JOURNAL Cancer Res. 48 (9), 2550-2554 (1988)
MEDLINE 88184584
PUBMED 3356015
REFERENCE 3 (bases 1 to 2974)
35 AUTHORS Barnett, T., Goebel, S.J., Nothdurft, M.A. and
Elting, J.J.
TITLE Carcinoembryonic antigen family: characterization of
cDNAs coding

for NCA and CEA and suggestion of nonrandom sequence
variation in

their conserved loop-domains

JOURNAL Genomics 3 (1), 59-66 (1988)

5 MEDLINE 89122014

REFERENCE 4 (bases 1 to 2974)

AUTHORS Barnett T and Zimmermann W.

TITLE Workshop report: proposed nomenclature for the
carcinoembryonic

10 antigen (CEA) gene family

JOURNAL Tumour Biol. 11 (1-2), 59-63 (1990)

MEDLINE 90176333

PUBMED 2309067

REFERENCE 5 (bases 1 to 2974)

15 AUTHORS Schrewe H, Thompson J, Bona M, Hefta LJ, Maruya A,
Hassauer M,

Shively JE, von Kleist S and Zimmermann W.

TITLE Cloning of the complete gene for carcinoembryonic
antigen: analysis

20 of its promoter indicates a region conveying cell
type-specific

expression

JOURNAL Mol. Cell. Biol. 10 (6), 2738-2748 (1990)

MEDLINE 90258861

25 PUBMED 2342461

COMMENT PROVISIONAL REFSEQ: This record has not yet been
subject to final

NCBI review. The reference sequence was derived from
M29540.1.

30 FEATURES Location/Qualifiers

source

1..2974

/organism="Homo sapiens"

/db_xref="taxon:9606"

/chromosome="19"

35 /map="19q13.1-q13.2"

gene

1..2974

/gene="CEACAM5"

/note="CEA; CD66e"

/db_xref="LocusID:1048"

5 /db_xref="MIM:114890"
CDS 115..2223
 /gene="CEACAM5"
 /note="carcinoembryonic antigen"
 /codon_start=1
 /db_xref="LocusID:1048"
 /db_xref="MIM:114890"
 /product="carcinoembryonic antigen-related
cell adhesion
10 molecule 5"
 /protein_id="NP_004354.1"
 /db_xref="GI:11386171"

15 /translation="MESPSAPPHRWCIPWQRLLLTASLLTFWNPPTAKLTIESTPFN
VAEGKEVLLLVHNLPOHLFGYSWYKGERVDGNRQIIGYVIGTQQATPGPAYSGREIYY
PNASLLIQNI IQNDTGFTLHVIKSDLVNEEATGQFRVYPELPKPSISSNNSKPVEDK
DAVAFTCEPETQDATYLWWVNNQSLPVSPRLQLSNGNRTLTLFNVTRNDTASYKCETQ
NPVSARRSDSVILNVLYGPDAPTISPLNTSYRSGENLNLSCHAASNPPAQYSWFVNGT
FQOSTQELFIPNITVNNSGSYTCQAHNSDTGLNRTTVTTITVYAEPPKPFITSNNSNP
20 VEDEDAVALTCEPEIQNTTYLWWVNNQSLPVSPRLQLSNDNRTLTLTLLSVTRNDVGPYE
CGIQNELSVDHSDPVILNVLYGPDPTISPSYTYRPGVNL SLSCHAASNPPAQYSWL
IDGNIQQHTQELFISNITEKNSGLYTCQANNSASGHSRTTVKTTITVSAELPKPSISSN
NSKPVEDKDAVAFTCEPEAQNTTYLWWVNGQSLPVSPRLQLSNGNRTLTLFNVTRNDA
RAYVCGIQNSVSANRSDPVTLDVLYGPDPTIISPDSYLSGANLNLSCHSASNPSPQ
25 YSWRINGIQQHTQVLFIAKITPNNNGTYACFVSNLATGRNNSIVKSITVSASGTSPG
LSAGATVGIMIGVLVGVALI"
 misc_feature 595..780
 /note="IG; Region: Immunoglobulin"
 misc_feature 868..1026
30 /note="IG; Region: Immunoglobulin"
 misc_feature 1129..1314
 /note="IG; Region: Immunoglobulin"
 misc_feature 1399..1566
 /note="IG; Region: Immunoglobulin"
35 misc_feature 1930..2091
 /note="IG; Region: Immunoglobulin"
BASE COUNT 840 a 847 c 613 g 674 t
ORIGIN

1 ctcagggcag agggaggaag gacagcagac cagacagtca cagcagcctt
 gacaaaacgt
 61 tcctggaact caagctcttc tccacagagg aggacagagc agacagcaga
 gaccatggag
 5 121 tctccctcgg cccctcccca cagatggtgc atccctggc agaggctcct
 gctcacagcc
 181 tcacttctaa ccttctggaa cccgccacc actgccaaagc tcactattga
 atccacgccg
 241 ttcaatgtcg cagaggggaa ggaggtgctt ctacttgtcc acaatctgcc
 10 ccagcatctt
 301 tttggctaca gctggtacaa aggtgaaaga gtggatggca accgtcaa
 tataggatat
 361 gtaataggaa ctcaacaagc taccacaggc cccgcataca gtggtcgaga
 gataatatac
 15 421 cccaatgcat cctgctgat ccagaacatc atccagaatg acacaggatt
 ctacacccta
 481 cagtcataa agtcagatct tgtgaatgaa gaagcaactg gccagttccg
 ggtatacccg
 541 gagctgccc aacctccat ctceagcaac aactccaaac ccgtggagga
 20 caaggatgct
 601 gtggccttca cctgtgaacc tgagactcag gacgcaacct acctgtggtg
 ggtaaacaat
 661 cagagcctcc cggtcagtcc caggctgcag ctgtccaatg gcaacaggac
 cctcactcta
 25 721 ttcaatgtca caagaaatga cacagcaagc taaaaatgtg aaaccagaa
 cccagtga
 781 gccaggcgca gtgattcagt catcctgaat gtccctctatg gcccgatgc
 cccaccatt
 841 tccctctaa acacatctta cagatcaggg gaaaatctga acctctctg
 30 ccacgcagcc
 901 tctaaccac ctgcacagta ctcttggtt gtcaatggga ctttcagca
 atccaccaa
 961 gagctcttta tcccaacat cactgtgaat aatagtggat cctatacgtg
 ccaagccat
 1021 aactcagaca ctggcctcaa taggaccaca gtcacgacga tcacagtcta
 35 tgcagagcca
 1081 cccaaacct tcatcaccag caacaactcc aaccccgagg aggatgagga
 tgctgtagcc

1141 ttaacctgtg aacctgagat tcagaacaca acctacctgt ggtgggtaaa
 taatcagagc
 1201 ctcccgggtca gtcccaggtc gcagctgtcc aatgacaaca ggaccctcac
 tctactcagt
 5 1261 gtcacaagga atgatgtagg accctatgag tgtggaatcc agaacgaatt
 aagtgttgac
 1321 cacagcgacc cagtcaccc cagtcatcct gaatgtcctc tatggcccag acgacccccc
 catttcccc
 1381 tcatacacct attaccgtcc aggggtgaac ctccagcctct cctgccatgc
 10 agcctctaac
 1441 ccacctgcac agtattcttg gctgattgat gggaacatcc agcaacacac
 acaagagctc
 1501 tttatctcca acatcactga gaagaacagc ggactctata cctgccaggc
 caataactca
 15 1561 gccagtggcc acagcaggac tacagtcaag acaatcacag tctctgcgga
 gctgcccag
 1621 ccctccatct ccagcaacaa ctccaaaccc gtggaggaca aggatgctgt
 ggccttcacc
 1681 tgtgaacctg aggctcagaa cacaacctac ctgtgggtggg taaatgggtca
 20 gagcctccca
 1741 gtcagtccca ggctgcagct gtccaatggc aacaggaccc tcactctatt
 caatgtcaca
 1801 agaaatgacg caagagccta tgtatgtgga atccagaact cagtgagtgc
 aaaccgcagt
 25 1861 gacccagtca ccctggatgt cctctatggg ccggacaccc ccatcatttc
 ccccccagac
 1921 tcgtcttacc ttctgggagc gaacctcaac ctctcctgcc actcggcctc
 taaccoatcc
 1981 ccgcagtatt cttggcgat caatgggata ccgcagcaac acacacaagt
 30 tctctttatc
 2041 gccaaaatca cgccaaataa taacgggacc tatgcctgtt ttgtctctaa
 cttggctact
 2101 ggccgcaata attccatagt caagagcatc acagtctctg catctggaac
 ttctcctggc
 35 2161 ctctcagctg gggccactgt cggcatcatg attggagtgc tgggtggggc
 tgctctgata
 2221 tagcagccct ggtgtagttt cttcatttca ggaagactga cagttgtttt
 gcttcttctc

2281 taaagcattt gcaacagcta cagtctaaaa ttgcttcttt accaaggata
 tttacagaaa
 2341 agactctgac cagagatoga gaccatccta gccaacatcg tgaaacccca
 tctctactaa
 5 2401 aaatacaaaa atgagctggg cttggtggcg cgcacctgta gtcccagtta
 ctcgggagggc
 2461 tgaggcagga gaatcgcttg aaccggggag gtggagattg cagtgaagccc
 agatcgaccc
 2521 actgcactcc agtctggcaa cagagcaaga ctccatctca aaaagaaaag
 10 aaaagaagac
 2581 tctgacctgt actcttgaat acaagtttct gataccactg cactgtctga
 gaatttccaa
 2641 aactttaatg aactaactga cagcttcatg aaactgtcca ccaagatcaa
 gcagagaaaa
 15 2701 taattaattt catgggacta aatgaactaa tgaggattgc tgattcttta
 aatgtcttgt
 2761 tttccagatt tcaggaaact ttttttcttt taagctatcc actcttacag
 caatttgata
 2821 aaatatactt ttgtgaacaa aaattgagac atttacattt tctccctatg
 20 tggtcgctcc
 2881 agacttggga aactattcat gaatatttat attgtatggg aatatagtta
 ttgcacaagt
 2941 tcaataaaaa tctgctcttt gtataacaga aaaa
 //

25

Her2/Neu
 LOCUS HUMHER2A 4530 bp mRNA linear
 PRI 18-SEP-1995
 30 DEFINITION Human tyrosine kinase-type receptor (HER2) mRNA,
 complete cds.
 ACCESSION M11730
 VERSION M11730.1 GI:183986
 KEYWORDS tyrosine kinase.
 35 SOURCE Homo sapiens (clone: lambda-HER2-436) fetal cDNA to
 mRNA.
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
 Euteleostomi;

Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
Homo.

REFERENCE 1 (bases 1 to 4530)

AUTHORS Coussens, L., Yang-Feng, T.L., Liao, Y.-C., Chen, E.,
5 Gray, A.,
McGrath, J., Seeburg, P.H., Libermann, T.A.,
Schlessinger, J.,
Francke, U., Levinson, A. and Ullrich, A.

TITLE Tyrosine kinase receptor with extensive homology to
10 EGF receptor
shares chromosomal location with neu oncogene

JOURNAL Science 230 (4730), 1132-1139 (1985)

MEDLINE 86070181

REFERENCE 2 (bases 1701 to 1719)

15 AUTHORS Ullrich, A.
JOURNAL Unpublished (1988)

FEATURES Location/Qualifiers

source 1..4530
/organism="Homo sapiens"
20 /db_xref="taxon:9606"
/clone="lambda-HER2-436"
/dev_stage="fetal"

mRNA <1..4530
/product="HER2 mRNA"

25 CDS 151..3918
/note="HER2 receptor"
/codon_start=1
/protein_id="AAA75493.1"
/db_xref="GI:306840"

30 /translation="MELAALCRWGLLLALLPPGAASTQVCTGTDMKLRLPASPETHLD
MLRHLYQGCQVVQGNLELTYLPTNASLSFLQDIQEVQGYVLIAHNQVRQVPLQRLRIV
RGTQLFEDNYALAVLDNGDPLNNTTPVTGASPGGLRELQLRSLTEILKGGVLIQRNPQ
LCYQDTILWKDIFHKNNQLALTLDITNRSRACHPCSPMCKGSRWGESSDCQSLTRT
35 VCAGGCARCKGPLPTDCCHEQCAAGCTGPKHSDCLACLFHNSGICELHCPALVTYNT
DTFESMPNPEGRYTFGASCVTACPYNYLSTDVGSCTLVCPHNPQEVTAEDGTQRCEKC
SKPCARVCYGLGMEHLREVRVAVTSANIQEFAGCKKIFGSLAFLPESFDGDPASNTAPL
QPEQLQVFETLEEITGYLYISAWPDSLPLSVFQNLQVIRGRILHNGAYSLTLQGLGI
SWLGLRSLRELGSGLALIHNTLHLCFVHTVPWDQLFRNPHQALLHTANRPEDECVGEG

LACHQLCARGHCWGPQTQCVNCSQFLRGQBCVEECRVLQGLPREYVNARHCLPCHPE
 CQPQNGSVTCFGPEADQCVACAHYKDPFFCVARCPGSKPDLSPYMPIWKFPDEEGACQ
 PCPINCTHSCVDLDDKGCPAEQRASPLTSIVSAVVGILLVVVLGVVFGILIKRRQQKI
 RKYTMRRLLQETELVEPLTPSGAMPNQAQMRILKETELRKVKVLGSGAFGTVYKGIWI
 5 PDGENVKIPVAIKVLRENTSPKANKEILDEAYVMAGVGSPYVSRLLGICLTSTVQLVT
 QLMPYGCLLDHVRENRLGSLQDLNWCMIKAGMSYLEDVRLVHRDLAARNVLVKSP
 NHVKITDFGLARLLDIDETEHADGGKVPIKWMALLESILRRRFTHQSDVWSYGVTVWE
 LMTFGAKPYDGIPIREIPDLLEKGERLPQPPICTIDVYIMVKCWMIDSECRPRFREL
 VSEFSRMARDPQRFVVIQNEDLGPASPLDSTFYRSLLEDDDMGDLVDAEEYLVPQQGF
 10 FCPDPAPGAGGMVHHRSSSTRSGGDLTLGLEPSEEEAPRSPAPSEGAGSDVFDG
 DLGMGAAGLQSLPHTDPSPLQRYSEDPTVPLPSETDGYVAPLTCSPQPEYVNQPDVR
 PQPPSPREGPLPAARPAGATLERAKTILSPGKNGVVKDVFAFGGAVENPEYLTPOGGAA
 PQPHPPPAFSPAFDNLYYWDQPPPERGAPPSTFKGTPTAENPEYLGLDVPV"
 old_sequence 1701..1719
 15 /citation=[1]
 BASE COUNT 922 a 1382 c 1346 g 880 t
 ORIGIN Chromosome 17q21-q22.
 1 aattctcgag ctctctgacc ggtcgacgag ctcgagggtc gacgagctcg
 agggcgcgcg
 20 61 cccggccccc acccctcgca gcaccccgcg ccccgcgccc tcccagccgg
 gtccagccgg
 121 agccatgggg ccggagccgc agtgagcacc atggagctgg cggccttggtg
 ccgctggggg
 181 ctctctctcg ccctcttgcc ccccgagacc gcgagcacc aagtgtgcac
 25 cggcacagac
 241 atgaagctgc ggctccctgc cagtcccgag acccacctgg acatgctccg
 ccacctctac
 301 cagggctgcc aggtggtgca gggaaacctg gaactcacct acctgcccac
 caatgccagc
 361 ctgtccttcc tgcaggatat ccaggaggtg cagggctacg tgctcatcgc
 tcacaaccaa
 421 gtgaggcagg tcccactgca gaggctgcgg attgtgagag gaacccagct
 ctttgaggac
 481 aactatgccc tggccgtgct agacaatgga gaccgctga acaataccac
 35 ccctgtcaca
 541 ggggcctccc caggaggcct gcgggagctg cagcttcgaa gcctcacaga
 gatcttgaaa
 601 ggaggggtct tgatccagcg gaacccccag ctctgctacc aggacacgat
 tttgtggaag

	661	gacatcttcc	acaagaacaa	ccagctggct	ctcacactga	tagacaccaa
		ccgctctcgg				
	721	gcctgccacc	cctgttctcc	gatgtgtaag	ggctcccgt	gctggggaga
		gagttctgag				
5	781	gattgtcaga	gcctgacgcg	cactgtctgt	gccggtggct	gtgcccgtg
		caaggggcca				
	841	ctgcccactg	actgctgcca	tgagcagtgt	gctgccggct	gcacggggccc
		caagcactct				
10	901	gactgcctgg	cctgcctoca	cttcaaccac	agtggcatct	gtgagctgca
		ctgcccagcc				
	961	ctggtcacct	acaacacaga	cacgtttgag	tccatgccca	atcccagagg
		ccggtataca				
	1021	ttcggcgcca	gctgtgtgac	tgctgtccc	tacaactacc	tttctacgga
		cgtgggatcc				
15	1081	tgcaccctcg	tctgccccct	gcacaaccaa	gaggtgacag	cagaggatgg
		aacacagcgg				
	1141	tgtgagaagt	gcagcaagcc	ctgtgcccga	gtgtgctatg	gtctgggcat
		ggagcacttg				
	1201	cgagaggtga	gggcagttac	cagtgccaat	atccaggagt	ttgctggctg
20		caagaagatc				
	1261	tttgggagcc	tggcatttct	gccggagagc	tttgatgggg	accagcctc
		caacactgcc				
	1321	ccgctccagc	cagagcagct	ccaagtgttt	gagactctgg	aagagatcac
		aggttaccta				
25	1381	tacatctcag	catggccgga	cagcctgcct	gacctcagcg	tcttccagaa
		cctgcaagta				
	1441	atccggggac	gaattctgca	caatggcgcc	tactcgctga	ccctgcaagg
		gctgggcatc				
	1501	agctggctgg	ggctgcgctc	actgagggaa	ctgggcagtg	gactggccct
30		catccaccat				
	1561	aacacccacc	tctgcttcgt	gcacacgggtg	ccctgggacc	agctctttcg
		gaacccgcac				
	1621	caagctctgc	tccacactgc	caaccggcca	gaggacgagt	gtgtgggcga
		gggcctggcc				
35	1681	tgccaccagc	tgtgcgcccc	agggcactgc	tggggtccag	ggcccaccca
		gtgtgtcaac				
	1741	tgcagccagt	tccttcgggg	ccaggagtgc	gtggaggaat	gccgagtact
		gcaggggctc				

1801 cccagggagt atgtgaatgc caggcactgt ttgccgtgcc accctgagtg
 tcagccccag
 1861 aatgggtcag tgacctgttt tggaccggag gctgaccagt gtgtggcctg
 tgcccactat
 5 1921 aaggaccctc ccttctgcgt ggcccgtgc cccagcggtg tgaaacctga
 cctctcctac
 1981 atgccccatct ggaagtttcc agatgaggag ggcgcatgcc agccttgccc
 catcaactgc
 2041 acccactcct gtgtggacct ggatgacaag ggctgccccg ccgagcagag
 10 agccagccct
 2101 ctgacgtcca tcgtctctgc ggtggttggc attctgctgg tcgtggtctt
 gggggtggtc
 2161 tttgggatcc tcatcaagcg acggcagcag aagatccgga agtacacgat
 gcggagactg
 15 2221 ctgcaggaaa cggagctggg ggagccgctg acacctagcg gagcgatgcc
 caaccaggcg
 2281 cagatgcgga tcctgaaaga gacggagctg aggaaggtga aggtgcttgg
 atctggcgct
 2341 tttggcacag tctacaaggg catctggatc cctgatgggg agaattgtgaa
 20 aattccagtg
 2401 gccatcaaag tgttgaggga aaacacatcc cccaaagcca acaaagaaat
 cttagacgaa
 2461 gcatacgtga tggctgggtg gggctcccca tatgtctccc gccttctggg
 catctgcctg
 25 2521 acatccacgg tgcagctggg gacacagctt atgccctatg gctgcctctt
 agaccatgtc
 2581 cgggaaaacc gcggacgcct gggctcccag gacctgctga actggtgtat
 gcagattgcc
 2641 aaggggatga gctacctgga ggatgtgcgg ctctacaca gggacttggc
 30 cgctcggaac
 2701 gtgctgggtca agagtcccaa ccatgtcaaa attacagact tcgggctggc
 tcggctgctg
 2761 gacattgacg agacagagta ccatgcagat gggggcaagg tgcccatcaa
 gtggatggcg
 35 2821 ctggagtcca ttctccgccg gcggttcacc caccagagtg atgtgtggag
 ttatggtgtg
 2881 actgtgtggg agctgatgac ttttggggcc aaaccttaag atgggatccc
 agcccgggag

2941 atccctgacc tgctggaaaa gggggagcgg ctgccccagc ccccatctg
caccattgat
3001 gtctacatga tcatggtaa atgttgatg attgactctg aatgtcggcc
aagattccgg
5 3061 gagttggtgt ctgaattctc ccgcatggcc agggaccccc agcgctttgt
ggtcatccag
3121 aatgaggact tgggcccagc cagtccttg gacagcacct tctaccgctc
actgctggag
3181 gacgatgaca tgggggacct ggtggatgct gaggagtatc tggtagccca
10 gcagggttc
3241 ttctgtccag accctgcccc gggcgctggg ggcattggtc accacaggca
ccgcagctca
3301 tctaccagga gtggcgtgg ggacctgaca ctagggtgg agccctctga
agaggaggcc
15 3361 cccaggtctc cactggcacc ctccgaagg gctggctccg atgtatttga
tggtagacctg
3421 ggaatggggg cagccaagg gctgcaaagc ctccccacac atgaccccag
ccctctacag
3481 cggtagagtg aggacccac agtaccctg ccctctgaga ctgatggcta
20 cgttgcccc
3541 ctgacctgca gccccagcc tgaatatgtg aaccagccag atgttcggcc
ccagccccct
3601 tcgccccgag agggccctct gcctgctgcc cgacctgctg gtgccactct
ggaaaggggc
25 3661 aagactctct cccagggaa gaatggggtc gtcaaagacg tttttgcctt
tgggggtgcc
3721 gtggagaacc ccgagtactt gacaccccag ggaggagctg ccctcagcc
ccaccctct
3781 cctgccttca gccagcctt cgacaacctc tattactggg accaggaccc
30 accagagcgg
3841 ggggtctccac ccagcacctt caaagggaca cctacggcag agaaccaga
gtacctgggt
3901 ctggacgtgc cagtgtgaac cagaaggcca agtccgcaga agccctgatg
tgtcctcagg
35 3961 gagcagggaa ggctgactt ctgctggcat caagaggtgg gaggggcctc
cgaccacttc
4021 caggggaacc tgccatgcca ggaacctgtc ctaaggaacc ttcttctctg
cttgagttcc

```

      4081 cagatggctg gaaggggtcc agcctcgttg gaagaggaac agcactgggg
      agtctttgtg
      4141 gattctgagg ccctgcccaa tgagactcta ggggccagtg gatgccacag
      cccagcttgg
5      4201 cccttttcctt ccagatcctg ggtactgaaa gccttaggga agctggcctg
      agagggggaag
      4261 cggccctaag ggagtgtcta agaacaaaag cgacccattc agagactgtc
      cctgaaacct
      4321 agtactgccc cccatgagga aggaacagca atgggtgtcag tatccaggct
10     ttgtacagag
      4381 tgcttttctg tttagttttt actttttttg ttttgttttt ttaaagacga
      aataaagacc
      4441 caggggagaa tgggtgttgt atggggaggc aagtgtgggg ggtccttctc
      cacaccact
15     4501 ttgtccattt gcaaatatat tttggaaaac
      //

```

SCP-1

```

20     LOCUS      HSSCP1PRT              3393 bp      mRNA      linear
      PRI 26-FEB-1997
      DEFINITION  H.sapiens mRNA for SCP1 protein.
      ACCESSION   X95654
      VERSION     X95654.1  GI:1212982
25     KEYWORDS   SCP1 gene.
      SOURCE      human.
      ORGANISM    Homo sapiens
                  Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
                  Euteleostomi;
30     Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
      Homo.
      REFERENCE   1  (bases 1 to 3393)
      AUTHORS     Meuwissen,R.L., Meerts,I., Hoovers,J.M., Leschot,N.J.
      and
35     Heyting,C.
      TITLE       Human synaptonemal complex protein 1 (SCP1): isolation
      and
                  characterization of the cDNA and chromosomal
                  localization of the

```

gene
JOURNAL Genomics 39 (3), 377-384 (1997)
MEDLINE 97224467
REFERENCE 2 (bases 1 to 3393)
5 AUTHORS Meuwissen, R.J.L.
TITLE Direct Submission
JOURNAL Submitted (13-FEB-1996) Dr. R.L.J. Meuwissen,
Agricultural
University, Genetics, Dreijenlaan 2, 6703 HA
10 WAGENINGEN,
NETHERLANDS
FEATURES Location/Qualifiers
source 1..3393
/organism="Homo sapiens"
15 /db_xref="taxon:9606"
/map="lp12-pl3"
/tissue_type="testis"
gene 95..3025
/gene="SCP1"
20 CDS 95..3025
/gene="SCP1"
/codon_start=1
/evidence=experimental
/product="polypeptide of 976 aa"
25 /protein_id="CAA64956.1"
/db_xref="GI:1212983"
/db_xref="SWISS-PROT:Q15431"

/translation="MEKQKPFALFVPPRSSSSQVSAVKPQTLGGDSTFFKSFNKCTED
30 DLEFPFAKTNLSKNGENIDSDPALQKVNFLPVLEQVGNSDCHYQEGDKSDLENSEGL
SRVFSKLYKEAEKIKWKVSTAEALRQKESKLQENRKII EAQRKAIQELQFGNEKVSL
KLEEGIQENKDLIKENNATRHLCNLLKETCARSAEKTKKYEYEREETRQVYMDLNNNI
EKMITAHGELRVQAENSRLMHFKLKEDYEKIQHLEQYKKEINDKEKQVSLLLIQIT
EKENKMKDLTFLLEESRDKNQLEEKTKLQSENKQSI EKQHHLTKELEDIKVSLQRS
35 VSTQKALBEDLQIATKTICQLTEBKETQMEESNKARAAHSFVVTEFETTVCSLEELLR
TEQQRLEKNEDQLKILTMELQKKSSLEEMTKLTNNKEVELEELKKVLGEKETLLYEN
KQFEKIAEELKGTEQELIGLLQAREKEVHDLEIQLTAITTSQYYSKEVKDLKTELEN
EKLKNTELTSHCNKLSLENKELTQETSDMTLELKNQOEDINNKKQEERMLKQIENLQ
ETETQLRNELEYVREELKQKRDEVKCKLDKSEENCNNLRKQVENKNKYIEELQQENKA

LKKKGTAESQQLNVYBIKVNKLELELES AKQKFGEITDTYQKEIEDKKISEENLLEEV
 EKAKVIAD EAVKLQKEIDKRCQH KIAMVALMEKHKHQYDKIIEERDSELGLYKSKEQ
 EQSSLRASLEIELSNLKAEILSVKKQLEIEREEKEKLKREAKENTATLKEKKDKKTQT
 FLLETPEIYWKLDSKAVPSQTVSRNFTSVDHGISKDKRDYLWTS AKNTLSTPLPKAYT
 5 VKTPTPKKLQORENLNIP IESKKKRKMAFEFDINS DSSETTDLLSMVSEETLKTLY
 RNNNPPASHLCVKTPPKAPSSLTTPGPTLKFGAIRKMREDRWAVIAKMDRKKKLKEAE
 KLFV"
 BASE COUNT 1440 a 483 c 648 g 822 t
 ORIGIN
 10 1 gccctcatag accgtttggt gtagttcgcg tgggaacagc aaccacaggt
 ttcccgatag
 61 ttcttcaaag atatttaciaa ccgtaacaga gaaaatggaa aagcaaaagc
 cctttgcatt
 121 gttcgtacca ccgagatcaa gcagcagtcg ggtgtctcg gtgaaacctc
 15 agaccctggg
 181 aggcgattcc actttcttca agagtttcaa caaatgtact gaagatgatt
 tggagtttcc
 241 atttgcaaag actaatctct ccaaaaatgg ggaaaacatt gattcagatc
 ctgctttaca
 20 301 aaaagttaat ttcttgcccg tgcttgagca ggttggaat tctgactgtc
 actatcagga
 361 aggactaaaa gactctgatt tggagaattc agagggattg agcagagtgt
 tttcaaaaact
 421 gtataaggag gctgaaaaga taaaaaatg gaaagtaagt acagaagctg
 25 aactgagaca
 481 gaaagaaagt aagttgcaag aaaacagaaa gataattgaa gcacagcgaa
 aagccattca
 541 ggaactgcaa tttggaaatg aaaaagtaag tttgaaatta gaagaaggaa
 tacaagaaaa
 30 601 taaagattta ataaaagaga ataatgccac aaggcattta tgtaatctac
 tcaaagaaac
 661 ctgtgctaga tctgcagaaa agacaaagaa atatgaatat gaacggggaag
 aaaccaggca
 721 agttttatatg gatctaaata ataacattga gaaaatgata acagctcatg
 35 gggaaacttcg
 781 tgtgcaagct gagaattcca gactggaaat gcattttaag ttaaaggaag
 attatgaaaa
 841 aatccaacac cttgaacaag aatacaagaa ggaaataaat gacaaggaaa
 agcaggtatc

901 actactattg atccaaatca ctgagaaaga aaataaaatg aaagatttaa
 cattttctgct
 961 agaggaatcc agagataaag ttaatcaatt agaggaaaag acaaaattac
 agagtgaaaa
 5 1021 cttaaaacaa tcaattgaga aacagcatca tttgactaaa gaactagaag
 atattaaagt
 1081 gtcattacaa agaagtgtga gtactcaaaa ggcttttagag gaagatttac
 agatagcaac
 1141 aaaaacaatt tgtcagctaa ctgaagaaaa agaaactcaa atggaagaat
 10 ctaataaagc
 1201 tagagctgct cattcgtttg tggttactga atttgaaact actgtctgca
 gcttggaaga
 1261 attattgaga acagaacagc aaagattgga aaaaaatgaa gatcaattga
 aaatacttac
 15 1321 catggagctt caaaagaaat caagtgaact ggaagagatg actaagctta
 caaataacaa
 1381 agaagtagaa cttgaagaat tgaaaaaagt cttgggagaa aaggaaacac
 ttttatatga
 1441 aaataaaciaa tttgagaaga ttgctgaaga attaaaagga acagaacaag
 20 aactaattgg
 1501 tcttctccaa gccagagaga aagaagtaca tgatttgga atacagttaa
 ctgccattac
 1561 cacaagtga cagtattatt caaaagaggt taaagatcta aaaactgagc
 ttgaaaacga
 25 1621 gaagcttaag aatactgaat taacttcaca ctgcaacaag ctttcactag
 aaaacaaaga
 1681 gctcacacag gaaacaagt atatgaccct agaactcaag aatcagcaag
 aagatattaa
 1741 taataacaaa aagcaagaag aaaggatgtt gaaacaaata gaaaatcttc
 30 aagaaacaga
 1801 aaccaatta agaaatgaac tagaatatgt gagagaagag ctaaaacaga
 aaagagatga
 1861 agttaaatgt aaattggaca agagtgaaga aaattgtaac aatttaagga
 aacaagttga
 35 1921 aaataaaaac aagtatatgt aagaacttca gcaggagaat aaggccttga
 aaaaaaaagg
 1981 tacagcagaa agcaagcaac tgaatgttta tgagataaag gtcaataaat
 tagagttaga

2041 actagaaagt gccaaacaga aatttgagaa aatcacagac acctatcaga
 aagaaattga
 2101 ggacaaaaag atatcagaag aaaatctttt ggaagagggtt gagaaagcaa
 aagtaatagc
 5 2161 tgatgaagca gtaaaattac agaaagaaat tgataagcga tgtcaacata
 aaatagctga
 2221 aatggtagca cttatggaaa aacataagca ccaatatgat aagatcattg
 aagaaagaga
 2281 ctcagaatta ggactttata agagcaaaga acaagaacag tcatcactga
 10 gagcatcttt
 2341 ggagattgaa ctatccaatc tcaaagctga acttttgtct gttaagaagc
 aacttgaaat
 2401 agaaagagaa gagaaggaaa aactcaaaag agaggcaaaa gaaaacacag
 ctactcttaa
 15 2461 agaaaaaaaa gacaagaaaa cacaaacatt tttattggaa acacctgaaa
 tttattggaa
 2521 attggattct aaagcagttc cttcacaac tgtatctcga aatttcacat
 cagttgatca
 2581 tggcatatcc aaagataaaa gagactatct gtggacatct gccaaaaata
 20 ctttatctac
 2641 accattgccca aaggcatata cagtgaagac accaacaaaa caaaactac
 agcaaagaga
 2701 aaacttgaat ataccattg aagaaagtaa aaaaaagaga aaaatggcct
 ttgaatttga
 25 2761 tattaattca gatagttcag aaactactga tcttttgagc atggtttcag
 aagaagagac
 2821 attgaaaaca ctgtatagga acaataatcc accagcttct catctttgtg
 tcaaaacacc
 2881 aaaaaaggcc cttcatctc taacaacccc tggacctaca ctgaagtttg
 30 gagctataag
 2941 aaaaatgcgg gaggaccgtt gggctgtaat tgctaaaatg gatagaaaaa
 aaaaactaaa
 3001 agaagctgaa aagttatttg ttttaattca gagaatcagt gtagttaagg
 agcctaataa
 35 3061 cgtgaaactt atagttaata tttgttctt atttgccaga gccacatttt
 atctggaagt
 3121 tgagacttaa aaaatacttg catgaatgat ttgtgtttct ttatatTTTT
 agcctaaatg

3181 ttaactacat attgtctgga aacctgtcat tgtattcaga taattagatg
attatatatt

3241 gttgttactt tttcttgat tcatgaaaac tgtttttact aagttttcaa
atttgtaaag

5 3301 ttagcctttg aatgctagga atgcattatt gagggtcatt ctttattctt
tactattaata

3361 atatttttgga tgcaaaaaaaaa aaaaaaaaaa aaa

//

10

SSX-4

LOCUS NM_005636 576 bp mRNA linear
PRI 10-DEC-2001

15 DEFINITION Homo sapiens synovial sarcoma, X breakpoint 4 (SSX4),
mRNA.

ACCESSION NM_005636

VERSION NM_005636.1 GI:5032122

KEYWORDS .

SOURCE human.

20 ORGANISM Homo sapiens

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata;
Euteleostomi;

Mammalia; Eutheria; Primates; Catarrhini; Hominidae;
Homo.

25 REFERENCE 1 (bases 1 to 576)

AUTHORS Gure,A.O., Tureci,O., Sahin,U., Tsang,S.,
Scanlan,M.J., Knuth,A.,

Pfreundschuh,M., Old,L.J. and Chen,Y.T.

30 TITLE SSX: a multigene family with several members
transcribed in normal

testis and human cancer

JOURNAL Int. J. Cancer 72 (6), 965-971 (1997)

MEDLINE 98021352

35 COMMENT PROVISIONAL REFSEQ: This record has not yet been
subject to final

NCBI review. The reference sequence was derived from
U90841.1.

FEATURES Location/Qualifiers

source 1..576

```

/organism="Homo sapiens"
/db_xref="taxon:9606"
/chromosome="X"
/map="Xp11.3"
5      gene      1..576
        /gene="SSX4"
        /db_xref="LocusID:6759"
        /db_xref="MIM:300326"
        CDS      1..567
10      /gene="SSX4"
        /note="Kruppel-associated box containing SSX
gene"
        /codon_start=1
        /db_xref="LocusID:6759"
15      /db_xref="MIM:300326"
        /product="synovial sarcoma, X breakpoint 4"
        /protein_id="NP_005627.1"
        /db_xref="GI:5032123"

20      /translation="MNGDDAFARRPRDDAQISEKLRKAFDDIAKYFSKKEWEKMSSEKIVY
VYMKLNIEVMTKLGFKVTLPPFMRSKRAADFHGNDFGNDRNHRNQVERPQMTFG
SLQRIFPKIMPKKPAEEENGLKEVPEASGPQNDGKQLCPPGNPSTLEKINKTSGPKRG
KHAWTHRLRERKQLVVYEEISDPPEEDDE"
        misc_feature      70..246
25      /note="KRAB; Region: krueppel associated box"
BASE COUNT      187 a      127 c      150 g      112 t
ORIGIN
        1 atgaacggag acgacgcctt tgcaaggaga cccagggatg atgctcaa
atcagagaag
30      61 ttacgaaagg ccttcgatga tattgccaaa tacttctcta agaaagagt
ggaaaagatg
        121 aaatcctcgg agaaaatcgt ctatgtgtat atgaagctaa actatgag
catgactaaa
        181 ctaggtttca aggtcaccct cccacctttc atgcgtagta aacgggctg
agacttccac
35      241 gggaatgatt ttggtaacga tcgaaaccac aggaatcagg ttgaacgtcc
tcagatgact
        301 ttcggcagcc tccagagaat cttcccgaag atcatgccca agaagccagc
agaggaagaa

```


361, aatggtttga aggaagtgcc agaggcatct ggcccacaaa atgatgggaa
acagctgtgc
421 cccccgggaa atccaagtac cttggagaag attaacaaga catctggacc
caaaaggggg
5 481 aaacatgcct ggaccacag actgcgtgag agaaagcagc tgggtggttta
tgaagagatc
541 agcgaccctg aggaagatga cgagtaactc ccctcg

10 All patents and publications mentioned in the specification
are indicative of the levels of those skilled in the art to which
the invention pertains. All patents and publications are herein
incorporated by reference to the same extent as if each individual
publication was specifically and individually indicated to be
incorporated by reference.

15 The invention illustratively described herein suitably may be
practiced in the absence of any element or elements, limitation or
limitations which is not specifically disclosed herein. The terms
and expressions which have been employed are used as terms of
description and not of limitation, and there is no intention that
20 in the use of such terms and expressions indicates the exclusion
of equivalents of the features shown and described or portions
thereof. It is recognized that various modifications are possible
within the scope of the invention claimed. Thus, it should be
understood that although the present invention has been
25 specifically disclosed by preferred embodiments and optional
features, modification and variation of the concepts herein
disclosed may be resorted to by those skilled in the art, and that
such modifications and variations are considered to be within the
scope of this invention as defined by the appended claims.

30

WHAT IS CLAIMED IS:

1. An isolated epitope, comprising a component selected from the group consisting of:
 - (i) a polypeptide having the sequence as disclosed in TABLE 1;
 - (ii) an epitope cluster comprising the polypeptide of (i);
 - (iii) a polypeptide having substantial similarity to (i) or (ii);
 - (iv) a polypeptide having functional similarity to any of (i) through (iii); and
 - (v) a nucleic acid encoding the polypeptide of any of (i) through (iv).
2. The epitope of claim 1, wherein the epitope is immunologically active.
3. The epitope of claim 1, wherein the polypeptide is less than about 30 amino acids in length.
4. The epitope of claim 1, wherein the polypeptide is 8 to 10 amino acids in length.
5. The epitope of claim 1, wherein the substantial or functional similarity comprises addition of at least one amino acid.
6. The epitope of claim 5, wherein the at least one additional amino acid is at an N-terminus of the polypeptide.
7. The epitope of claim 1, wherein the substantial or functional similarity comprises a substitution of at least one amino acid.
8. The epitope of claim 1, the polypeptide having affinity to an HLA-A2 molecule.
9. The epitope of claim 8, wherein the affinity is determined by an assay of binding.
10. The epitope of claim 8, wherein the affinity is determined by an assay of restriction of epitope recognition.
11. The epitope of claim 8, wherein the affinity is determined by a prediction algorithm.
12. The epitope of claim 1, the polypeptide having affinity to an HLA-B7 or HLA-B51 molecule.
13. The epitope of claim 1, wherein the polypeptide is a housekeeping epitope.
14. The epitope of claim 1, wherein the polypeptide corresponds to an epitope displayed on a tumor cell.
15. The epitope of claim 1, wherein the polypeptide corresponds to an epitope displayed on a neovasculature cell.
16. The epitope of claim 1, wherein the peptide is an immune epitope.
17. The epitope of claim 1 wherein the epitope is a nucleic acid.
18. A pharmaceutical composition comprising the peptide of claim 1 and a pharmaceutically acceptable adjuvant, carrier, diluent, or excipient.
19. The composition of claim 18, where the adjuvant is a polynucleotide.

20. The composition of claim 19 wherein the polynucleotide comprises a dinucleotide.
21. The composition of claim 20 wherein the dinucleotide is CpG.
22. The composition of claim 18, wherein the adjuvant is encoded by a polynucleotide.
23. The composition of claim 18 wherein the adjuvant is a cytokine.
- 5 24. The composition of claim 23 wherein the cytokine is GM-CSF.
25. The composition of claim 18 further comprising a professional antigen-presenting cell (pAPC).
26. The composition of claim 25, wherein the pAPC is a dendritic cell.
27. The composition of claim 18, further comprising a second epitope.
- 10 28. The composition of claim 27, wherein the second epitope is a polypeptide.
29. The composition of claim 27, wherein the second epitope is a nucleic acid.
30. The composition of claim 27, wherein the second epitope is a housekeeping epitope.
31. The composition of claim 27, wherein the second epitope is an immune epitope.
- 15 32. A pharmaceutical composition comprising the nucleic acid of claim 1 and a pharmaceutically acceptable adjuvant, carrier, diluent, or excipient.
33. A recombinant construct comprising the nucleic acid of Claim 1.
34. The construct of claim 33, further comprising a plasmid, a viral vector, or an artificial chromosome.
- 20 35. The construct of claim 33, further comprising a sequence encoding at least one feature selected from the group consisting of a second epitope, an IRES, an ISS, an NIS, and ubiquitin.
36. A purified antibody that specifically binds to the epitope of claim 1.
37. A purified antibody that specifically binds to a peptide-MHC protein complex
- 25 comprising the epitope of claim 1.
38. The antibody of claim 36 or claim 37, wherein the antibody is a monoclonal antibody.
39. A multimeric MHC-peptide complex comprising the epitope of claim 1.
40. An isolated T cell expressing a T cell receptor specific for an MHC-peptide
- 30 complex, the complex comprising the epitope of claim 1.
41. The T cell of claim 40, produced by an *in vitro* immunization.
42. The T cell of claim 40, isolated from an immunized animal.
43. A T cell clone comprising the T cell of claim 40.
44. A polyclonal population of T cells comprising the T cell of claim 40.
- 35 45. A pharmaceutical composition comprising the T cell of claim 40 and a pharmaceutically acceptable adjuvant, carrier, diluent, or excipient.

46. An isolated protein molecule comprising the binding domain of a T cell receptor specific for an MHC-peptide complex, the complex comprising the epitope of claim 1.
47. The protein of claim 46, wherein the protein is multivalent.
48. An isolated nucleic acid encoding the protein of claim 46.
- 5 49. A recombinant construct comprising the nucleic acid of claim 48.
50. A host cell expressing the recombinant construct of claim 33 or 49.
51. The host cell of claim 50, wherein the host cell is a dendritic cell, macrophage, tumor cell, or tumor-derived cell.
52. The host cell of claim 50, wherein the host cell is a bacterium, fungus, or
10 protozoan.
53. A pharmaceutical composition comprising the host cell of claim 50 and a pharmaceutically acceptable adjuvant, carrier, diluent, or excipient.
54. A vaccine or immunotherapeutic composition comprising at least one component selected from the group consisting of the epitope of claim 1; the composition of claim 18, 32, 45, or
15 53; the construct of claim 33; the T cell of claim 40, and the host cell of claim 50.
55. A method of treating an animal, comprising:
administering to an animal the vaccine or immunotherapeutic composition of claim
54.
56. The method of claim 55, wherein the administering step comprises a mode of
20 delivery selected from the group consisting of transdermal, intranodal, perinodal, oral, intravenous, intradermal, intramuscular, intraperitoneal, mucosal, aerosol inhalation, and instillation.
57. The method of claim 55, further comprising a step of assaying to determine a characteristic indicative of a state of a target cell or target cells.
58. The method of claim 57, comprising a first assaying step and a second assaying
25 step, wherein the first assaying step precedes the administering step, and wherein the second assaying step follows the administering step.
59. The method of claim 58, further comprising a step of comparing the characteristic determined in the first assaying step with the characteristic determined in the second assaying step to obtain a result.
- 30 60. The method of claim 59, wherein the result is selected from the group consisting of: evidence of an immune response, a diminution in number of target cells, a loss of mass or size of a tumor comprising target cells, a decrease in number or concentration of an intracellular parasite infecting target cells.
61. A method of evaluating immunogenicity of a vaccine or immunotherapeutic
35 composition, comprising:

- administering to an animal the vaccine or immunotherapeutic composition of claim 54; and
- evaluating immunogenicity based on a characteristic of the animal.
62. The method of claim 61, wherein the animal is HLA-transgenic.
- 5 63. A method of evaluating immunogenicity, comprising:
- in vitro* stimulation of a T cell with the vaccine or immunotherapeutic composition of claim 54; and
- evaluating immunogenicity based on a characteristic of the T cell.
64. The method of claim 63, wherein the stimulation is a primary stimulation.
- 10 65. A method of making a passive/adoptive immunotherapeutic, comprising:
- combining the T cell of claim 40 or the host cell of claim 50 with a pharmaceutically acceptable adjuvant, carrier, diluent, or excipient.
66. A method of determining specific T cell frequency comprising the step of contacting T cells with a MHC-peptide complex comprising the epitope of claim 1.
- 15 67. The method of claim 66, wherein the contacting step comprises at least one feature selected from the group consisting of immunization, restimulation, detection, and enumeration.
68. The method of Claim 66, further comprising ELISPOT analysis, limiting dilution analysis, flow cytometry, *in situ* hybridization, the polymerase chain reaction or any combination thereof.
- 20 69. A method of evaluating immunologic response, comprising the method of claim 66 carried out prior to and subsequent to an immunization step.
70. A method of evaluating immunologic response, comprising:
- determining frequency, cytokine production, or cytolytic activity of T cells, prior to and subsequent to a step of stimulation with MHC-peptide complexes comprising the epitope of claim 1.
- 25 71. A method of diagnosing a disease comprising:
- contacting a subject tissue with at least one component selected from the group consisting of the T cell of claim 40, the host cell of claim 50, the antibody of claim 36, the protein of claim 46; and
- 30 diagnosing the disease based on a characteristic of the tissue or of the component.
72. The method of claim 71, wherein the contacting step takes place *in vivo*.
73. The method of claim 71, wherein the contacting step takes place *in vitro*.
74. A method of making a vaccine, comprising:
- 35 combining at least one component selected from the group consisting of the epitope of claim 1; the composition of claim 18, 32, 45, or 53; the construct of claim 33;

the T cell of claim 40, and the host cell of claim 50, with a pharmaceutically acceptable adjuvant, carrier, diluent, or excipient.

5 75. A computer readable medium having recorded thereon the sequence of any one of SEQ ID NOS: X -Y, in a machine having a hardware or software that calculates the physical, biochemical, immunologic, or molecular genetic properties of a molecule embodying said sequence.

 76. A method of treating an animal comprising combining the method of claim 55 combined with at least one mode of treatment selected from the group of radiation therapy, chemotherapy, biochemotherapy, and surgery.

10 77. An isolated polypeptide comprising an epitope cluster from a target-associated antigen having the sequence as disclosed in Tables 25-44, wherein the amino acid sequence consists of not more than about 80% of the amino acid sequence of the antigen.

 78. A vaccine or immunotherapeutic product comprising the polypeptide of claim 78.

 79. An isolated polynucleotide encoding the polypeptide of claim 78.

15 80. A vaccine or immunotherapeutic product comprising the polynucleotide of claim 80.

 81. The polynucleotide of claim 79 or 80, wherein the polynucleotide is DNA.

 82. The polynucleotide of claim 79 or 80, wherein the polynucleotide is RNA.

2/22

FIG. 1B

CTAG_HUMAN	NY-ESO	(101)	EAATAPSEINODPPEIVPGAKNTTSGNNTT	150
AAD05202 - CAG-3		(101)	EAATAPSEINODPPEIVPGAKNTTSGNNTT	
CAA11044 -LAGE-1a		(101)	EAATAPSEINODPPEIVPGAKNTTSGNNTT	
CAA10194 - LAGE-1s		(101)	EAATAPSEINODPPEIVPGAKNTTSGNNTT	
CAA11043 - LAGE-1b		(101)	EAATAPSEINODPPEIVPGAKNTTSGNNTT	
CAA10196 - LAGE-1L		(101)	EAATAPSEINODPPEIVPGAKNTTSGNNTT	
AAH02833 CT-2		(101)	EAATAPSEINODPPEIVPGAKNTTSGNNTT	
Consensus		(101)	EAEIVRRILSRDAAPLPRPGAVLKDFTVSGNLLFIRLTAADHRQLQLSIS	
CTAG_HUMAN	NY-ESO	(151)	VGWGLGASPEGQKARDLRTPKHKVGEPTPGPPPPPEGAQGDGCRGVA	200
AAD05202 - CAG-3		(151)	VGWGLGASPEGQKARDLRTPKHKVGEPTPGPPPPPEGAQGDGCRGVA	
CAA11044 -LAGE-1a		(151)	VGWGLGASPEGQKARDLRTPKHKVGEPTPGPPPPPEGAQGDGCRGVA	
CAA10194 - LAGE-1s		(151)	VGWGLGASPEGQKARDLRTPKHKVGEPTPGPPPPPEGAQGDGCRGVA	
CAA11043 - LAGE-1b		(151)	VGWGLGASPEGQKARDLRTPKHKVGEPTPGPPPPPEGAQGDGCRGVA	
CAA10196 - LAGE-1L		(151)	VGWGLGASPEGQKARDLRTPKHKVGEPTPGPPPPPEGAQGDGCRGVA	
AAH02833 CT-2		(151)	VGWGLGASPEGQKARDLRTPKHKVGEPTPGPPPPPEGAQGDGCRGVA	
Consensus		(151)	SCLQQLSLLMWITQCFLPVFLAQ PSGQR	

FIG. 1C

CTAG HUMAN	NY-ESO	(181)	-----
AAD05202	- CAG-3	(181)	-----
CAA11044	-LAGE-1a	(181)	-----
CAA10194	- LAGE-1s	(181)	-----
CAA11043	- LAGE-1b	(201)	FNVMSAPHI
CAA10196	- LAGE-1L	(201)	FNVMSAPHI
AAH02833	CT-2	(201)	FNVMSAPHI
	Consensus	(201)	

4/22





FIG. 2

5/22

FIG. 3A

**FACscan Analysis of Binding Assay to Determine the Binding
Ability of Tyrosinase 208-216 Peptide to MHC Class 1**




F1 (HLA A2 Peptide) = 3.13

F1 (TYR 208-216 Peptide) = 0.01

6/22

FIG. 3B


**FACscan Analysis of Binding Assay to Determine the Binding
Ability of Tyrosinase 207-215 Peptide to MHC Class 1**



F1 (HLA A2 Peptide) = 3.13
F1 (TYR 207-215 Peptide) = 2.00


7/22

FIG. 4



8/22


FIG. 5
Comparison of Peptides Binding Affinity to HLA A2




9/22

FIG. 6

**SSX2₄₁₋₄₉ specific lysis by CTL from peptide
injected HHD1 mice**




10/22



Pool sequencing of PSMA_163–192 Digested for 60 min by proteasome


11/22

FIG. 7B



Pool sequencing of PSMA_163-192 Digested for 60 min by proteasome

12/22




Pool sequencing of PSMA_163–192 Digested for 60 min by proteasome


FIG. 7C

13/22

FIG. 8




281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310
R G I A E A V G L P S I P V H P I G Y Y D A Q K L L E K M G



Pool sequencing of PSMA_281_310 Digested for 60 min by Proteasome


FIG. 9

FIG. 10
Comparison of Peptides Binding
Affinity to HLA A2
by Binding Assay




16/22

Autologous DC Present A1 Peptide to CD8 T cell

*FIG. 11*


17/22

Secretion of IFN γ Was Blocked by Anti-A1 Antibody

*FIG. 12*


18/22

FIG. 13
Comparison of Peptides Binding Affinity
to HLA A2 by Binding Assay




19/22

FIG. 14
Comparison of Peptides Binding Affinity
to HLA A2 by Binding Assay




20/22



Graphs show lysis of unpulsed EL4 cells (open circles) and EL4 cells pulsed with gp33 peptide (solid triangles). Symbols represent individual mice and one of three similar experiments is shown.

FIG. 15

21/22



Mean tumor volumes \pm 1SD are shown for mice immunized with pEFGPL33A DNA (solid circles) or control pEGFP-N3 DNA (open triangles). Numbers in brackets indicate number of mice with tumors/total number of mice in group. One of two similar experiments is shown.

FIG. 16

22/22




FIG. 17

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.